

Full wwPDB NMR Structure Validation Report (i)

May 28, 2020 – 07:21 pm BST

PDB ID : 1BWY

Title : NMR STUDY OF BOVINE HEART FATTY ACID BINDING PROTEIN Authors : Lassen, D.; Luecke, C.; Kveder, M.; Mesgarzadeh, A.; Schmidt, J.M.; Specht,

B.; Lezius, A.; Spener, F.; Rueterjans, H.

Deposited on : 1998-09-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)

NmrClust : Kelley et al. (1996)

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	${ m NMR~archive} \ (\#{ m Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	A	132	86%	14%		



2 Ensemble composition and analysis (i)

This entry contains 25 models. Model 5 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid				
1 A:1-A:132 (132)		0.80	5	

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 4 single-model clusters were found.

Cluster number	$egin{array}{c} \mathbf{Models} \end{array}$
1	3, 5, 6, 8, 10, 11, 14, 16, 17, 20
2	1, 15, 18, 21, 22
3	9, 24, 25
4	2, 7, 19
Single-model clusters	4; 12; 13; 23



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2076 atoms, of which 1047 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called PROTEIN (HEART FATTY ACID BINDING PROTEIN).

Mol	Chain	Residues	Atoms				Trace		
1	Λ	132	Total	С	Н	N	О	S	0
1	A	152	2076	649	1047	173	203	4	U




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)



4.2.2 Score per residue for model 2






4.2.3 Score per residue for model 3

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.4 Score per residue for model 4

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.5 Score per residue for model 5 (medoid)

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)



4.2.6 Score per residue for model 6


• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)



4.2.7 Score per residue for model 7









4.2.8 Score per residue for model 8

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.9 Score per residue for model 9

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.10 Score per residue for model 10

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.11 Score per residue for model 11






4.2.12 Score per residue for model 12

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.13 Score per residue for model 13

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.14 Score per residue for model 14

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)



4.2.15 Score per residue for model 15

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)



4.2.16 Score per residue for model 16








4.2.17 Score per residue for model 17

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.18 Score per residue for model 18

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.19 Score per residue for model 19

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.20 Score per residue for model 20

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.21 Score per residue for model 21






4.2.22 Score per residue for model 22

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.23 Score per residue for model 23

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)




4.2.24 Score per residue for model 24

• Molecule 1: PROTEIN (HEART FATTY ACID BINDING PROTEIN)



4.2.25 Score per residue for model 25





5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: DISTANCE GEOMETRY, ENERGY MIN-IMIZATION.

Of the 100 calculated structures, 25 were deposited, based on the following criterion: LOWEST VIOLATION OF EXPERIMENTAL DISTANCE CONSTRAINTS.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
SYBYL	refinement	6.0
DIANA	structure solution	
SYBYL	structure solution	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1029	1047	1045	1±1
All	All	25725	26175	26125	27

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$\mathbf{Distance}(\mathbf{\mathring{A}})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:18:ASP:OD2	1:A:124:CYS:SG	0.49	2.71	20	3
1:A:10:LEU:HD23	1:A:11:VAL:N	0.49	2.23	15	1
1:A:31:GLN:NE2	1:A:32:VAL:HG23	0.47	2.25	19	1
1:A:106:ARG:CG	1:A:115:LEU:HD23	0.45	2.41	10	1
1:A:8:TRP:CD2	1:A:42:ILE:HD13	0.45	2.46	7	1
1:A:19:TYR:CD1	1:A:124:CYS:SG	0.45	3.09	9	1
1:A:12:ASP:HB3	1:A:127:THR:HG22	0.44	1.88	13	1
1:A:9:LYS:NZ	1:A:129:GLU:OE2	0.44	2.51	4	1



Continued from previous page...

Atom-1	Atom-2	Clock (Å)	$\mathbf{Distance}(\mathbf{\mathring{A}})$	${f Models}$	
Atom-1	Atom-2	$\operatorname{Clash}(\mathrm{\AA})$	Distance(A)	Worst	Total
1:A:9:LYS:NZ	1:A:129:GLU:OE1	0.44	2.48	1	1
1:A:64:PHE:CD2	1:A:84:VAL:HG11	0.43	2.48	12	1
1:A:8:TRP:CG	1:A:42:ILE:HD13	0.43	2.49	7	1
1:A:19:TYR:CE2	1:A:117:LEU:HD12	0.43	2.49	17	1
1:A:112:LYS:NZ	1:A:129:GLU:OE2	0.42	2.51	13	1
1:A:18:ASP:OD1	1:A:21:LYS:NZ	0.42	2.52	7	2
1:A:12:ASP:OD2	1:A:14:LYS:NZ	0.42	2.52	4	1
1:A:113:LEU:C	1:A:113:LEU:HD12	0.42	2.35	24	1
1:A:91:LEU:HD22	1:A:91:LEU:C	0.41	2.35	13	1
1:A:71:ASP:OD2	1:A:79:LYS:NZ	0.41	2.54	4	1
1:A:90:LYS:NZ	1:A:107:GLU:OE1	0.41	2.54	18	1
1:A:8:TRP:CZ3	1:A:113:LEU:HB2	0.41	2.51	19	1
1:A:19:TYR:CE2	1:A:117:LEU:HD21	0.41	2.51	22	1
1:A:86:LEU:C	1:A:86:LEU:HD12	0.40	2.37	6	1
1:A:87:ASP:OD2	1:A:90:LYS:NZ	0.40	2.54	8	1
1:A:91:LEU:HD23	1:A:91:LEU:C	0.40	2.37	7	1

5.2 Torsion angles (i)

5.2.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	${f Analysed}$	Favoured	Allowed	${f Outliers}$	Percentiles	
1	A	130/132 (98%)	98±5 (75±4%)	22±5 (17±4%)	$10\pm 2 \ (8\pm 2\%)$	2 15	
All	All	$3250/3300 \ (98\%)$	2451 (75%)	553 (17%)	246 (8%)	2 15	

All 42 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	${f Res}$	Type	Models (Total)
1	A	22	SER	22
1	A	46	GLY	19
1	A	3	ALA	17
1	A	98	ASN	15
1	A	5	VAL	14
1	A	78	ARG	14



Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	67	GLY	13
1	A	88	GLY	11
1	A	25	VAL	11
1	A	34	ASN	8
1	A	26	GLY	7
1	A	13	SER	7
1	A	121	THR	7
1	A	56	THR	7
1	A	66	LEU	7
1	A	2	ASP	6
1	A	28	ALA	5
1	A	76	ASP	5
1	A	37	LYS	5
1	A	120	GLY	5
1	A	27	PHE	4
1	A	17	ASP	4
1	A	57	PHE	4
1	A	11	VAL	3
1	A	36	THR	3
1	A	90	LYS	3
1	A	110	ASP	2
1	A	45	ASN	2
1	A	89	GLY	2
1	A	29	THR	2
1	A	18	ASP	1
1	A	75	ALA	1
1	A	122	ALA	1
1	A	59	ASN	1
1	A	73	THR	1
1	A	68	VAL	1
1	A	55	SER	1
1	A	12	ASP	1
1	A	23	LEU	1
1	A	33	GLY	1
1	A	54	GLN	1
1	A	111	GLY	1

5.2.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	116/116 (100%)	100±3 (86±2%)	16±3 (14±2%)	7 47
All	All	2900/2900 (100%)	2505 (86%)	395 (14%)	7 47

All 83 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	66	LEU	25
1	A	100	GLN	19
1	A	48	THR	19
1	A	117	LEU	13
1	A	127	THR	11
1	A	20	MET	11
1	A	16	PHE	9
1	A	60	THR	9
1	A	87	ASP	9
1	A	83	ILE	9
1	A	51	ILE	8
1	A	90	LYS	8
1	A	9	LYS	8
1	A	68	VAL	8
1	A	105	VAL	7
1	A	30	ARG	7
1	A	25	VAL	7
1	A	53	THR	6
1	A	97	TRP	6
1	A	14	LYS	6
1	A	65	LYS	6
1	A	12	ASP	6
1	A	35	MET	6
1	A	78	ARG	6
1	A	131	GLN	6
1	A	27	PHE	5
1	A	37	LYS	5
1	A	121	THR	5
1	A	91	LEU	5
1	A	104	LEU	5
1	A	58	LYS	5
1	A	19	TYR	4
1	A	15	ASN	4
1	A	21	LYS	4
1	A	77	ASP	4
1	A	18	ASP	4



Continued from previous page...

Mol	nued fron Chain	Res	$\overline{ ext{Type}}$	Models (Total)
1	A	39	THR	4
1	A	54	GLN	4
1	A	106	ARG	4
1	A	47	ASP	4
1	A	124	CYS	4
1	A	59	ASN	4
1	A	50	ILE	3
1	A	103	SER	3
1	A	113	LEU	3
1	A	98	ASN	3
1	A	69	GLU	3
1	A	74	THR	3
1	A	61	GLU	3
1	A	102	THR	3
1	A	85	THR	3
1	A	101	GLU	3
1	A	23	LEU	3
1	A	82	SER	3
1	A	32	VAL	3
1	A	119	HIS	3
1	A	41	ILE	3
1	A	40	THR	2
1	A	108	MET	2
1	A	84	VAL	2
1	A	114	ILE	2
1	A	31	GLN	2
1	A	81	LYS	2
1	A	86	LEU	2
1	A	118	THR	2
1	A	125	THR	2
1	A	57	PHE	2
1	A	52	LYS	2
1	A	115	LEU	2
1	A	126	ARG	2
1	A	34	ASN	2
1	A	17	ASP	2
1	A	130	LYS	1
1	A	76	ASP	1
1	A	73	THR	1
1	A	96	LYS	1
1	A	8	TRP	1
1	A	56	THR	1



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Models (Total)
1	A	62	ILE	1
1	A	63	SER	1
1	A	29	THR	1
1	A	79	LYS	1
1	A	43	GLU	1

5.2.3 RNA (i)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates (i)

There are no carbohydrates in this entry.

5.5 Ligand geometry (i)

There are no ligands in this entry.

5.6 Other polymers (i)

There are no such molecules in this entry.

5.7 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Chemical shift validation (i)

No chemical shift data were provided

