

Full wwPDB X-ray Structure Validation Report (i)

Jan 27, 2024 - 06:40 PM EST

PDB ID	:	1CLC
Title	:	THREE-DIMENSIONAL STRUCTURE OF ENDOGLUCANASE D AT 1.9
		ANGSTROMS RESOLUTION
Authors	:	Alzari, P.M.; Lascombe, M.B.
Deposited on		
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	А	639	73%	10%	•	15%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ENDOGLUCANASE CELD; EC: 3.2.1.4.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	541	Total 4213	C 2685	N 702	O 808	S 18	0	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	13	ILE	THR	conflict	UNP P04954
А	14	THR	LEU	conflict	UNP P04954
А	15	ASN	LYS	conflict	UNP P04954
А	?	-	SER	deletion	UNP P04954
А	?	-	MET	deletion	UNP P04954
А	?	-	LYS	deletion	UNP P04954
А	?	-	LYS	deletion	UNP P04954
А	?	-	VAL	deletion	UNP P04954
А	18	GLY	LEU	conflict	UNP P04954
А	?	-	LEU	deletion	UNP P04954
А	?	-	LEU	deletion	UNP P04954
А	?	-	ILE	deletion	UNP P04954
А	?	-	ALA	deletion	UNP P04954
А	?	-	VAL	deletion	UNP P04954
А	21	ASP	PHE	conflict	UNP P04954
А	23	GLN	-	insertion	UNP P04954
А	24	PRO	-	insertion	UNP P04954

There are 17 discrepancies between the modelled and reference sequences:

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	3	Total Ca 3 3	0	0

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).



Mol	Chain	Residues	Atom	ns	ZeroOcc	AltConf
3	А	1	Total 1	Zn 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	369	Total O 369 369	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- MM
 MM
 MM
 MM
 MM
 MM

 VLIN
 VIL
 VIL
 VIL
 VIL
 VIL
 VIL

 VLIN
 VLIN
 VIL
 VIL
 VIL
 VIL
 VIL
 VIL

 VLIN
 VLIN
 VIL
 VI
- Molecule 1: ENDOGLUCANASE CELD; EC: 3.2.1.4



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 31 2 1	Depositor	
Cell constants	98.90Å 98.90Å 191.40Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor	
Resolution (Å)	7.00 - 1.90	Depositor	
% Data completeness	(Not available) (7.00-1.90)	Depositor	
(in resolution range)	(100 available) (1.00-1.50)	Depositor	
R_{merge}	0.07	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	X-PLOR	Depositor	
R, R_{free}	0.204 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	4586	wwPDB-VP	
Average B, all atoms $(Å^2)$	34.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain		lengths	Bond angles		
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.50	0/4330	0.71	0/5904	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	1	0

There are no bond length outliers.

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	А	166	ILE	CB

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4213	0	3931	43	0

Continued on next page...



001000	continuou front proto de pagom						
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes	
2	А	3	0	0	0	0	
3	А	1	0	0	0	0	
4	А	369	0	0	5	0	
All	All	4586	0	3931	43	0	

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:64:ASN:H	1:A:114:ASN:ND2	1.90	0.69
1:A:532:TYR:HA	1:A:561:ASN:HD21	1.59	0.67
1:A:76:SER:O	1:A:92:THR:HA	1.95	0.65
1:A:177:ASP:O	4:A:939:HOH:O	2.14	0.65
1:A:445:HIS:CD2	1:A:447:TYR:H	2.16	0.64
1:A:445:HIS:HD2	1:A:447:TYR:H	1.45	0.64
1:A:180:LEU:HD23	1:A:188:THR:HG23	1.79	0.64
1:A:471:GLN:HE21	1:A:471:GLN:HA	1.64	0.61
1:A:137:ASN:HD22	1:A:137:ASN:H	1.49	0.61
1:A:538:TRP:HB2	1:A:539:PRO:HA	1.84	0.59
1:A:479:ASN:ND2	1:A:481:ASP:H	1.98	0.59
1:A:224:LYS:HE3	4:A:1016:HOH:O	2.03	0.58
1:A:376:ASP:HB2	1:A:379:TYR:CD2	2.40	0.57
1:A:166:ILE:HG22	1:A:168:TYR:CE2	2.40	0.56
1:A:384:GLU:HG2	1:A:423:LEU:HD23	1.89	0.55
1:A:55:ARG:HG2	1:A:500:ASN:HB2	1.91	0.53
1:A:256:LEU:HD13	1:A:323:LYS:HE3	1.92	0.52
1:A:275:ASN:HB3	4:A:845:HOH:O	2.11	0.51
1:A:402:ASN:HB3	4:A:915:HOH:O	2.10	0.50
1:A:180:LEU:HD21	1:A:190:LYS:HB2	1.94	0.49
1:A:492:HIS:HE1	4:A:659:HOH:O	1.94	0.49
1:A:479:ASN:HD22	1:A:480:ASN:N	2.11	0.48
1:A:212:THR:HG23	1:A:562:ALA:HB2	1.95	0.48
1:A:479:ASN:HD22	1:A:479:ASN:C	2.16	0.48
1:A:533:LEU:H	1:A:561:ASN:ND2	2.12	0.47
1:A:77:THR:O	1:A:124:PRO:HD2	2.14	0.47
1:A:275:ASN:HA	1:A:350:SER:OG	2.15	0.47
1:A:181:ASP:OD1	1:A:187:HIS:HA	2.16	0.46
1:A:479:ASN:HD22	1:A:481:ASP:H	1.63	0.46
1:A:474:ASN:HD22	1:A:479:ASN:H	1.64	0.45

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:474:ASN:ND2	1:A:479:ASN:H	2.14	0.45
1:A:156:GLY:HA2	1:A:173:CYS:O	2.16	0.45
1:A:193:THR:O	1:A:194:LYS:HB2	2.17	0.45
1:A:533:LEU:H	1:A:561:ASN:HD21	1.63	0.45
1:A:295:SER:HB3	1:A:344:ALA:HB2	2.00	0.43
1:A:64:ASN:H	1:A:114:ASN:HD21	1.64	0.43
1:A:53:ARG:HD2	1:A:73:ALA:HB2	2.00	0.43
1:A:120:TYR:HB3	1:A:131:ASN:HD22	1.83	0.42
1:A:368:ALA:HB2	1:A:383:PHE:CD1	2.55	0.41
1:A:63:PRO:O	1:A:64:ASN:HB2	2.20	0.41
1:A:300:ALA:HA	1:A:335:LEU:HD11	2.02	0.41
1:A:73:ALA:O	1:A:106:TYR:CD1	2.74	0.40
1:A:371:TRP:CD1	1:A:380:LEU:HB2	2.56	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	539/639~(84%)	522~(97%)	16 (3%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	199	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mo	l Cha	ain	Analysed	Rotameric	Outliers	Percentiles
1	A	L	432/539~(80%)	415 (96%)	17~(4%)	32 23

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	45	THR
1	А	56	LEU
1	А	96	MET
1	А	137	ASN
1	А	181	ASP
1	А	189	LYS
1	А	211	ILE
1	А	227	LEU
1	А	237	LYS
1	А	259	GLN
1	А	471	GLN
1	А	479	ASN
1	А	480	ASN
1	А	487	LEU
1	А	493	VAL
1	А	510	ASN
1	А	561	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	114	ASN
1	А	131	ASN
1	А	137	ASN
1	А	176	ASN
1	А	259	GLN
1	А	326	ASN
1	А	346	GLN
1	А	443	GLN
1	А	445	HIS
1	А	460	ASN
1	А	466	GLN
1	А	471	GLN
1	А	474	ASN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	А	479	ASN
1	А	480	ASN
1	А	492	HIS
1	А	510	ASN
1	А	561	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

