



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 10:41 am BST

PDB ID : 1F3U
Title : CRYSTAL STRUCTURE OF THE RAP30/74 INTERACTION DOMAINS OF HUMAN TFIIF
Authors : Gaiser, F.; Tan, S.; Richmond, T.J.
Deposited on : 2000-06-06
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	118	
1	C	118	
1	E	118	
1	G	118	
2	B	171	
2	D	171	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	171	
2	H	171	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9192 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTION INITIATION FACTOR IIF, BETA SUB-UNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	118	895	557	163	174	1	16	0	0
1	C	118	894	557	163	173	1	21	0	0
1	E	118	895	557	163	174	1	0	0	0
1	G	118	895	557	163	174	1	5	0	0

- Molecule 2 is a protein called TRANSCRIPTION INITIATION FACTOR IIF, ALPHA SUB-UNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	139	1158	737	213	205	3	22	0	0
2	D	153	1284	812	235	234	3	14	0	0
2	F	154	1310	827	240	240	3	90	8	0
2	H	137	1169	743	215	208	3	52	6	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	95	Total	O	0	0
			95	95		
3	B	89	Total	O	0	0
			89	89		
3	C	52	Total	O	0	0
			52	52		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	91	Total 91	O 91	0	0
3	E	118	Total 118	O 118	0	0
3	F	119	Total 119	O 119	0	0
3	G	71	Total 71	O 71	0	0
3	H	57	Total 57	O 57	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRANSCRIPTION INITIATION FACTOR IIF, BETA SUBUNIT





- Molecule 1: TRANSCRIPTION INITIATION FACTOR IIF, BETA SUBUNIT



- Molecule 1: TRANSCRIPTION INITIATION FACTOR IIF, BETA SUBUNIT




- Molecule 1: TRANSCRIPTION INITIATION FACTOR IIF, BETA SUBUNIT




- Molecule 2: TRANSCRIPTION INITIATION FACTOR IIF, ALPHA SUBUNIT







● Molecule 2: TRANSCRIPTION INITIATION FACTOR IIF, ALPHA SUBUNIT



● Molecule 2: TRANSCRIPTION INITIATION FACTOR IIF, ALPHA SUBUNIT



● Molecule 2: TRANSCRIPTION INITIATION FACTOR IIF, ALPHA SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	48.02Å 72.29Å 82.27Å 104.51° 93.32° 104.32°	Depositor
Resolution (Å)	6.00 – 1.70 67.37 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.4 (6.00-1.70) 96.7 (67.37-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.70Å)	Xtrriage
Refinement program	CNS 0.5	Depositor
R, R_{free}	0.225 , 0.260 0.228 , 0.261	Depositor DCC
R_{free} test set	4381 reflections (4.02%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtrriage
Anisotropy	0.145	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 71.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9192	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1801e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/909	0.65	0/1227
1	C	0.30	0/908	0.67	0/1226
1	E	0.38	0/909	0.71	1/1227 (0.1%)
1	G	0.33	0/909	0.63	0/1227
2	B	0.35	0/1186	0.68	0/1594
2	D	0.31	0/1314	0.60	0/1768
2	F	0.38	0/1340	0.65	0/1804
2	H	0.34	0/1196	0.61	0/1608
All	All	0.34	0/8671	0.65	1/11681 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	38	GLY	N-CA-C	5.22	126.16	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	895	0	904	25	0
1	C	894	0	901	33	0
1	E	895	0	904	29	0
1	G	895	0	904	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1158	0	1151	35	0
2	D	1284	0	1261	42	0
2	F	1310	0	1283	34	0
2	H	1169	0	1162	47	0
3	A	95	0	0	1	0
3	B	89	0	0	7	0
3	C	52	0	0	2	0
3	D	91	0	0	2	0
3	E	118	0	0	4	0
3	F	119	0	0	2	0
3	G	71	0	0	2	0
3	H	57	0	0	1	0
All	All	9192	0	8470	236	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 236 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:56:ILE:HD11	2:D:161:GLU:HG2	1.52	0.88
1:A:59:ASN:HD21	1:A:62:LEU:HD13	1.40	0.86
2:B:51:LEU:HD22	2:B:91:PHE:HZ	1.43	0.83
2:D:21:ASN:HD22	2:D:21:ASN:C	1.84	0.80
1:G:59:ASN:HD22	1:G:59:ASN:C	1.87	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/118 (98%)	105 (90%)	6 (5%)	5 (4%)	2	0
1	C	116/118 (98%)	103 (89%)	7 (6%)	6 (5%)	2	0
1	E	116/118 (98%)	114 (98%)	2 (2%)	0	100	100
1	G	116/118 (98%)	109 (94%)	7 (6%)	0	100	100
2	B	135/171 (79%)	128 (95%)	5 (4%)	2 (2%)	10	2
2	D	149/171 (87%)	139 (93%)	8 (5%)	2 (1%)	12	2
2	F	153/171 (90%)	140 (92%)	7 (5%)	6 (4%)	3	0
2	H	136/171 (80%)	121 (89%)	10 (7%)	5 (4%)	3	0
All	All	1037/1156 (90%)	959 (92%)	52 (5%)	26 (2%)	5	1

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	PRO
1	A	76	SER
2	B	119	THR
1	C	102	ASP
2	F	77	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/96 (100%)	94 (98%)	2 (2%)	53	36
1	C	95/96 (99%)	94 (99%)	1 (1%)	73	63
1	E	96/96 (100%)	92 (96%)	4 (4%)	30	12
1	G	96/96 (100%)	92 (96%)	4 (4%)	30	12
2	B	122/147 (83%)	117 (96%)	5 (4%)	30	12
2	D	135/147 (92%)	131 (97%)	4 (3%)	41	22
2	F	138/147 (94%)	130 (94%)	8 (6%)	20	6
2	H	124/147 (84%)	118 (95%)	6 (5%)	25	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	902/972 (93%)	868 (96%)	34 (4%)	33 14

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	48	THR
2	F	94	GLU
2	H	91	PHE
2	F	55	LYS
2	B	95	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	21	ASN
2	D	53	ASN
2	H	121	ASN
2	D	38	ASN
2	D	44	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	118/118 (100%)	0.72	10 (8%) 10 12	22, 39, 69, 80	6 (5%)
1	C	118/118 (100%)	1.22	27 (22%) 0 0	29, 52, 76, 85	8 (6%)
1	E	118/118 (100%)	0.63	7 (5%) 22 24	20, 32, 58, 75	0
1	G	118/118 (100%)	1.04	16 (13%) 3 3	21, 42, 69, 77	1 (0%)
2	B	135/171 (78%)	0.84	14 (10%) 6 7	21, 42, 68, 82	0
2	D	151/171 (88%)	1.02	26 (17%) 1 1	27, 49, 72, 79	0
2	F	145/171 (84%)	1.03	22 (15%) 2 2	19, 38, 70, 78	10 (6%)
2	H	133/171 (77%)	1.39	31 (23%) 0 0	25, 48, 76, 83	10 (7%)
All	All	1036/1156 (89%)	0.99	153 (14%) 2 2	19, 42, 73, 85	35 (3%)

The worst 5 of 153 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	55	LYS	10.3
2	D	91	PHE	10.2
2	F	164	TRP	10.2
1	G	75	VAL	8.6
1	C	73	ALA	8.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.