



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 03:16 PM EDT

PDB ID : 1FY1
Title : [R23S,F25E]HBP, A MUTANT OF HUMAN HEPARIN BINDING PROTEIN (CAP37)
Authors : Kastrup, J.S.; Linde, V.; Pedersen, A.K.; Stoffer, B.; Iversen, L.F.; Larsen, I.K.; Rasmussen, P.B.; Flodgaard, H.J.; Bjorn, S.E.
Deposited on : 2000-09-28
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	
2	B	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	B	2	X	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 1809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEPARIN-BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	225	1697	1045	324	317	11	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	SER	ARG	engineered mutation	UNP P20160
A	25	GLU	PHE	engineered mutation	UNP P20160

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0

- Molecule 4 is ETHANOL (three-letter code: EOH) (formula: C_2H_6O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	3	2	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	53	Total 53	O 53	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HEPARIN-BINDING PROTEIN



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	38.72Å 66.56Å 102.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 19.68 – 2.52	Depositor EDS
% Data completeness (in resolution range)	98.2 (20.00-2.50) 97.8 (19.68-2.52)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.55 (at 2.53Å)	Xtrriage
Refinement program	TNT	Depositor
R, R_{free}	0.207 , 0.294 0.212 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	37.9	Xtrriage
Anisotropy	0.415	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 82.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1809	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EOH, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	6/1731 (0.3%)	1.49	27/2346 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	118	GLU	CD-OE2	6.21	1.32	1.25
1	A	151	GLU	CD-OE2	6.21	1.32	1.25
1	A	25	GLU	CD-OE2	6.19	1.32	1.25
1	A	98	GLU	CD-OE2	5.77	1.31	1.25
1	A	66	GLN	CG-CD	5.72	1.64	1.51
1	A	182	GLU	CD-OE1	5.29	1.31	1.25

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	199	GLY	C-N-CD	-23.07	69.85	120.60
1	A	47	ASN	CB-CA-C	11.85	134.09	110.40
1	A	46	GLN	CA-C-N	-8.65	98.18	117.20
1	A	59	ASP	CB-CG-OD2	-7.14	111.87	118.30
1	A	142	ARG	NE-CZ-NH2	-7.05	116.77	120.30
1	A	59	ASP	CB-CG-OD1	6.90	124.51	118.30
1	A	82	ASP	CB-CG-OD2	-6.81	112.17	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	166	ARG	N-CA-CB	6.72	122.71	110.60
1	A	152	ASP	CB-CG-OD2	-6.60	112.36	118.30
1	A	47	ASN	C-N-CD	-6.58	106.11	120.60
1	A	96	ASP	CB-CG-OD2	-6.38	112.55	118.30
1	A	96	ASP	CB-CG-OD1	6.38	124.04	118.30
1	A	193	LEU	CB-CA-C	6.29	122.14	110.20
1	A	142	ARG	NE-CZ-NH1	6.12	123.36	120.30
1	A	200	PRO	N-CA-CB	6.01	110.52	103.30
1	A	89	ASP	CB-CG-OD2	-5.81	113.07	118.30
1	A	46	GLN	CB-CA-C	5.80	122.00	110.40
1	A	155	ARG	NE-CZ-NH1	5.72	123.16	120.30
1	A	214	ASP	CB-CG-OD2	-5.69	113.18	118.30
1	A	48	PRO	N-CD-CG	5.64	111.65	103.20
1	A	201	ASP	CB-CG-OD2	-5.33	113.50	118.30
1	A	194	GLY	C-N-CD	-5.26	109.03	120.60
1	A	46	GLN	O-C-N	5.21	131.04	122.70
1	A	46	GLN	N-CA-CB	5.16	119.89	110.60
1	A	151	GLU	N-CA-CB	5.13	119.83	110.60
1	A	62	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	A	161	THR	N-CA-CB	-5.05	100.71	110.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	47	ASN	CA

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1697	0	1656	58	0
2	B	28	0	25	1	0
3	A	28	0	26	2	0
4	A	3	0	6	0	0
5	A	53	0	0	1	0
All	All	1809	0	1713	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:VAL:O	1:A:168:GLY:HA2	1.85	0.76
1:A:47:ASN:O	1:A:50:VAL:HG22	1.90	0.72
1:A:165:THR:O	1:A:166:ARG:C	2.24	0.70
1:A:43:PHE:C	1:A:44:GLN:O	2.33	0.65
1:A:85:GLN:HG3	1:A:155:ARG:HH12	1.63	0.63
1:A:5:ARG:HD2	3:A:1451:NAG:H83	1.80	0.63
1:A:47:ASN:H	1:A:48:PRO:HD3	1.64	0.62
1:A:52:THR:HG22	1:A:53:VAL:N	2.15	0.62
1:A:130:SER:HB2	5:A:1515:HOH:O	1.98	0.61
1:A:143:PHE:CE2	3:A:1451:NAG:H82	2.36	0.60
1:A:163:VAL:HG12	1:A:165:THR:H	1.67	0.59
1:A:85:GLN:HG3	1:A:85:GLN:O	2.01	0.59
1:A:63:ARG:HD3	1:A:63:ARG:N	2.18	0.57
1:A:163:VAL:HG21	1:A:168:GLY:HA3	1.85	0.57
1:A:66:GLN:N	1:A:66:GLN:OE1	2.38	0.57
1:A:111:PRO:O	1:A:210:ARG:NH1	2.37	0.57
1:A:49:GLY:C	1:A:51:SER:H	2.06	0.56
1:A:147:THR:HG22	1:A:162:GLY:O	2.05	0.56
1:A:29:ALA:HB2	1:A:178:PRO:HG3	1.87	0.56
1:A:84:GLN:HA	1:A:84:GLN:OE1	2.08	0.52
1:A:149:THR:HB	1:A:150:PRO:HD2	1.93	0.51
1:A:161:THR:HG22	1:A:201:ASP:HB2	1.92	0.51
1:A:43:PHE:O	1:A:44:GLN:O	2.29	0.50
1:A:219:ASN:N	1:A:219:ASN:ND2	2.59	0.50
1:A:63:ARG:O	1:A:64:GLU:C	2.51	0.48
1:A:44:GLN:HA	1:A:44:GLN:OE1	2.13	0.48
1:A:166:ARG:CZ	1:A:198:ARG:HG3	2.44	0.48
1:A:165:THR:HG22	1:A:166:ARG:O	2.13	0.48
1:A:168:GLY:O	1:A:197:GLY:HA2	2.13	0.48
1:A:47:ASN:N	1:A:48:PRO:HD3	2.20	0.48
1:A:219:ASN:N	1:A:219:ASN:HD22	2.12	0.47
1:A:166:ARG:HB3	1:A:167:ARG:HG3	1.97	0.47
1:A:195:PRO:O	1:A:196:CYS:C	2.54	0.47
1:A:218:ASN:C	1:A:220:PRO:HD3	2.35	0.47
1:A:163:VAL:HG12	1:A:164:LEU:N	2.30	0.46
2:B:1:NAG:H61	2:B:2:NAG:C1	2.46	0.46
1:A:52:THR:HG22	1:A:53:VAL:H	1.81	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:LEU:HD23	1:A:101:LEU:N	2.32	0.45
1:A:101:LEU:N	1:A:101:LEU:CD2	2.78	0.45
1:A:35:PHE:CD1	1:A:217:LEU:CD2	3.00	0.45
1:A:219:ASN:N	1:A:220:PRO:HD3	2.32	0.44
1:A:63:ARG:N	1:A:63:ARG:CD	2.80	0.44
1:A:47:ASN:N	1:A:47:ASN:OD1	2.49	0.44
1:A:113:GLN:O	1:A:114:ASN:HB2	2.18	0.43
1:A:52:THR:CG2	1:A:53:VAL:N	2.78	0.43
1:A:32:HIS:O	1:A:34:ARG:N	2.52	0.42
1:A:171:CYS:O	1:A:174:ASP:HB2	2.19	0.42
1:A:193:LEU:HA	1:A:193:LEU:HD12	1.73	0.42
1:A:35:PHE:HA	1:A:93:LEU:O	2.20	0.42
1:A:54:VAL:CG1	1:A:68:ARG:HD3	2.49	0.41
1:A:205:ARG:HG2	1:A:205:ARG:HH11	1.85	0.41
1:A:108:LEU:HA	1:A:109:PRO:HD2	1.85	0.41
1:A:55:LEU:HD12	1:A:71:PHE:HE2	1.86	0.41
1:A:101:LEU:HA	1:A:101:LEU:HD22	1.80	0.40
1:A:18:ILE:HG13	1:A:53:VAL:HG22	2.03	0.40
1:A:172:ASN:OD1	1:A:172:ASN:N	2.50	0.40
1:A:15:LEU:HA	1:A:15:LEU:HD12	1.92	0.40
1:A:164:LEU:HD23	1:A:164:LEU:HA	1.99	0.40
1:A:166:ARG:NH2	1:A:198:ARG:HG3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	223/225 (99%)	197 (88%)	18 (8%)	8 (4%)	3 4

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	47	ASN
1	A	200	PRO
1	A	44	GLN
1	A	46	GLN
1	A	48	PRO
1	A	33	ALA
1	A	50	VAL
1	A	134	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	183/183 (100%)	171 (93%)	12 (7%)	16 32

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ASN
1	A	63	ARG
1	A	78	GLU
1	A	101	LEU
1	A	133	SER
1	A	151	GLU
1	A	152	ASP
1	A	161	THR
1	A	166	ARG
1	A	172	ASN
1	A	180	VAL
1	A	219	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	HIS
1	A	158	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	219	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	1.22	1 (7%)	17,19,21	2.73	4 (23%)
2	NAG	B	2	2	14,14,15	1.12	0	17,19,21	3.32	10 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	1/1/5/7	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	O4-C4	2.41	1.48	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-O5-C5	7.12	121.84	112.19
2	B	2	NAG	O5-C1-C2	-6.63	100.82	111.29
2	B	2	NAG	C1-O5-C5	6.48	120.97	112.19
2	B	2	NAG	C2-N2-C7	-6.34	113.87	122.90
2	B	1	NAG	C8-C7-N2	-5.15	107.38	116.10
2	B	1	NAG	O5-C1-C2	-5.10	103.24	111.29
2	B	2	NAG	C4-C3-C2	-3.73	105.55	111.02
2	B	2	NAG	C6-C5-C4	-3.43	104.96	113.00
2	B	1	NAG	O7-C7-C8	3.22	128.04	122.06
2	B	2	NAG	C3-C4-C5	2.70	115.06	110.24
2	B	2	NAG	C1-C2-N2	2.69	115.09	110.49
2	B	2	NAG	O4-C4-C3	2.57	116.29	110.35
2	B	2	NAG	O3-C3-C4	-2.51	104.55	110.35
2	B	2	NAG	C8-C7-N2	-2.24	112.30	116.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	2	NAG	C1

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C8-C7-N2-C2
2	B	1	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0
2	B	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	A	1001	1	14,14,15	1.39	3 (21%)	17,19,21	2.46	7 (41%)
3	NAG	A	1451	1	14,14,15	1.25	2 (14%)	17,19,21	3.19	7 (41%)
4	EOH	A	300	-	2,2,2	0.71	0	1,1,1	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1001	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1451	1	-	0/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	NAG	O4-C4	2.45	1.48	1.43
3	A	1001	NAG	C8-C7	2.37	1.55	1.50
3	A	1451	NAG	O4-C4	2.23	1.48	1.43
3	A	1001	NAG	C3-C2	2.17	1.57	1.52
3	A	1451	NAG	C8-C7	2.09	1.54	1.50

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1451	NAG	C8-C7-N2	-8.48	101.75	116.10
3	A	1451	NAG	O7-C7-C8	5.69	132.62	122.06
3	A	1001	NAG	C4-C3-C2	5.07	118.45	111.02
3	A	1451	NAG	C1-O5-C5	4.63	118.47	112.19
3	A	1001	NAG	C1-O5-C5	4.63	118.46	112.19
3	A	1001	NAG	O5-C5-C6	3.76	113.09	107.20
3	A	1451	NAG	O4-C4-C3	3.58	118.63	110.35
3	A	1451	NAG	O3-C3-C2	-3.05	103.16	109.47
3	A	1001	NAG	C3-C4-C5	-3.03	104.83	110.24
3	A	1001	NAG	C8-C7-N2	-3.03	110.97	116.10
3	A	1001	NAG	O5-C5-C4	-2.81	104.00	110.83
3	A	1451	NAG	O5-C5-C4	-2.51	104.72	110.83
3	A	1001	NAG	O7-C7-N2	-2.22	117.86	121.95
3	A	1451	NAG	O5-C1-C2	-2.15	107.90	111.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1451	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/225 (100%)	0.45	17 (7%) 13 14	15, 30, 69, 87	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	PRO	7.1
1	A	49	GLY	6.8
1	A	45	SER	5.3
1	A	46	GLN	5.1
1	A	222	PRO	4.4
1	A	152	ASP	3.9
1	A	50	VAL	3.6
1	A	44	GLN	3.2
1	A	167	ARG	3.2
1	A	221	GLY	3.0
1	A	47	ASN	3.0
1	A	165	THR	2.9
1	A	84	GLN	2.7
1	A	132	ARG	2.3
1	A	225	ALA	2.3
1	A	134	GLY	2.2
1	A	166	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	2	14/15	0.46	0.63	56,100,100,100	0
2	NAG	B	1	14/15	0.73	0.30	12,54,100,100	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	1001	14/15	0.65	0.38	54,100,100,100	0
3	NAG	A	1451	14/15	0.83	0.31	22,61,91,100	0
4	EOH	A	300	3/3	0.84	0.30	23,23,36,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.