



wwPDB X-ray Structure Validation Summary Report i

Sep 23, 2025 – 09:29 pm BST

PDB ID : 1GZV / pdb_00001gsv
Title : The crystal structure of phosphoglucose isomerase from pig muscle complexed with 5-phosphoarabinonate
Authors : Davies, C.; Muirhead, H.
Deposited on : 2002-06-06
Resolution : 3.51 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbit	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	1165 (3.58-3.46)
Ramachandran outliers	177936	1150 (3.58-3.46)
Sidechain outliers	177891	1151 (3.58-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	A	557		59%	34%	6%

2 Entry composition [\(i\)](#)


There are 2 unique types of molecules in this entry. The entry contains 4437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUCOSE-6-PHOSPHATE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	554	Total	C	N	O	S	0	0	0
			4422	2828	768	810	16			

- Molecule 2 is 5-PHOSPHOARABINONIC ACID (CCD ID: PA5) (formula: C₅H₁₁O₉P).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			15	5	9	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: GLUCOSE-6-PHOSPHATE ISOMERASE



4 Data and refinement statistics i

Xtriaze (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value			Source
Space group	P 43 21 2			Depositor
Cell constants a, b, c, α , β , γ	95.60 \AA 90.00 $^{\circ}$	95.60 \AA 90.00 $^{\circ}$	138.70 \AA 90.00 $^{\circ}$	Depositor
Resolution (\AA)	15.00 – 3.51			Depositor
% Data completeness (in resolution range)	92.8 (15.00-3.51)			Depositor
R_{merge}	0.04			Depositor
R_{sym}	(Not available)			Depositor
Refinement program	REFMAC 5.0.32			Depositor
R , R_{free}	0.190, 0.249			Depositor
Estimated twinning fraction	No twinning to report.			Xtriaze
Total number of atoms	4437			wwPDB-VP
Average B, all atoms (\AA^2)	42.0			wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PA5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.12	6/4533 (0.1%)	1.36	35/6136 (0.6%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	148	ILE	CA-CB	-6.70	1.45	1.54
1	A	183	VAL	CA-CB	-6.22	1.46	1.54
1	A	165	MET	SD-CE	-5.79	1.65	1.79
1	A	492	MET	SD-CE	-5.35	1.66	1.79
1	A	524	ILE	CA-CB	-5.14	1.47	1.54

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	388	HIS	N-CA-C	-8.94	101.92	112.92
1	A	383	GLY	N-CA-C	-7.43	101.21	113.02
1	A	359	ASN	N-CA-C	7.06	121.61	112.92
1	A	91	PHE	N-CA-C	6.99	119.93	111.82
1	A	353	GLN	N-CA-C	-6.92	103.67	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4422	0	4376	149	0
2	A	15	0	8	4	0
All	All	4437	0	4384	149	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 149 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:THR:HG22	1:A:366:SER:H	1.20	1.03
1:A:214:THR:OG1	2:A:600:PA5:O2P	1.96	0.82
1:A:448:LEU:HD13	1:A:458:PHE:HB2	1.63	0.81
1:A:328:ASN:HD21	1:A:504:TRP:HE1	1.29	0.79
1:A:103:ARG:HD2	1:A:303:ASP:OD2	1.82	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	552/557 (99%)	516 (94%)	34 (6%)	2 (0%)	30 63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	520	LEU
1	A	233	LYS

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	476/479 (99%)	422 (89%)	54 (11%)	4 22

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	249	THR
1	A	364	THR
1	A	523	LYS
1	A	254	GLU
1	A	319	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	300	HIS
1	A	474	ASN
1	A	507	ASN
1	A	500	GLN
1	A	57	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PA5	A	600	-	14,14,14	1.19	2 (14%)	19,20,20	1.56	5 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PA5	A	600	-	-	1/18/18/18	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	PA5	O1-C1	-2.85	1.21	1.30
2	A	600	PA5	C2-C1	-2.58	1.49	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	PA5	O3P-P-O2P	3.59	121.37	107.64
2	A	600	PA5	O2-C2-C3	-3.05	103.98	110.45
2	A	600	PA5	O2-C2-C1	-2.51	105.41	110.66
2	A	600	PA5	O3-C3-C2	-2.18	105.24	109.21
2	A	600	PA5	O1-C1-O1A	-2.03	119.49	124.09

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	600	PA5	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	PA5	4	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [\(i\)](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [\(i\)](#)

EDS was not executed - this section is therefore empty.