



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 3, 2023 – 06:01 AM EDT

PDB ID : 1HR9  
Title : Yeast Mitochondrial Processing Peptidase beta-E73Q Mutant Complexed with Malate Dehydrogenase Signal Peptide  
Authors : Taylor, A.B.; Smith, B.S.; Kitada, S.; Kojima, K.; Miyaura, H.; Otwinowski, Z.; Ito, A.; Deisenhofer, J.  
Deposited on : 2000-12-21  
Resolution : 3.01 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.34  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.34


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2399 (3.04-3.00)
Clashscore	141614	2734 (3.04-3.00)
Ramachandran outliers	138981	2640 (3.04-3.00)
Sidechain outliers	138945	2643 (3.04-3.00)
RSRZ outliers	127900	2287 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	 61% 31% • 5%
1	C	475	 63% 28% • 6%
1	E	475	 62% 30% • 6%
1	G	475	 63% 30% • 5%
2	B	443	 67% 27% 5% •

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	443	 <p>% 67% 28% 5%</p>
2	F	443	 <p>% 66% 29% 5%</p>
2	H	443	 <p>2% 67% 27% 5%</p>
3	O	8	 <p>75% 25%</p>
3	P	8	 <p>50% 50%</p>
3	Q	8	 <p>50% 50%</p>
3	R	8	 <p>25% 75%</p>

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 27925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOCHONDRIAL PROCESSING PEPTIDASE ALPHA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	452	3499	2213	594	673	19	0	0	0
1	C	448	3473	2199	587	668	19	0	0	0
1	E	446	3465	2193	587	666	19	0	0	0
1	G	451	3493	2210	593	671	19	0	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	177	GLY	GLU	SEE REMARK 999	UNP P11914
A	217	GLY	GLU	SEE REMARK 999	UNP P11914
A	483	HIS	-	expression tag	UNP P11914
A	484	HIS	-	expression tag	UNP P11914
A	485	HIS	-	expression tag	UNP P11914
A	486	HIS	-	expression tag	UNP P11914
A	487	HIS	-	expression tag	UNP P11914
A	488	HIS	-	expression tag	UNP P11914
C	177	GLY	GLU	SEE REMARK 999	UNP P11914
C	217	GLY	GLU	SEE REMARK 999	UNP P11914
C	483	HIS	-	expression tag	UNP P11914
C	484	HIS	-	expression tag	UNP P11914
C	485	HIS	-	expression tag	UNP P11914
C	486	HIS	-	expression tag	UNP P11914
C	487	HIS	-	expression tag	UNP P11914
C	488	HIS	-	expression tag	UNP P11914
E	177	GLY	GLU	SEE REMARK 999	UNP P11914
E	217	GLY	GLU	SEE REMARK 999	UNP P11914
E	483	HIS	-	expression tag	UNP P11914
E	484	HIS	-	expression tag	UNP P11914

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	485	HIS	-	expression tag	UNP P11914
E	486	HIS	-	expression tag	UNP P11914
E	487	HIS	-	expression tag	UNP P11914
E	488	HIS	-	expression tag	UNP P11914
G	177	GLY	GLU	SEE REMARK 999	UNP P11914
G	217	GLY	GLU	SEE REMARK 999	UNP P11914
G	483	HIS	-	expression tag	UNP P11914
G	484	HIS	-	expression tag	UNP P11914
G	485	HIS	-	expression tag	UNP P11914
G	486	HIS	-	expression tag	UNP P11914
G	487	HIS	-	expression tag	UNP P11914
G	488	HIS	-	expression tag	UNP P11914

- Molecule 2 is a protein called MITOCHONDRIAL PROCESSING PEPTIDASE BETA SUB-UNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	439	3414	2148	591	668	7	0	0	0
2	D	441	3431	2159	594	671	7	0	0	0
2	F	443	3442	2165	596	674	7	0	0	0
2	H	440	3422	2154	592	669	7	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	20	ALA	-	cloning artifact	UNP P10507
B	73	GLN	GLU	engineered mutation	UNP P10507
B	84	PRO	SER	SEE REMARK 999	UNP P10507
B	350	ARG	GLN	SEE REMARK 999	UNP P10507
D	20	ALA	-	cloning artifact	UNP P10507
D	73	GLN	GLU	engineered mutation	UNP P10507
D	84	PRO	SER	SEE REMARK 999	UNP P10507
D	350	ARG	GLN	SEE REMARK 999	UNP P10507
F	20	ALA	-	cloning artifact	UNP P10507
F	73	GLN	GLU	engineered mutation	UNP P10507
F	84	PRO	SER	SEE REMARK 999	UNP P10507
F	350	ARG	GLN	SEE REMARK 999	UNP P10507
H	20	ALA	-	cloning artifact	UNP P10507

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
H	73	GLN	GLU	engineered mutation	UNP P10507
H	84	PRO	SER	SEE REMARK 999	UNP P10507
H	350	ARG	GLN	SEE REMARK 999	UNP P10507


- Molecule 3 is a protein called MALATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	O	8	Total	C	N	O	0	0	0
			63	38	15	10			
3	P	8	Total	C	N	O	0	0	0
			63	38	15	10			
3	Q	8	Total	C	N	O	0	0	0
			63	38	15	10			
3	R	8	Total	C	N	O	0	0	0
			63	38	15	10			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	D	1	Total	Zn	0	0
			1	1		
4	F	1	Total	Zn	0	0
			1	1		
4	H	1	Total	Zn	0	0
			1	1		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
5	E	1	Total	8	2	4	1	0	0
5	G	1	Total	8	2	4	1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: MITOCHONDRIAL PROCESSING PEPTIDASE ALPHA SUBUNIT




#### • Molecule 1: MITOCHONDRIAL PROCESSING PEPTIDASE ALPHA SUBUNIT








● Molecule 1: MITOCHONDRIAL PROCESSING PEPTIDASE ALPHA SUBUNIT




● Molecule 1: MITOCHONDRIAL PROCESSING PEPTIDASE ALPHA SUBUNIT




● Molecule 2: MITOCHONDRIAL PROCESSING PEPTIDASE BETA SUBUNIT






● Molecule 2: MITOCHONDRIAL PROCESSING PEPTIDASE BETA SUBUNIT




● Molecule 2: MITOCHONDRIAL PROCESSING PEPTIDASE BETA SUBUNIT




- Molecule 2: MITOCHONDRIAL PROCESSING PEPTIDASE BETA SUBUNIT

Chain H:  2% 67% 27% 5%




- Molecule 3: MALATE DEHYDROGENASE

Chain O:  75% 25%




- Molecule 3: MALATE DEHYDROGENASE

Chain P:  50% 50%




- Molecule 3: MALATE DEHYDROGENASE

Chain Q:  50% 50%



- Molecule 3: MALATE DEHYDROGENASE

Chain R:  25% 75%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.37Å 178.52Å 201.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.55 – 3.01 47.55 – 3.01	Depositor EDS
% Data completeness (in resolution range)	93.3 (47.55-3.01) 93.4 (47.55-3.01)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.52 (at 3.01Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.217 , 0.256 0.204 , 0.240	Depositor DCC
$R_{free}$ test set	2035 reflections (2.25%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	69.5	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 64.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	27925	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	0/3570	0.81	2/4830 (0.0%)
1	C	0.66	0/3544	0.79	3/4795 (0.1%)
1	E	0.61	0/3535	0.87	4/4784 (0.1%)
1	G	0.65	2/3564 (0.1%)	0.78	3/4822 (0.1%)
2	B	0.72	4/3478 (0.1%)	0.84	5/4720 (0.1%)
2	D	0.61	2/3495 (0.1%)	0.93	9/4744 (0.2%)
2	F	0.59	2/3506 (0.1%)	0.78	6/4759 (0.1%)
2	H	0.59	3/3486 (0.1%)	1.06	12/4732 (0.3%)
3	O	0.82	0/62	0.96	0/79
3	P	0.68	0/62	0.90	0/79
3	Q	0.66	0/62	0.86	0/79
3	R	0.74	0/62	0.79	0/79
All	All	0.64	13/28426 (0.0%)	0.86	44/38502 (0.1%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	81	GLN	CD-OE1	-10.06	1.01	1.24
2	B	81	GLN	CD-NE2	-9.57	1.08	1.32
2	D	354	ASN	CG-OD1	-8.19	1.05	1.24
2	H	354	ASN	CG-OD1	-7.75	1.06	1.24
2	F	354	ASN	CG-OD1	-7.44	1.07	1.24

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	253	ARG	NE-CZ-NH2	22.20	131.40	120.30
2	D	253	ARG	NE-CZ-NH1	-22.01	109.30	120.30
2	H	149	ARG	NE-CZ-NH2	-21.94	109.33	120.30
2	H	207	ARG	NE-CZ-NH1	-21.93	109.33	120.30
2	H	149	ARG	NE-CZ-NH1	21.33	130.97	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3460	134	0
1	C	3473	0	3435	119	0
1	E	3465	0	3426	119	0
1	G	3493	0	3455	129	0
2	B	3414	0	3414	116	0
2	D	3431	0	3432	124	0
2	F	3442	0	3442	119	0
2	H	3422	0	3424	111	0
3	O	63	0	73	3	0
3	P	63	0	73	4	0
3	Q	63	0	73	5	0
3	R	63	0	73	6	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
4	F	1	0	0	0	0
4	H	1	0	0	0	0
5	E	15	0	17	0	0
5	G	15	0	17	0	0
All	All	27925	0	27814	931	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 931 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:295:GLY:HA2	2:D:93:GLU:HG2	1.40	1.03
1:C:230:ILE:HD12	1:C:230:ILE:H	1.15	1.02
1:G:230:ILE:H	1:G:230:ILE:HD12	1.21	1.01
1:E:230:ILE:HD12	1:E:230:ILE:H	1.25	1.01
1:A:230:ILE:HD12	1:A:230:ILE:H	1.20	1.00

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/475 (94%)	417 (93%)	27 (6%)	4 (1%)	17	53
1	C	444/475 (94%)	413 (93%)	27 (6%)	4 (1%)	17	53
1	E	442/475 (93%)	412 (93%)	27 (6%)	3 (1%)	22	59
1	G	447/475 (94%)	412 (92%)	32 (7%)	3 (1%)	22	59
2	B	437/443 (99%)	403 (92%)	27 (6%)	7 (2%)	9	38
2	D	439/443 (99%)	401 (91%)	31 (7%)	7 (2%)	9	38
2	F	441/443 (100%)	408 (92%)	26 (6%)	7 (2%)	9	38
2	H	438/443 (99%)	397 (91%)	34 (8%)	7 (2%)	9	38
3	O	6/8 (75%)	6 (100%)	0	0	100	100
3	P	6/8 (75%)	6 (100%)	0	0	100	100
3	Q	6/8 (75%)	6 (100%)	0	0	100	100
3	R	6/8 (75%)	6 (100%)	0	0	100	100
All	All	3560/3704 (96%)	3287 (92%)	231 (6%)	42 (1%)	13	46

5 of 42 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	250	PHE
1	C	250	PHE
1	E	250	PHE
1	G	250	PHE
2	B	367	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/401 (95%)	357 (94%)	24 (6%)	18	49
1	C	379/401 (94%)	354 (93%)	25 (7%)	16	47
1	E	378/401 (94%)	357 (94%)	21 (6%)	21	54
1	G	380/401 (95%)	360 (95%)	20 (5%)	22	57
2	B	376/379 (99%)	353 (94%)	23 (6%)	18	51
2	D	378/379 (100%)	353 (93%)	25 (7%)	16	47
2	F	379/379 (100%)	358 (94%)	21 (6%)	21	55
2	H	377/379 (100%)	354 (94%)	23 (6%)	18	51
3	O	6/6 (100%)	6 (100%)	0	100	100
3	P	6/6 (100%)	6 (100%)	0	100	100
3	Q	6/6 (100%)	5 (83%)	1 (17%)	2	10
3	R	6/6 (100%)	5 (83%)	1 (17%)	2	10
All	All	3052/3144 (97%)	2868 (94%)	184 (6%)	19	51

5 of 184 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	363	THR
1	G	151	GLU
2	F	31	LEU
2	F	219	GLU
1	G	307	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 73 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	305	HIS
2	H	374	ASN
1	G	381	ASN
2	H	73	GLN

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	C	320	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EPE	E	489	-	15,15,15	1.69	5 (33%)	18,20,20	0.90	1 (5%)
5	EPE	G	489	-	15,15,15	1.79	5 (33%)	18,20,20	0.89	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	E	489	-	-	2/9/19/19	0/1/1/1
5	EPE	G	489	-	-	0/9/19/19	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	489	EPE	C10-S	4.06	1.83	1.77
5	E	489	EPE	C10-S	3.06	1.81	1.77
5	E	489	EPE	C6-N1	2.79	1.54	1.46
5	E	489	EPE	C9-N1	2.54	1.53	1.47
5	G	489	EPE	C7-N4	2.46	1.53	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	489	EPE	C7-N4-C5	-2.13	105.79	111.23
5	E	489	EPE	C7-N4-C5	-2.06	105.96	111.23

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	489	EPE	C10-C9-N1-C2
5	E	489	EPE	C10-C9-N1-C6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	452/475 (95%)	-0.29	5 (1%) 80 55	24, 52, 92, 101	0
1	C	448/475 (94%)	-0.34	1 (0%) 95 87	30, 54, 90, 101	0
1	E	446/475 (93%)	-0.23	3 (0%) 87 68	36, 63, 98, 101	0
1	G	451/475 (94%)	-0.28	3 (0%) 87 68	28, 57, 94, 101	0
2	B	439/443 (99%)	-0.40	0 100 100	26, 51, 83, 101	0
2	D	441/443 (99%)	-0.16	3 (0%) 87 68	23, 75, 101, 101	0
2	F	443/443 (100%)	-0.14	3 (0%) 87 68	47, 84, 101, 101	0
2	H	440/443 (99%)	0.10	9 (2%) 65 36	50, 95, 101, 101	0
3	O	8/8 (100%)	-0.44	0 100 100	36, 50, 81, 88	0
3	P	8/8 (100%)	-0.29	0 100 100	79, 94, 101, 101	0
3	Q	8/8 (100%)	-0.05	0 100 100	65, 96, 101, 101	0
3	R	8/8 (100%)	-0.14	0 100 100	84, 97, 101, 101	0
All	All	3592/3704 (96%)	-0.22	27 (0%) 86 65	23, 66, 101, 101	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	252	ASN	4.2
1	E	252	ASN	3.6
1	G	250	PHE	3.0
1	C	252	ASN	3.0
2	H	440	SER	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EPE	E	489	15/15	0.92	0.30	84,84,84,84	0
5	EPE	G	489	15/15	0.92	0.31	91,91,91,91	0
4	ZN	H	504	1/1	0.98	0.10	101,101,101,101	0
4	ZN	D	502	1/1	0.98	0.11	99,99,99,99	0
4	ZN	F	503	1/1	0.98	0.11	91,91,91,91	0
4	ZN	B	501	1/1	0.99	0.12	51,51,51,51	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.