

# wwPDB X-ray Structure Validation Summary Report (i)

#### Aug 3, 2023 – 06:59 AM EDT

PDB ID : 1I6V

Title : THERMUS AQUATICUS CORE RNA POLYMERASE-RIFAMPICIN

**COMPLEX** 

Authors: Campbell, E.A.; Korzheva, N.; Mustaev, A.; Murakami, K.; Goldfarb, A.;

Darst, S.A.

Deposited on : 2001-03-05

Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.34

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution		
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$		
Clashscore	141614	1205 (3.34-3.26)		
Ramachandran outliers	138981	1183 (3.34-3.26)		
Sidechain outliers	138945	1182 (3.34-3.26)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length		Quality of chain								
1	A	314	12%	40%	15% •	29%						
1	В	314	11%	46%	16%	• 27%						
2	С	1118	14%	60%		24%						
3	D	1264	18%	52%		21% • 7%						
4	Е	99	19%	62%		18%						



# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 21292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	224	Total 1741	C 1109	11	O 330	S 3	0	0	0
1	В	230	Total 1761	C 1122	- '	O 337	S 3	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	112	VAL	GLY	$\operatorname{conflict}$	UNP Q9KWU8
A	232	SER	LEU	conflict	UNP Q9KWU8
В	112	VAL	GLY	conflict	UNP Q9KWU8
В	232	SER	LEU	conflict	UNP Q9KWU8

• Molecule 2 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
2	С	1113	Total 8508	C 5386	N 1514	O 1585	S 23	12	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	2	LYS	GLU	$\operatorname{conflict}$	UNP Q9KWU7
С	?	-	GLU	deletion	UNP Q9KWU7
С	1111	VAL	ILE	conflict	UNP Q9KWU7

• Molecule 3 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
3	D	1174	Total 8502	C 5329	N 1550	O 1596	S 27	17	0	0




There are 9	discrepancies	between	the modelled	and	reference sequences:
I HOLO OLO U	ansoropanions	DCCWCCII	une moderned	alla	reference bequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	70	ALA	GLY	$\operatorname{conflict}$	UNP Q9KWU6
D	77	ALA	GLY	conflict	UNP Q9KWU6
D	91	ALA	GLY	$\operatorname{conflict}$	UNP Q9KWU6
D	113	ALA	GLY	$\operatorname{conflict}$	UNP Q9KWU6
D	139	ALA	GLY	$\operatorname{conflict}$	UNP Q9KWU6
D	144	ALA	GLY	$\operatorname{conflict}$	UNP Q9KWU6
D	863	THR	VAL	conflict	UNP Q9KWU6
D	866	THR	VAL	conflict	UNP Q9KWU6
D	1009	ASN	LYS	conflict	UNP Q9KWU6

• Molecule 4 is a protein called DNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
4	E	98	Total	С	N	О	S	0	0	0
T		30	719	453	132	130	4			U

 $\bullet$  Molecule 5 is RIFAMPICIN (three-letter code: RFP) (formula:  $\mathrm{C_{43}H_{58}N_4O_{12}}).$ 



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	С	1	Total 59	C 43	N 4	O 12	0	0

• Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	1	Total Mg 1 1	0	0

 $\bullet$  Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total Zn 1 1	0	0




# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: DNA-DIRECTED RNA POLYMERASE




• Molecule 1: DNA-DIRECTED RNA POLYMERASE









• Molecule 2: DNA-DIRECTED RNA POLYMERASE









• Molecule 3: DNA-DIRECTED RNA POLYMERASE











# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 41 21 2	Depositor	
Cell constants	199.45Å 199.45Å 289.13Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	8.00 - 3.30	Depositor	
% Data completeness	(Not available) (8.00-3.30)	Depositor	
(in resolution range)	(110t available) (0.00 9.90)		
$R_{merge}$	0.08	Depositor	
$R_{sym}$	(Not available)	Depositor	
Refinement program	CNS	Depositor	
$R, R_{free}$	0.276 , $0.359$	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	21292	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP	



## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: RFP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5	
1	A	0.43	0/1775	0.77	1/2417 (0.0%)	
1	В	0.42	1/1795~(0.1%)	0.77	$1/2447 \ (0.0\%)$	
2	С	0.41	0/8672	0.80	7/11752 (0.1%)	
3	D	0.44	1/8439~(0.0%)	0.84	19/11447 (0.2%)	
4	Е	0.37	0/730	0.71	0/991	
All	All	0.42	$2/21411 \ (0.0\%)$	0.81	28/29054 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	С	0	1
3	D	0	3
All	All	0	5

#### All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
3	D	1270	ALA	C-N	-12.09	1.06	1.34
1	В	94	MET	SD-CE	5.75	2.10	1.77

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
3	D	1270	ALA	O-C-N	-19.05	92.23	122.70
3	D	1270	ALA	C-N-CA	13.67	155.87	121.70
3	D	1270	ALA	CA-C-N	12.08	143.77	117.20

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
2	С	580	MET	N-CA-C	6.98	129.85	111.00
3	D	1166	LEU	CA-CB-CG	6.22	129.61	115.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	139	TYR	Sidechain
2	С	1013	TYR	Sidechain
3	D	1165	TYR	Sidechain
3	D	1268	PRO	Mainchain
3	D	1270	ALA	Mainchain

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1741	0	1747	348	0
1	В	1761	0	1754	316	0
2	С	8508	0	8421	1817	0
3	D	8502	0	8002	1671	0
4	Е	719	0	685	138	0
5	С	59	0	56	6	0
6	D	1	0	0	0	0
7	D	1	0	0	0	0
All	All	21292	0	20665	4041	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 96.

The worst 5 of 4041 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:94:MET:CE	1:B:94:MET:SD	2.10	1.39
2:C:690:ILE:HB	2:C:852:ILE:HG22	1.19	1.18

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{aligned}$	Clash overlap (Å)
3:D:1280:VAL:HG12	3:D:1281:VAL:HG23	1.25	1.16
2:C:491:GLU:HA	2:C:531:PHE:HA	1.29	1.15
3:D:772:PRO:HG3	3:D:778:LEU:HB2	1.28	1.14

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	222/314~(71%)	111 (50%)	61 (28%)	50 (22%)	0 0
1	В	228/314 (73%)	123 (54%)	62 (27%)	43 (19%)	0 1
2	С	1111/1118 (99%)	591 (53%)	277 (25%)	243 (22%)	0 0
3	D	1126/1264 (89%)	588 (52%)	291 (26%)	247 (22%)	0 0
4	E	96/99 (97%)	50 (52%)	24 (25%)	22 (23%)	0 0
All	All	2783/3109 (90%)	1463 (53%)	715 (26%)	605 (22%)	0 0

5 of 605 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	PHE
1	A	19	HIS
1	A	26	GLU
1	A	59	GLU
1	A	73	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the	number of r	residues for	which the	${\rm sidechain}$	conformation	was
analysed, and the total number of	residues.					

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	A	189/271 (70%)	161 (85%)	28 (15%)	3	13
1	В	190/271 (70%)	169 (89%)	21 (11%)	6	23
2	С	870/935 (93%)	770 (88%)	100 (12%)	5	22
3	D	782/1035 (76%)	671 (86%)	111 (14%)	3	15
4	E	67/88 (76%)	62 (92%)	5 (8%)	13	39
All	All	2098/2600 (81%)	1833 (87%)	265 (13%)	4	19

5 of 265 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	1166	LEU
3	D	1210	SER
4	Е	32	ARG
2	С	438	ILE
2	С	433	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 71 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	861	GLN
3	D	909	ASN
3	D	1374	GLN
2	С	565	GLN
2	С	545	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



### 5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Pos	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
IVIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	RFP	С	1640	-	63,63,63	1.26	9 (14%)	94,94,94	0.95	4 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	RFP	С	1640	-	-	13/60/85/85	0/5/5/5

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
5	С	1640	RFP	O5-C29	3.47	1.48	1.39
5	С	1640	RFP	O4-C11	3.12	1.27	1.21
5	С	1640	RFP	C5-C10	2.99	1.49	1.43
5	С	1640	RFP	C39-N4	2.98	1.52	1.46
5	С	1640	RFP	C8-C9	2.77	1.51	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
5	С	1640	RFP	C34-C26-C25	-2.37	107.15	111.40
5	С	1640	RFP	C5-C10-C9	-2.33	115.41	119.66
5	С	1640	RFP	O12-C4-C10	2.16	124.19	119.00
5	С	1640	RFP	C24-C23-C22	2.12	118.97	115.43

There are no chirality outliers.

5 of 13 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
5	С	1640	RFP	C17-C18-C19-C20
5	С	1640	RFP	C26-C27-C28-C29
5	С	1640	RFP	O6-C27-C28-C29
5	С	1640	RFP	C32-C22-C23-C24
5	С	1640	RFP	C21-C22-C23-C24

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	С	1640	RFP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	D	4

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	155:ASP	С	2(U):UNK	N	54.72
1	D	46(U):UNK	С	452:ILE	N	47.08
1	D	10(U):UNK	С	20(U):UNK	N	15.58
1	D	1270:ALA	С	1271:LYS	N	1.06



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

