



wwPDB X-ray Structure Validation Summary Report

Dec 19, 2023 – 07:06 AM EST

PDB ID : 1IQC
Title : Crystal structure of Di-Heme Peroxidase from Nitrosomonas europaea
Authors : Shimizu, H.
Deposited on : 2001-07-20
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	308	
1	B	308	
1	C	308	
1	D	308	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10767 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called di-heme peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	S	0	0	0
			2388	1497	411	467	13			
1	B	308	Total	C	N	O	S	0	0	0
			2388	1497	411	467	13			
1	C	308	Total	C	N	O	S	0	0	0
			2388	1497	411	467	13			
1	D	308	Total	C	N	O	S	0	0	0
			2388	1497	411	467	13			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	B	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		
2	D	1	Total	Ca	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is HEME C (three-letter code: HEC) (formula: C₃₄H₃₄FeN₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	C O	0	0
			6	3 3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	260	Total	O	0	0
			260	260		
6	B	199	Total	O	0	0
			199	199		
6	C	279	Total	O	0	0
			279	279		
6	D	120	Total	O	0	0
			120	120		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: di-heme peroxidase



- Molecule 1: di-heme peroxidase



- Molecule 1: di-heme peroxidase



- Molecule 1: di-heme peroxidase



A1	D83	K164	P241
N2	G84	M165	Y242
E3	R85	A166	F243
P4	A86	L167	H244
I5	P96	N168	L251
I8	P100	Q169	E256
V11	K101	D170	T257
T12	E102	E171	M258
T12	M103	L172	G259
M15	A104	E173	R260
M18	E108	L177	I261
A19	I109	C183	Q262
E20	A110	V184	L263
L21	A110	Q185	N264
G22	V113	C186	N268
K23	V114	H187	K269
M24	A115	M188	D270
L25	Q119	G189	E271
F26	Y120	V192	K274
D28	R121	G193	T281
P29	E122	G194	K290
R30	R123	S195	L291
L31	F124	S196	P292
S32	K125	Y197	I293
K33	K126	Q198	L294
S34	V127	K199	P295
G35	F128	M200	P296
F36	G128	G201	S297
I37	S130	K209	N298
S38	D131	M210	N299
C39	E132	P211	D300
N40	V133	T134	T301
S41	T134	R215	P302
C42	I135	M216	P306
L45	D136	D217	Y307
D51	R137	V218	E308
H59	I138	T219	
K60	T139	G220	
W61	A141	N221	
Q62	I142	E222	
Q63	A143	A223	
I66	Q144	D224	
L72	F145	R225	
N73	E146	N226	
N77	L149	V227	
L78	V150	F228	
A79	T151	K229	
Q80	P152	V230	
F81	K155	N235	
W82	F156	I236	
	D157	E237	
	E161	L238	
		T239	
		Y240	

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.13Å 55.11Å 144.00Å 90.00° 103.60° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80	Depositor
% Data completeness (in resolution range)	95.9 (30.00-1.80)	Depositor
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.222 , 0.257	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	10767	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, MG, GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2441	0.58	0/3304
1	B	0.29	0/2441	0.56	0/3304
1	C	0.32	0/2441	0.58	0/3304
1	D	0.29	0/2441	0.52	0/3304
All	All	0.30	0/9764	0.56	0/13216

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	240	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2388	0	2331	53	0
1	B	2388	0	2331	81	0
1	C	2388	0	2331	40	0
1	D	2388	0	2331	179	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	D	1	0	0	0	0
4	A	86	0	64	15	0
4	B	86	0	64	15	0
4	C	86	0	64	14	0
4	D	86	0	64	17	0
5	B	6	0	8	0	0
6	A	260	0	0	4	0
6	B	199	0	0	4	0
6	C	279	0	0	0	0
6	D	120	0	0	6	0
All	All	10767	0	9588	348	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 348 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:186:CYS:SG	4:B:402:HEC:HAC	1.50	1.50
1:D:42:CYS:SG	4:D:401:HEC:HAC	1.70	1.31
1:C:39:CYS:SG	4:C:401:HEC:HAB	1.72	1.25
1:A:183:CYS:SG	4:A:402:HEC:HAB	1.75	1.21
1:A:39:CYS:SG	4:A:401:HEC:HAB	1.77	1.15

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/308 (99%)	296 (97%)	10 (3%)	0	100	100
1	B	306/308 (99%)	287 (94%)	16 (5%)	3 (1%)	15	5
1	C	306/308 (99%)	298 (97%)	8 (3%)	0	100	100
1	D	306/308 (99%)	285 (93%)	19 (6%)	2 (1%)	22	10
All	All	1224/1232 (99%)	1166 (95%)	53 (4%)	5 (0%)	34	21

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2	ASN
1	D	2	ASN
1	B	163	ASP
1	D	215	ARG
1	B	3	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/260 (100%)	255 (98%)	5 (2%)	57	46
1	B	260/260 (100%)	254 (98%)	6 (2%)	50	37
1	C	260/260 (100%)	255 (98%)	5 (2%)	57	46
1	D	260/260 (100%)	250 (96%)	10 (4%)	33	18
All	All	1040/1040 (100%)	1014 (98%)	26 (2%)	47	34

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	264	ASN
1	D	23	LYS
1	D	268	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	15	ASN
1	D	126	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 56 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	15	ASN
1	D	299	ASN
1	C	176	ASN
1	D	298	ASN
1	D	144	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 7 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	HEC	C	402	1	32,50,50	1.44	3 (9%)	24,82,82	1.30	3 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	1410	-	5,5,5	0.29	0	5,5,5	0.27	0
4	HEC	A	401	1,6	32,50,50	1.24	3 (9%)	24,82,82	1.13	1 (4%)
4	HEC	B	402	1	32,50,50	1.40	3 (9%)	24,82,82	0.90	0
4	HEC	B	401	1,6	32,50,50	1.36	6 (18%)	24,82,82	0.97	1 (4%)
4	HEC	A	402	1	32,50,50	1.49	2 (6%)	24,82,82	1.26	1 (4%)
4	HEC	C	401	1,6	32,50,50	1.44	4 (12%)	24,82,82	1.15	2 (8%)
4	HEC	D	401	1,6	32,50,50	1.40	4 (12%)	24,82,82	0.91	0
4	HEC	D	402	1	32,50,50	1.43	4 (12%)	24,82,82	1.75	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEC	C	402	1	-	2/10/54/54	-
5	GOL	B	1410	-	-	0/4/4/4	-
4	HEC	A	401	1,6	-	1/10/54/54	-
4	HEC	B	402	1	-	2/10/54/54	-
4	HEC	B	401	1,6	-	2/10/54/54	-
4	HEC	A	402	1	-	2/10/54/54	-
4	HEC	C	401	1,6	-	2/10/54/54	-
4	HEC	D	401	1,6	-	5/10/54/54	-
4	HEC	D	402	1	-	3/10/54/54	-

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	402	HEC	C3C-C2C	-4.63	1.35	1.40
4	A	402	HEC	C2B-C3B	-4.60	1.35	1.40
4	A	402	HEC	C3C-C2C	-4.48	1.36	1.40
4	C	401	HEC	C2B-C3B	-4.30	1.36	1.40
4	D	402	HEC	C3C-C2C	-4.05	1.36	1.40

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	402	HEC	CBD-CAD-C3D	6.97	124.51	112.62
4	C	401	HEC	CBD-CAD-C3D	-2.99	107.52	112.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	402	HEC	CBD-CAD-C3D	-2.97	107.56	112.62
4	A	401	HEC	CBD-CAD-C3D	-2.90	107.66	112.62
4	C	402	HEC	CMB-C2B-C1B	-2.52	124.58	128.46

There are no chirality outliers.

5 of 19 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	402	HEC	C4D-C3D-CAD-CBD
4	D	401	HEC	C3A-C2A-CAA-CBA
4	C	401	HEC	CAA-CBA-CGA-O2A
4	D	401	HEC	CAA-CBA-CGA-O2A
4	D	402	HEC	CAD-CBD-CGD-O2D


There are no ring outliers.


8 monomers are involved in 61 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	402	HEC	7	0
4	A	401	HEC	7	0
4	B	402	HEC	6	0
4	B	401	HEC	9	0
4	A	402	HEC	8	0
4	C	401	HEC	7	0
4	D	401	HEC	6	0
4	D	402	HEC	11	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.