



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 19, 2023 – 02:37 PM EST

PDB ID : 1J3L  
Title : Structure of the RNA-processing inhibitor RraA from *Thermus thermophilis*  
Authors : Rehse, P.H.; Miyano, M.; Tahirov, T.H.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2003-02-04  
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*






The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	 2% 70% 28%
1	B	164	 4% 69% 28%
1	C	164	 3% 71% 25%
1	D	164	 9% 46% 48% 6%
1	E	164	 9% 45% 46% 9%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	164	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into four segments: a red segment (9%), a green segment (37%), a yellow segment (55%), and a small grey segment (2%). The percentages are labeled below the corresponding segments.</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Demethylmenaquinone Methyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	164	1224	775	218	229	1	1	0	0	0
1	B	164	1223	773	218	230	1	1	0	0	0
1	C	161	1211	769	215	225	1	1	0	0	0
1	D	164	1226	776	218	230	1	1	0	0	0
1	E	164	1226	776	218	230	1	1	0	0	0
1	F	157	1181	752	208	219	1	1	0	0	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	90	Total	O	0	0
			90	90		

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	116	Total 116	O 116	0	0
4	C	72	Total 72	O 72	0	0
4	D	60	Total 60	O 60	0	0
4	E	28	Total 28	O 28	0	0
4	F	32	Total 32	O 32	0	0






● Molecule 1: Demethylmenaquinone Methyltransferase



● Molecule 1: Demethylmenaquinone Methyltransferase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.87Å 109.07Å 270.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.40 – 2.30 38.39 – 2.29	Depositor EDS
% Data completeness (in resolution range)	85.3 (38.40-2.30) 88.7 (38.39-2.29)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 2.29Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.206 , 0.274 0.210 , 0.279	Depositor DCC
$R_{free}$ test set	3996 reflections (5.62%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.8	Xtrriage
Anisotropy	0.219	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 58.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.029 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.064 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7692	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1243	0.68	0/1676
1	B	0.38	0/1242	0.69	1/1674 (0.1%)
1	C	0.35	0/1230	0.62	0/1661
1	D	0.38	0/1245	0.65	0/1679
1	E	0.28	0/1245	0.57	0/1679
1	F	0.27	0/1200	0.55	0/1621
All	All	0.35	0/7405	0.63	1/9990 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	39	ARG	NE-CZ-NH1	-5.79	117.41	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1224	0	1253	37	0
1	B	1223	0	1249	41	0
1	C	1211	0	1246	33	0
1	D	1226	0	1257	73	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1226	0	1257	90	0
1	F	1181	0	1215	88	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	90	0	0	6	0
4	B	116	0	0	4	0
4	C	72	0	0	1	0
4	D	60	0	0	1	0
4	E	28	0	0	0	0
4	F	32	0	0	2	0
All	All	7692	0	7477	343	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 343 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:117:LYS:H	1:E:117:LYS:HE2	1.11	1.12
1:E:117:LYS:HE2	1:E:117:LYS:N	1.84	0.91
1:F:138:LEU:HD12	1:F:139:PRO:HD2	1.53	0.88
1:E:117:LYS:H	1:E:117:LYS:CE	1.86	0.88
1:C:8:LEU:HD12	1:C:144:LEU:HD22	1.56	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	162/164 (99%)	153 (94%)	6 (4%)	3 (2%)	<b>8</b> <b>7</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	162/164 (99%)	153 (94%)	5 (3%)	4 (2%)	5	4
1	C	159/164 (97%)	147 (92%)	10 (6%)	2 (1%)	12	12
1	D	162/164 (99%)	140 (86%)	17 (10%)	5 (3%)	4	2
1	E	162/164 (99%)	127 (78%)	19 (12%)	16 (10%)	0	0
1	F	155/164 (94%)	129 (83%)	19 (12%)	7 (4%)	2	1
All	All	962/984 (98%)	849 (88%)	76 (8%)	37 (4%)	3	1

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	SER
1	B	160	VAL
1	B	161	ARG
1	D	2	GLU
1	D	157	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/124 (100%)	120 (97%)	4 (3%)	39	54
1	B	124/124 (100%)	121 (98%)	3 (2%)	49	66
1	C	124/124 (100%)	120 (97%)	4 (3%)	39	54
1	D	125/124 (101%)	118 (94%)	7 (6%)	21	29
1	E	125/124 (101%)	120 (96%)	5 (4%)	31	44
1	F	121/124 (98%)	117 (97%)	4 (3%)	38	53
All	All	743/744 (100%)	716 (96%)	27 (4%)	35	49

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	34	ARG

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	127	ASP
1	F	34	ARG
1	D	114	THR
1	E	60	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	E	44	ASN
1	E	58	GLN
1	F	44	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	163/164 (99%)	-0.30	4 (2%) 57 64	13, 29, 64, 103	0
1	B	163/164 (99%)	-0.28	7 (4%) 35 42	15, 26, 53, 107	0
1	C	160/164 (97%)	-0.37	5 (3%) 49 56	21, 34, 59, 98	0
1	D	163/164 (99%)	0.33	15 (9%) 9 12	23, 50, 94, 110	0
1	E	163/164 (99%)	0.57	15 (9%) 9 12	36, 66, 98, 117	0
1	F	156/164 (95%)	0.53	15 (9%) 8 10	39, 73, 94, 107	0
All	All	968/984 (98%)	0.08	61 (6%) 20 25	13, 44, 92, 117	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	160	VAL	11.6
1	D	163	GLY	11.1
1	E	160	VAL	8.2
1	E	159	GLY	6.9
1	E	158	SER	6.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	1001	1/1	0.96	0.10	34,34,34,34	0
3	CL	A	1002	1/1	0.98	0.07	30,30,30,30	0
3	CL	B	1003	1/1	0.98	0.05	27,27,27,27	0

## 6.5 Other polymers

There are no such residues in this entry.