



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 7, 2023 – 04:56 AM EDT

PDB ID : 1N8J  
Title : Crystal Structure of AhpC with Active Site Cysteine mutated to Serine (C46S)  
Authors : Wood, Z.A.; Poole, L.B.; Karplus, P.A.  
Deposited on : 2002-11-20  
Resolution : 2.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	
1	B	186	
1	C	186	
1	D	186	
1	E	186	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	186	 5% 78% 22%
1	G	186	 2% 82% 18%
1	H	186	 0% 81% 18%
1	I	186	 4% 77% 20%
1	J	186	 3% 79% 21%
1	K	186	 0% 78% 22%
1	L	186	 2% 76% 23%
1	M	186	 4% 71% 28%
1	N	186	 4% 73% 26%
1	O	186	 0% 78% 22%
1	P	186	 2% 81% 18%
1	Q	186	 2% 83% 16%
1	R	186	 0% 78% 20%
1	S	186	 8% 67% 33%
1	T	186	 6% 66% 33%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 30710 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkyl hydroperoxide reductase C22 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	186	1456	928	241	284	3	0	0	0
1	B	186	1456	928	241	284	3	0	0	0
1	C	186	1456	928	241	284	3	0	0	0
1	D	186	1456	928	241	284	3	0	0	0
1	E	186	1456	928	241	284	3	0	0	0
1	F	186	1456	928	241	284	3	0	0	0
1	G	186	1456	928	241	284	3	0	0	0
1	H	186	1456	928	241	284	3	0	0	0
1	I	186	1456	928	241	284	3	0	0	0
1	J	186	1456	928	241	284	3	0	0	0
1	K	186	1456	928	241	284	3	0	0	0
1	L	186	1456	928	241	284	3	0	0	0
1	M	186	1456	928	241	284	3	0	0	0
1	N	186	1456	928	241	284	3	0	0	0
1	O	186	1456	928	241	284	3	0	0	0
1	P	186	1456	928	241	284	3	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	Q	186	1456	928	241	284	3	0	0	0
1	R	186	1456	928	241	284	3	0	0	0
1	S	186	1456	928	241	284	3	0	0	0
1	T	186	1456	928	241	284	3	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	SER	CYS	engineered mutation	UNP P0A251
B	46	SER	CYS	engineered mutation	UNP P0A251
C	46	SER	CYS	engineered mutation	UNP P0A251
D	46	SER	CYS	engineered mutation	UNP P0A251
E	46	SER	CYS	engineered mutation	UNP P0A251
F	46	SER	CYS	engineered mutation	UNP P0A251
G	46	SER	CYS	engineered mutation	UNP P0A251
H	46	SER	CYS	engineered mutation	UNP P0A251
I	46	SER	CYS	engineered mutation	UNP P0A251
J	46	SER	CYS	engineered mutation	UNP P0A251
K	46	SER	CYS	engineered mutation	UNP P0A251
L	46	SER	CYS	engineered mutation	UNP P0A251
M	46	SER	CYS	engineered mutation	UNP P0A251
N	46	SER	CYS	engineered mutation	UNP P0A251
O	46	SER	CYS	engineered mutation	UNP P0A251
P	46	SER	CYS	engineered mutation	UNP P0A251
Q	46	SER	CYS	engineered mutation	UNP P0A251
R	46	SER	CYS	engineered mutation	UNP P0A251
S	46	SER	CYS	engineered mutation	UNP P0A251
T	46	SER	CYS	engineered mutation	UNP P0A251

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	77	Total	O	0	0
			77	77		
2	B	71	Total	O	0	0
			71	71		
2	C	58	Total	O	0	0
			58	58		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	56	Total O 56 56	0	0
2	E	77	Total O 77 77	0	0
2	F	82	Total O 82 82	0	0
2	G	94	Total O 94 94	0	0
2	H	92	Total O 92 92	0	0
2	I	72	Total O 72 72	0	0
2	J	86	Total O 86 86	0	0
2	K	106	Total O 106 106	0	0
2	L	105	Total O 105 105	0	0
2	M	49	Total O 49 49	0	0
2	N	58	Total O 58 58	0	0
2	O	95	Total O 95 95	0	0
2	P	116	Total O 116 116	0	0
2	Q	104	Total O 104 104	0	0
2	R	93	Total O 93 93	0	0
2	S	52	Total O 52 52	0	0
2	T	47	Total O 47 47	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein



- Molecule 1: Alkyl hydroperoxide reductase C22 protein





- Molecule 1: Alkyl hydroperoxide reductase C22 protein








- Molecule 1: Alkyl hydroperoxide reductase C22 protein





- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein





- Molecule 1: Alkyl hydroperoxide reductase C22 protein











- Molecule 1: Alkyl hydroperoxide reductase C22 protein





- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein




- Molecule 1: Alkyl hydroperoxide reductase C22 protein



- Molecule 1: Alkyl hydroperoxide reductase C22 protein



- Molecule 1: Alkyl hydroperoxide reductase C22 protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.08Å 107.33Å 120.08Å 102.43° 116.22° 98.81°	Depositor
Resolution (Å)	19.99 – 2.17 20.40 – 2.16	Depositor EDS
% Data completeness (in resolution range)	89.3 (19.99-2.17) 88.8 (20.40-2.16)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 2.17Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.188 , 0.234 0.187 , 0.234	Depositor DCC
$R_{free}$ test set	10527 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.7	Xtrriage
Anisotropy	0.267	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	30710	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/1490	0.61	0/2019
1	B	0.38	0/1490	0.60	0/2019
1	C	0.35	0/1490	0.57	0/2019
1	D	0.34	0/1490	0.58	0/2019
1	E	0.38	0/1490	0.62	0/2019
1	F	0.40	0/1490	0.63	0/2019
1	G	0.43	0/1490	0.64	0/2019
1	H	0.40	0/1490	0.64	0/2019
1	I	0.40	0/1490	0.60	0/2019
1	J	0.38	0/1490	0.60	0/2019
1	K	0.42	0/1490	0.63	0/2019
1	L	0.43	0/1490	0.63	0/2019
1	M	0.35	0/1490	0.57	0/2019
1	N	0.35	0/1490	0.57	0/2019
1	O	0.42	0/1490	0.63	0/2019
1	P	0.43	0/1490	0.64	0/2019
1	Q	0.41	0/1490	0.63	0/2019
1	R	0.42	0/1490	0.63	0/2019
1	S	0.36	0/1490	0.57	0/2019
1	T	0.34	0/1490	0.56	0/2019
All	All	0.39	0/29800	0.61	0/40380

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1456	0	1418	43	0
1	B	1456	0	1418	50	0
1	C	1456	0	1418	47	0
1	D	1456	0	1418	37	0
1	E	1456	0	1418	46	0
1	F	1456	0	1418	40	0
1	G	1456	0	1418	31	0
1	H	1456	0	1418	30	0
1	I	1456	0	1418	36	0
1	J	1456	0	1418	36	0
1	K	1456	0	1418	39	0
1	L	1456	0	1418	42	0
1	M	1456	0	1418	53	0
1	N	1456	0	1418	42	0
1	O	1456	0	1418	36	0
1	P	1456	0	1418	33	0
1	Q	1456	0	1418	36	0
1	R	1456	0	1418	36	0
1	S	1456	0	1418	66	0
1	T	1456	0	1418	70	0
2	A	77	0	0	3	0
2	B	71	0	0	3	0
2	C	58	0	0	4	0
2	D	56	0	0	2	0
2	E	77	0	0	4	0
2	F	82	0	0	2	0
2	G	94	0	0	0	0
2	H	92	0	0	0	0
2	I	72	0	0	2	0
2	J	86	0	0	0	0
2	K	106	0	0	0	0
2	L	105	0	0	6	0
2	M	49	0	0	2	0
2	N	58	0	0	2	0
2	O	95	0	0	1	0
2	P	116	0	0	3	0
2	Q	104	0	0	2	0
2	R	93	0	0	4	0
2	S	52	0	0	4	0
2	T	47	0	0	2	0
All	All	30710	0	28360	720	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 720 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:180:LEU:O	1:J:79:LYS:HE3	1.66	0.96
1:S:109:ASN:HD21	1:S:118:ASP:HB2	1.32	0.94
1:B:180:LEU:O	1:D:79:LYS:HE2	1.69	0.92
1:Q:79:LYS:HE2	1:T:180:LEU:O	1.71	0.91
1:T:109:ASN:HD21	1:T:118:ASP:HB2	1.38	0.89

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	184/186 (99%)	177 (96%)	6 (3%)	1 (0%)	29 28
1	B	184/186 (99%)	179 (97%)	5 (3%)	0	100 100
1	C	184/186 (99%)	177 (96%)	6 (3%)	1 (0%)	29 28
1	D	184/186 (99%)	178 (97%)	6 (3%)	0	100 100
1	E	184/186 (99%)	178 (97%)	6 (3%)	0	100 100
1	F	184/186 (99%)	182 (99%)	2 (1%)	0	100 100
1	G	184/186 (99%)	180 (98%)	4 (2%)	0	100 100
1	H	184/186 (99%)	180 (98%)	4 (2%)	0	100 100
1	I	184/186 (99%)	177 (96%)	7 (4%)	0	100 100
1	J	184/186 (99%)	179 (97%)	5 (3%)	0	100 100
1	K	184/186 (99%)	180 (98%)	4 (2%)	0	100 100
1	L	184/186 (99%)	181 (98%)	3 (2%)	0	100 100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	M	184/186 (99%)	178 (97%)	4 (2%)	2 (1%)	14	11
1	N	184/186 (99%)	179 (97%)	4 (2%)	1 (0%)	29	28
1	O	184/186 (99%)	179 (97%)	5 (3%)	0	100	100
1	P	184/186 (99%)	180 (98%)	4 (2%)	0	100	100
1	Q	184/186 (99%)	181 (98%)	3 (2%)	0	100	100
1	R	184/186 (99%)	178 (97%)	6 (3%)	0	100	100
1	S	184/186 (99%)	178 (97%)	6 (3%)	0	100	100
1	T	184/186 (99%)	176 (96%)	8 (4%)	0	100	100
All	All	3680/3720 (99%)	3577 (97%)	98 (3%)	5 (0%)	51	58

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	182	LEU
1	M	17	ASN
1	C	39	PRO
1	M	39	PRO
1	N	39	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/154 (100%)	149 (97%)	5 (3%)	39	47
1	B	154/154 (100%)	150 (97%)	4 (3%)	46	55
1	C	154/154 (100%)	152 (99%)	2 (1%)	69	79
1	D	154/154 (100%)	149 (97%)	5 (3%)	39	47
1	E	154/154 (100%)	149 (97%)	5 (3%)	39	47
1	F	154/154 (100%)	152 (99%)	2 (1%)	69	79
1	G	154/154 (100%)	151 (98%)	3 (2%)	57	68

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	154/154 (100%)	151 (98%)	3 (2%)	57	68
1	I	154/154 (100%)	149 (97%)	5 (3%)	39	47
1	J	154/154 (100%)	152 (99%)	2 (1%)	69	79
1	K	154/154 (100%)	151 (98%)	3 (2%)	57	68
1	L	154/154 (100%)	150 (97%)	4 (3%)	46	55
1	M	154/154 (100%)	149 (97%)	5 (3%)	39	47
1	N	154/154 (100%)	149 (97%)	5 (3%)	39	47
1	O	154/154 (100%)	153 (99%)	1 (1%)	86	92
1	P	154/154 (100%)	151 (98%)	3 (2%)	57	68
1	Q	154/154 (100%)	152 (99%)	2 (1%)	69	79
1	R	154/154 (100%)	149 (97%)	5 (3%)	39	47
1	S	154/154 (100%)	151 (98%)	3 (2%)	57	68
1	T	154/154 (100%)	152 (99%)	2 (1%)	69	79
All	All	3080/3080 (100%)	3011 (98%)	69 (2%)	52	62

5 of 69 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Q	103	LEU
1	R	87	THR
1	S	79	LYS
1	G	103	LEU
1	G	87	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 112 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	78	HIS
1	T	155	GLN
1	M	127	GLN
1	T	109	ASN
1	R	109	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	186/186 (100%)	-0.12	9 (4%) 30 32	27, 41, 85, 124	0
1	B	186/186 (100%)	-0.10	3 (1%) 72 72	28, 47, 73, 92	0
1	C	186/186 (100%)	0.11	10 (5%) 25 27	32, 49, 85, 109	0
1	D	186/186 (100%)	0.11	10 (5%) 25 27	30, 53, 94, 116	0
1	E	186/186 (100%)	-0.11	2 (1%) 80 80	27, 46, 68, 95	0
1	F	186/186 (100%)	-0.17	9 (4%) 30 32	23, 37, 83, 113	0
1	G	186/186 (100%)	-0.32	4 (2%) 62 62	24, 35, 63, 87	0
1	H	186/186 (100%)	-0.38	0 100 100	24, 36, 62, 82	0
1	I	186/186 (100%)	-0.07	7 (3%) 40 41	28, 43, 84, 106	0
1	J	186/186 (100%)	-0.23	5 (2%) 54 55	25, 39, 74, 100	0
1	K	186/186 (100%)	-0.34	0 100 100	22, 35, 60, 86	0
1	L	186/186 (100%)	-0.33	3 (1%) 72 72	23, 35, 66, 89	0
1	M	186/186 (100%)	0.26	7 (3%) 40 41	38, 58, 83, 95	0
1	N	186/186 (100%)	0.16	7 (3%) 40 41	35, 56, 87, 110	0
1	O	186/186 (100%)	-0.21	1 (0%) 91 91	22, 40, 67, 80	0
1	P	186/186 (100%)	-0.37	4 (2%) 62 62	21, 33, 64, 92	0
1	Q	186/186 (100%)	-0.34	3 (1%) 72 72	23, 35, 64, 87	0
1	R	186/186 (100%)	-0.34	1 (0%) 91 91	24, 38, 63, 85	0
1	S	186/186 (100%)	0.28	14 (7%) 14 15	33, 51, 109, 121	0
1	T	186/186 (100%)	0.33	12 (6%) 18 19	40, 60, 96, 123	0
All	All	3720/3720 (100%)	-0.11	111 (2%) 50 51	21, 44, 79, 124	0

The worst 5 of 111 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	S	186	ILE	6.0
1	D	186	ILE	5.9
1	A	182	LEU	5.7
1	N	180	LEU	5.7
1	A	181	ASP	5.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.