



wwPDB NMR Structure Validation Summary Report ⓘ

Aug 10, 2020 – 03:30 AM BST

PDB ID : 1NXN
Title : SOLUTION STRUCTURE OF CONTRYPHAN-VN
Authors : Eliseo, T.; Cicero, D.O.; Polticelli, F.; Schinina, M.E.; Massilia, G.R.; Paci, M.; Ascenzi, P.
Deposited on : 2003-02-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.13.1
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | NMR archive (#Entries) |
|-----------------------|--------------------------|------------------------|
| Clashscore | 158937 | 12864 |
| Ramachandran outliers | 154571 | 11451 |
| Sidechain outliers | 154315 | 11428 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 10 | |

2 Ensemble composition and analysis

This entry contains 20 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores.

Cyrange was unable to find well-defined residues.

Error message: The number of core atoms (6) was below the domain threshold value (8).

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 142 atoms, of which 66 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CONTRYPHAN-VN, MAJOR FORM (CIS CONFORMER).

| Mol | Chain | Residues | Atoms | | | | | | Trace |
|-----|-------|----------|-------|----|----|----|----|---|-------|
| | | | Total | C | H | N | O | S | |
| 1 | A | 10 | 142 | 50 | 66 | 13 | 11 | 2 | 1 |

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: CONTRYPHAN-VN, MAJOR FORM (CIS CONFORMER)



4.2 Residue scores for the representative (author defined) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: CONTRYPHAN-VN, MAJOR FORM (CIS CONFORMER)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *FEWEST RESTRAINTS VIOLATIONS, LOWEST ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---------------|--------------------|---------|
| XwinNMR | structure solution | |
| NMRPipe | structure solution | |
| NMRView | structure solution | |
| XPLOR | structure solution | 1.5 |
| XPLOR | refinement | 1.5 |

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: DTR, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------------------|-------------|---------------------|
| | | RMSZ | #Z>5 | RMSZ | #Z>5 |
| 1 | A | 1.24±1.38 | 1±4/63 (1.3± 5.9%) | 1.41±0.65 | 0±2/84 (0.4± 1.8%) |
| All | All | 1.85 | 17/1260 (1.3%) | 1.56 | 7/1680 (0.4%) |

5 of 17 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) | Models | |
|-----|-------|-----|------|-------|--------|-------------|----------|--------|-------|
| | | | | | | | | Worst | Total |
| 1 | A | 9 | CYS | CA-CB | -24.64 | 0.99 | 1.53 | 1 | 1 |
| 1 | A | 2 | ASP | CA-CB | -19.88 | 1.10 | 1.53 | 1 | 1 |
| 1 | A | 6 | LYS | CE-NZ | -18.36 | 1.03 | 1.49 | 1 | 1 |
| 1 | A | 2 | ASP | N-CA | -17.66 | 1.11 | 1.46 | 1 | 1 |
| 1 | A | 6 | LYS | CG-CD | -17.39 | 0.93 | 1.52 | 1 | 1 |

5 of 7 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) | Models | |
|-----|-------|-----|------|----------|-------|-------------|----------|--------|-------|
| | | | | | | | | Worst | Total |
| 1 | A | 6 | LYS | CB-CG-CD | 21.51 | 167.54 | 111.60 | 1 | 1 |
| 1 | A | 6 | LYS | CG-CD-CE | 20.97 | 174.81 | 111.90 | 1 | 1 |
| 1 | A | 6 | LYS | CD-CE-NZ | 10.88 | 136.73 | 111.70 | 1 | 1 |
| 1 | A | 9 | CYS | N-CA-CB | 10.14 | 128.85 | 110.60 | 1 | 1 |
| 1 | A | 9 | CYS | CA-CB-SG | 9.65 | 131.37 | 114.00 | 1 | 1 |

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|-----|-------|-------|----------|----------|---------|
| 1 | A | 76 | 65 | 64 | 3±4 |
| All | All | 1520 | 1319 | 1280 | 68 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

5 of 50 unique clashes are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|--------------|--------------|----------|-------------|--------|-------|
| | | | | Worst | Total |
| 1:A:6:LYS:CG | 1:A:6:LYS:CE | 1.20 | 2.18 | 1 | 1 |
| 1:A:6:LYS:CB | 1:A:6:LYS:CD | 1.07 | 2.32 | 1 | 1 |
| 1:A:6:LYS:CG | 1:A:6:LYS:CD | 0.90 | 0.93 | 1 | 1 |
| 1:A:1:GLY:O | 1:A:3:CYS:N | 0.89 | 2.05 | 17 | 1 |
| 1:A:3:CYS:SG | 1:A:9:CYS:CB | 0.89 | 2.59 | 1 | 1 |

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|--------------|-------------|-------------|-------------|----|
| 1 | A | 7/10 (70%) | 5±1 (70±13%) | 2±1 (23±8%) | 1±1 (7±12%) | 2 | 16 |
| All | All | 140/200 (70%) | 98 (70%) | 32 (23%) | 10 (7%) | 2 | 16 |

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | A | 2 | ASP | 4 |
| 1 | A | 6 | LYS | 3 |
| 1 | A | 9 | CYS | 3 |

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|-----|-------|----------------|--------------|--------------|-------------|
| 1 | A | 7/7 (100%) | 6±1 (83±10%) | 1±1 (17±10%) | 5 40 |
| All | All | 140/140 (100%) | 116 (83%) | 24 (17%) | 5 40 |

All 4 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|-----|-------|-----|------|----------------|
| 1 | A | 6 | LYS | 9 |
| 1 | A | 3 | CYS | 7 |
| 1 | A | 9 | CYS | 4 |
| 1 | A | 2 | ASP | 4 |

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

| Mol | Chain | Number of breaks |
|-----|-------|------------------|
| 1 | 1-A | 2 |

All chain breaks are listed below:

| Model | Chain | Residue-1 | Atom-1 | Residue-2 | Atom-2 | Distance (Å) |
|-------|-------|-----------|--------|-----------|--------|--------------|
| 1 | A | 8:TRP | C | 9:CYS | N | 1.16 |
| 1 | A | 5:DTR | C | 6:LYS | N | 1.06 |

7 Chemical shift validation

No chemical shift data were provided