

Full wwPDB X-ray Structure Validation Report (i)

Aug 8, 2023 – 01:02 AM EDT

PDB ID : 1P63

Title : Human Acidic Fibroblast Growth Factor. 140 Amino Acid Form with Amino

Terminal His Tag and Leu111 Replaced with Ile (L111I)

Authors: Brych, S.R.; Kim, J.; Logan, T.M.; Blaber, M.

Deposited on : 2003-04-28

Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution		
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(\mathring{A}))$		
Clashscore	141614	3665 (1.60-1.60)		
Ramachandran outliers	138981	3564 (1.60-1.60)		
Sidechain outliers	138945	3563 (1.60-1.60)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain						
1	A	144	79%	17%					
1	В	144	70%	26%					



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ACIDIC FIBROBLAST GROWTH FACTOR.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	140	Total	С	N	О	S	0	0	0
1	1 A	140	1126	714	200	208	4	U	U	U
1	B	141	Total	С	N	О	S	0	0	0
1	1 B	141	1136	720	203	209	4		U	U

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	HIS	-	expression tag	UNP P05230
A	1B	HIS	-	expression tag	UNP P05230
A	1C	HIS	-	expression tag	UNP P05230
A	1D	HIS	-	expression tag	UNP P05230
A	111	ILE	LEU	engineered mutation	UNP P05230
В	1A	HIS	-	expression tag	UNP P05230
В	1B	HIS	-	expression tag	UNP P05230
В	1C	HIS	-	expression tag	UNP P05230
В	1D	HIS	-	expression tag	UNP P05230
В	111	ILE	LEU	engineered mutation	UNP P05230


• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 5	O 4	S 1	0	0

 \bullet Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: $\mathrm{CH_2O_2}).$



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	В	1	Total 3	C 1	O 2	0	0

• Molecule 4 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	118	Total O 118 118	0	0
4	В	101	Total O 101 101	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	C 2 2 21	Depositor	
Cell constants	74.41Å 96.72Å 107.79Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	26.90 - 1.60	Depositor	
% Data completeness	(Not available) (26.90-1.60)	Depositor	
(in resolution range)	(1000 available) (20.30 1.00)		
R_{merge}	0.05	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	TNT	Depositor	
R, R_{free}	0.219 , 0.251	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2489	wwPDB-VP	
Average B, all atoms (Å ²)	24.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
IVIOI	Cham	RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.87	5/1156 (0.4%)	1.06	10/1562 (0.6%)	
1	В	0.87	8/1167 (0.7%)	1.04	11/1577 (0.7%)	
All	All	0.87	$13/2323 \ (0.6\%)$	1.05	21/3139 (0.7%)	

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	В	82	GLU	CD-OE2	6.86	1.33	1.25
1	A	104	GLU	CD-OE2	5.70	1.31	1.25
1	A	91	GLU	CD-OE2	5.67	1.31	1.25
1	A	82	GLU	CD-OE2	5.67	1.31	1.25
1	A	60	GLU	CD-OE2	5.65	1.31	1.25
1	A	49	GLU	CD-OE2	5.63	1.31	1.25
1	В	91	GLU	CD-OE2	5.50	1.31	1.25
1	В	104	GLU	CD-OE2	5.46	1.31	1.25
1	В	87	GLU	CD-OE2	5.45	1.31	1.25
1	В	81	GLU	CD-OE2	5.25	1.31	1.25
1	В	90	GLU	CD-OE1	5.23	1.31	1.25
1	В	60	GLU	CD-OE2	5.15	1.31	1.25
1	В	49	GLU	CD-OE1	5.10	1.31	1.25

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
1	A	68	ASP	CB-CG-OD2	-7.40	111.64	118.30
1	A	36	ASP	CB-CG-OD2	-7.02	111.98	118.30
1	A	36	ASP	CB-CG-OD1	6.93	124.54	118.30
1	В	36	ASP	CB-CG-OD2	-6.80	112.18	118.30
1	В	36	ASP	CB-CG-OD1	6.71	124.34	118.30
1	A	28	ASP	CB-CG-OD2	-6.45	112.50	118.30

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	A	68	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	32	ASP	CB-CG-OD1	6.23	123.91	118.30
1	A	39	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	В	28	ASP	CB-CG-OD2	-6.15	112.76	118.30
1	В	68	ASP	CB-CG-OD2	-6.07	112.84	118.30
1	В	32	ASP	CB-CG-OD1	5.97	123.67	118.30
1	A	32	ASP	CB-CG-OD2	-5.91	112.98	118.30
1	В	32	ASP	CB-CG-OD2	-5.79	113.08	118.30
1	A	28	ASP	CB-CG-OD1	5.69	123.42	118.30
1	В	68	ASP	CB-CG-OD1	5.67	123.40	118.30
1	В	39	ASP	CB-CG-OD2	-5.67	113.20	118.30
1	В	70	ASP	CB-CG-OD2	-5.19	113.63	118.30
1	В	28	ASP	CB-CG-OD1	5.17	122.95	118.30
1	В	94	TYR	CB-CG-CD2	-5.04	117.97	121.00
1	A	39	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1126	0	1102	14	0
1	В	1136	0	1109	21	0
2	A	5	0	0	0	0
3	В	3	0	1	1	0
4	A	118	0	0	2	0
4	В	101	0	0	2	1
All	All	2489	0	2212	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
	1100111-2	$\operatorname{distance}\left(\mathrm{\AA}\right)$	overlap (Å)
1:A:49:GLU:HG3	1:A:84:LEU:HD21	1.44	0.97
1:A:49:GLU:HG3	1:A:84:LEU:CD2	2.15	0.77
1:B:18:ASN:HB2	1:B:129:ALA:HA	1.71	0.73
1:B:39:ASP:O	1:B:43:GLN:NE2	2.30	0.64
1:B:72:LEU:HD13	1:B:119:ARG:HG2	1.86	0.57
1:A:81:GLU:HG3	1:A:101:LYS:HD2	1.86	0.57
1:B:8:TYR:HB3	1:B:46:LEU:O	2.04	0.57
1:B:72:LEU:CD1	1:B:119:ARG:HG2	2.35	0.56
1:A:124:HIS:CD2	1:A:127:GLN:HG3	2.41	0.55
1:B:10:LYS:HB3	1:B:11:PRO:HD2	1.88	0.55
1:B:119:ARG:NH1	4:B:2420:HOH:O	2.38	0.55
1:B:7:ASN:HB2	4:B:2250:HOH:O	2.10	0.51
1:A:118:LYS:HE2	1:A:127:GLN:HE22	1.76	0.50
1:A:81:GLU:CD	1:A:101:LYS:HD2	2.31	0.49
1:A:10:LYS:HB3	1:A:11:PRO:HD2	1.94	0.49
1:A:18:ASN:HB2	1:A:129:ALA:HA	1.95	0.49
1:A:89:LEU:HD21	1:A:92:ASN:HA	1.93	0.49
1:B:37:ARG:NH2	1:B:136:PRO:O	2.46	0.49
1:B:72:LEU:HD22	1:B:72:LEU:N	2.27	0.49
1:A:81:GLU:CG	1:A:101:LYS:HD2	2.44	0.47
1:B:122:ARG:HA	1:B:122:ARG:HD3	1.59	0.46
1:B:134:PRO:O	1:B:135:LEU:HD23	2.16	0.45
1:B:40:GLN:O	1:B:43:GLN:OE1	2.35	0.45
1:A:57:LYS:CE	1:A:64:TYR:CZ	3.00	0.45
1:A:81:GLU:HG2	4:A:1337:HOH:O	2.17	0.45
1:B:43:GLN:H	1:B:43:GLN:CD	2.20	0.44
1:B:1(B):HIS:H	3:B:160:FMT:C	2.30	0.44
1:B:72:LEU:N	1:B:72:LEU:CD2	2.80	0.44
1:B:24:ARG:NH1	1:B:39:ASP:OD2	2.45	0.43
1:B:1(D):HIS:CE1	1:B:88:ARG:HB3	2.54	0.42
1:B:128:LYS:HA	1:B:128:LYS:HD3	1.81	0.42
1:A:98:ILE:HD11	4:A:1278:HOH:O	2.21	0.41
1:B:105:LYS:HD3	1:B:107:TRP:CZ2	2.56	0.41
1:B:121:PRO:O	1:B:122:ARG:NH1	2.54	0.41
1:A:128:LYS:HA	1:A:128:LYS:HD3	1.74	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)	
4:B:2403:HOH:O	4:B:2419:HOH:O[3_556]	2.00	0.20	



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	138/144 (96%)	132 (96%)	6 (4%)	0	100	100
1	В	139/144~(96%)	133 (96%)	6 (4%)	0	100	100
All	All	277/288 (96%)	265 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	123/127 (97%)	123 (100%)	0	100	100	
1	В	124/127~(98%)	121 (98%)	3 (2%)	49	24	
All	All	$247/254 \ (97\%)$	244 (99%)	3 (1%)	71	54	

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	37	ARG
1	В	118	LYS
1	В	122	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	Res	Type
1	В	21	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	${ m gths}$	В	ond ang	gles
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	162	-	4,4,4	0.43	0	6,6,6	0.12	0
3	FMT	В	160	-	2,2,2	0.45	0	1,1,1	0.28	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	160	FMT	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

