



wwPDB X-ray Structure Validation Summary Report

Aug 15, 2023 – 10:43 PM EDT

PDB ID : 1TWI
Title : Crystal structure of Diaminopimelate Decarboxylase from *m. jannaschii* in co-complex with L-lysine
Authors : Rajashankar, K.R.; Ray, S.S.; Bonanno, J.B.; Pinho, M.G.; He, G.; De Lencastre, H.; Tomasz, A.; Burley, S.K.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2004-07-01
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	434	 85% 13% 2% 2% 3%
1	B	434	 82% 16% 2% 2% 3%
1	C	434	 86% 13% 2% 2% 2%
1	D	434	 85% 14% 2% 2% 3%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diaminopimelate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	434	3395	2166	568	640	21	0	0	0
1	B	434	3395	2166	568	640	21	0	0	0
1	C	434	3395	2166	568	640	21	0	0	0
1	D	434	3395	2166	568	640	21	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	15	MET	-	initiating methionine	UNP Q58497
B	15	MET	-	initiating methionine	UNP Q58497
C	15	MET	-	initiating methionine	UNP Q58497
D	15	MET	-	initiating methionine	UNP Q58497

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	B	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	C	1	Total	C	N	O	P	0	0
			16	8	1	6	1		
2	D	1	Total	C	N	O	P	0	0
			16	8	1	6	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is LYSINE (three-letter code: LYS) (formula: C₆H₁₅N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			10	6	2	2		
4	D	1	Total	C	N	O	0	0
			10	6	2	2		
4	D	1	Total	C	N	O	0	0
			10	6	2	2		
4	D	1	Total	C	N	O	0	0
			10	6	2	2		
4	D	1	Total	C	N	O	0	0
			10	6	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	369	Total	O	0	0
			369	369		
5	B	301	Total	O	0	0
			301	301		
5	C	370	Total	O	0	0
			370	370		
5	D	383	Total	O	0	0
			383	383		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Diaminopimelate decarboxylase




- Molecule 1: Diaminopimelate decarboxylase




- Molecule 1: Diaminopimelate decarboxylase



- Molecule 1: Diaminopimelate decarboxylase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.24Å 147.07Å 89.39Å 90.00° 97.25° 90.00°	Depositor
Resolution (Å)	29.25 – 2.00 29.25 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.3 (29.25-2.00) 98.5 (29.25-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.56 (at 2.00Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.157 , 0.202 0.151 , 0.195	Depositor DCC
R_{free} test set	6043 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtrriage
Anisotropy	0.183	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 61.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15128	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	2/3452 (0.1%)	0.91	4/4654 (0.1%)
1	B	0.90	4/3452 (0.1%)	0.88	2/4654 (0.0%)
1	C	0.95	0/3452	0.89	0/4654
1	D	0.96	2/3452 (0.1%)	0.92	4/4654 (0.1%)
All	All	0.94	8/13808 (0.1%)	0.90	10/18616 (0.1%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	69	GLU	CG-CD	6.47	1.61	1.51
1	D	372	CYS	CB-SG	6.32	1.93	1.82
1	B	132	GLU	CD-OE1	5.40	1.31	1.25
1	A	132	GLU	CD-OE1	5.33	1.31	1.25
1	B	47	TYR	CD2-CE2	5.32	1.47	1.39

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	343	ARG	NE-CZ-NH1	6.82	123.71	120.30
1	A	92	ARG	NE-CZ-NH1	6.51	123.56	120.30
1	D	307	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	A	92	ARG	NE-CZ-NH2	-5.91	117.34	120.30
1	D	307	ARG	NE-CZ-NH2	-5.90	117.35	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3395	0	3457	56	0
1	B	3395	0	3457	67	0
1	C	3395	0	3457	54	0
1	D	3395	0	3457	54	0
2	A	16	0	7	2	0
2	B	16	0	7	2	0
2	C	16	0	8	1	0
2	D	16	0	7	2	0
3	C	1	0	0	0	0
4	D	60	0	72	18	0
5	A	369	0	0	10	0
5	B	301	0	0	11	0
5	C	370	0	0	9	0
5	D	383	0	0	9	0
All	All	15128	0	13929	219	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 219 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:270:TYR:HB2	1:C:273:LYS:HE2	1.48	0.96
1:B:430:GLU:HG2	5:B:1559:HOH:O	1.76	0.86
1:A:407:ASN:HD22	1:A:409:TYR:H	1.24	0.85
1:C:414:ARG:HH12	1:D:408:ASN:HD22	1.25	0.85
1:B:47:TYR:HB2	1:B:417:MET:HG2	1.57	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	432/434 (100%)	425 (98%)	6 (1%)	1 (0%)	47	44
1	B	432/434 (100%)	422 (98%)	9 (2%)	1 (0%)	47	44
1	C	432/434 (100%)	425 (98%)	6 (1%)	1 (0%)	47	44
1	D	432/434 (100%)	423 (98%)	8 (2%)	1 (0%)	47	44
All	All	1728/1736 (100%)	1695 (98%)	29 (2%)	4 (0%)	47	44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	D	85	ASN
1	C	85	ASN
1	B	85	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/367 (100%)	355 (97%)	12 (3%)	38	37
1	B	367/367 (100%)	355 (97%)	12 (3%)	38	37
1	C	367/367 (100%)	356 (97%)	11 (3%)	41	41
1	D	367/367 (100%)	354 (96%)	13 (4%)	36	35
All	All	1468/1468 (100%)	1420 (97%)	48 (3%)	36	37

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	176	ASN
1	D	83	LYS
1	C	299	PRO
1	C	410	ASN
1	D	116	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	408	ASN
1	D	228	GLN
1	D	18	ASN
1	D	141	ASN
1	D	352	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	LYS	D	1601	-	8,9,9	1.12	1 (12%)	9,10,10	0.93	0
4	LYS	D	1602	-	8,9,9	0.88	0	9,10,10	0.75	0
4	LYS	D	1605	-	8,9,9	0.69	0	9,10,10	0.90	0
2	PLP	B	1502	-	16,16,16	1.61	1 (6%)	20,23,23	2.14	5 (25%)
4	LYS	D	1604	-	8,9,9	0.97	1 (12%)	9,10,10	0.82	0
4	LYS	D	1606	-	8,9,9	0.79	0	9,10,10	0.98	0
2	PLP	A	1501	-	16,16,16	1.88	6 (37%)	20,23,23	2.04	4 (20%)
4	LYS	D	1603	2	8,9,9	0.94	0	9,10,10	0.54	0
2	PLP	D	1504	-	16,16,16	2.87	7 (43%)	20,23,23	2.25	5 (25%)
2	PLP	C	1503	4	16,16,16	2.44	5 (31%)	20,23,23	2.01	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LYS	D	1601	-	-	2/9/9/9	-
4	LYS	D	1602	-	-	2/9/9/9	-
4	LYS	D	1605	-	-	2/9/9/9	-
2	PLP	B	1502	-	-	0/8/8/8	0/1/1/1
4	LYS	D	1604	-	-	1/9/9/9	-
4	LYS	D	1606	-	-	3/9/9/9	-
2	PLP	A	1501	-	-	0/8/8/8	0/1/1/1
4	LYS	D	1603	2	-	2/9/9/9	-
2	PLP	D	1504	-	-	0/8/8/8	0/1/1/1
2	PLP	C	1503	4	-	0/8/8/8	0/1/1/1

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1504	PLP	O4A-C4A	-7.31	0.96	1.21
2	C	1503	PLP	C4-C4A	4.77	1.57	1.46
2	D	1504	PLP	C4-C4A	4.75	1.57	1.46
2	B	1502	PLP	C4-C4A	4.62	1.57	1.46
2	A	1501	PLP	C4-C4A	4.60	1.57	1.46

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	1504	PLP	C4-C3-C2	7.42	124.78	120.19
2	C	1503	PLP	C4-C3-C2	6.47	124.19	120.19
2	A	1501	PLP	C4-C3-C2	5.76	123.75	120.19
2	B	1502	PLP	C2A-C2-C3	5.72	127.96	120.89
2	B	1502	PLP	C4-C3-C2	4.55	123.01	120.19

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	1606	LYS	N-CA-CB-CG
4	D	1602	LYS	CG-CD-CE-NZ
4	D	1601	LYS	CE-CD-CG-CB
4	D	1605	LYS	CE-CD-CG-CB
4	D	1603	LYS	CA-CB-CG-CD

There are no ring outliers.

10 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1601	LYS	4	0
4	D	1602	LYS	5	0
4	D	1605	LYS	4	0
2	B	1502	PLP	2	0
4	D	1604	LYS	1	0
4	D	1606	LYS	3	0
2	A	1501	PLP	2	0
4	D	1603	LYS	1	0
2	D	1504	PLP	2	0
2	C	1503	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	434/434 (100%)	-0.63	0 100 100	8, 15, 27, 33	0
1	B	434/434 (100%)	-0.33	15 (3%) 44 43	9, 18, 36, 42	0
1	C	434/434 (100%)	-0.58	9 (2%) 63 62	8, 15, 30, 40	0
1	D	434/434 (100%)	-0.54	11 (2%) 57 56	8, 15, 31, 39	0
All	All	1736/1736 (100%)	-0.52	35 (2%) 65 63	8, 15, 32, 42	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	178	ASN	4.0
1	B	180	LYS	3.8
1	C	181	THR	3.6
1	B	181	THR	3.4
1	B	360	ASN	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	LYS	D	1606	10/10	0.83	0.16	33,34,35,36	0
4	LYS	D	1602	10/10	0.84	0.20	33,34,35,36	0
4	LYS	D	1601	10/10	0.85	0.20	34,35,37,42	0
4	LYS	D	1604	10/10	0.87	0.13	42,47,49,49	0
4	LYS	D	1605	10/10	0.90	0.15	25,28,38,39	0
4	LYS	D	1603	10/10	0.92	0.13	24,27,35,36	0
2	PLP	A	1501	16/16	0.97	0.10	9,12,15,31	0
2	PLP	B	1502	16/16	0.97	0.11	15,17,20,30	0
2	PLP	C	1503	16/16	0.97	0.09	11,12,17,26	0
2	PLP	D	1504	16/16	0.98	0.08	9,11,16,27	0
3	MG	C	1607	1/1	0.98	0.02	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.