

# Full wwPDB X-ray Structure Validation Report (i)

#### Aug 15, 2023 – 08:17 PM EDT

PDB ID : 1U4R

Title: Crystal Structure of human RANTES mutant 44-AANA-47

Authors: Shaw, J.P.; Johnson, Z.; Borlat, F.; Zwahlen, C.; Kungl, A.; Roulin, K.;

Harrenga, A.; Wells, T.N.C.; Proudfoot, A.E.I.

Deposited on : 2004-07-26

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	A	68	68% 19%		9%		
1	В	68	62% 359	)	•		
1	С	68	71% 2	l%	• 6%		
1	D	68	82%	189	%		



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2227 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Small inducible cytokine A5.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	62	Total	С	N	О	S	0	0	0
1	A	02	482	310	82	85	5	0	U	U
1	В	68	Total	С	N	О	S	0	0	0
1	Ъ	00	522	335	84	98	5		U	U
1	С	64	Total	С	N	О	S	0	0	0
1		04	494	316	82	91	5	0	U	U
1	1 D	68	Total	С	N	О	S	0	0	0
1	ט	00	521	335	85	96	5		U	U

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	ALA	ARG	engineered mutation	UNP P13501
A	45	ALA	LYS	engineered mutation	UNP P13501
A	47	ALA	ARG	engineered mutation	UNP P13501
В	44	ALA	ARG	engineered mutation	UNP P13501
В	45	ALA	LYS	engineered mutation	UNP P13501
В	47	ALA	ARG	engineered mutation	UNP P13501
С	44	ALA	ARG	engineered mutation	UNP P13501
С	45	ALA	LYS	engineered mutation	UNP P13501
С	47	ALA	ARG	engineered mutation	UNP P13501
D	44	ALA	ARG	engineered mutation	UNP P13501
D	45	ALA	LYS	engineered mutation	UNP P13501
D	47	ALA	ARG	engineered mutation	UNP P13501

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	С	1	Total O S 5 4 1	0	0

### • Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	41	Total O 41 41	0	0
3	В	59	Total O 59 59	0	0
3	С	42	Total O 42 42	0	0
3	D	51	Total O 51 51	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Small inducible cytokine A5





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	22.82Å 80.47Å 65.44Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $94.52^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	15.63 - 2.20	Depositor	
% Data completeness	93.8 (15.63-2.20)	Depositor	
(in resolution range)	35.0 (15.05 2.20)	Depositor	
$R_{merge}$	0.10	Depositor	
$R_{sym}$	(Not available)	Depositor	
Refinement program	CNX 2002	Depositor	
$R, R_{free}$	0.222 , $0.306$	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2227	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP	



## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
IVIOI	Chain	RMSZ	RMSZ $\# Z  > 5$		# Z  > 5	
1	A	0.77	2/495~(0.4%)	0.87	5/673 (0.7%)	
1	В	0.32	0/537	0.57	0/733	
1	С	0.33	0/508	0.59	0/692	
1	D	0.33	0/537	0.60	0/733	
All	All	0.47	$2/2077 \ (0.1\%)$	0.66	5/2831 (0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

#### All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
1	A	42	VAL	C-N	12.60	1.63	1.34
1	A	46	ASN	C-N	8.38	1.53	1.34

#### All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	A	42	VAL	C-N-CA	-9.52	97.91	121.70
1	A	46	ASN	C-N-CA	-6.81	104.67	121.70
1	A	42	VAL	O-C-N	6.36	132.88	122.70
1	A	42	VAL	CA-C-N	-5.66	104.74	117.20
1	A	43	THR	N-CA-C	-5.26	96.79	111.00

There are no chirality outliers.



All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	46	ASN	Mainchain

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	482	0	463	15	0
1	В	522	0	493	19	0
1	С	494	0	471	14	0
1	D	521	0	493	11	0
2	A	10	0	0	1	0
2	С	5	0	0	1	0
3	A	41	0	0	1	0
3	В	59	0	0	1	0
3	С	42	0	0	3	0
3	D	51	0	0	4	0
All	All	2227	0	1920	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance}  (\mathring{\rm A}) \end{array}$	Clash overlap (Å)
1:B:25:LYS:HE3	1:B:44:ALA:HA	1.68	0.74
1:D:25:LYS:HE3	1:D:44:ALA:HA	1.73	0.70
1:A:43:THR:C	1:A:46:ASN:N	2.46	0.69
1:D:67:MET:HE1	3:D:119:HOH:O	1.93	0.67
1:B:15:ILE:HD13	1:B:19:LEU:HD13	1.78	0.66
1:C:15:ILE:HD13	1:C:16:ALA:N	2.10	0.66
1:A:19:LEU:HD12	1:A:24:ILE:HD11	1.78	0.65
1:C:14:TYR:CD2	1:C:52:ASN:HB2	2.32	0.64
1:D:1:SER:HA	3:D:93:HOH:O	1.98	0.63
1:A:55:LYS:O	1:A:59:ARG:HD3	2.00	0.60
1:A:56:LYS:HA	1:A:59:ARG:HH11	1.67	0.59

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Continued from pres		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ ({\rm \AA})$	overlap (Å)	
1:A:15:ILE:HD11	1:B:3:TYR:HE2	1.69	0.58	
1:D:67:MET:O	1:D:68:SER:CB	2.51	0.57	
1:A:56:LYS:HA	1:A:59:ARG:NH1	2.19	0.57	
1:B:52:ASN:HB3	1:B:55:LYS:HG2	1.86	0.57	
1:A:21:ARG:HH11	1:A:21:ARG:HG3	1.72	0.55	
1:C:13:ALA:HB1	3:C:204:HOH:O	2.06	0.54	
1:C:55:LYS:O	1:C:59:ARG:HD3	2.07	0.53	
1:B:55:LYS:O	1:B:59:ARG:HD3	2.09	0.52	
1:A:63:ASN:HB3	1:A:67:MET:CE	2.40	0.52	
1:B:14:TYR:CD2	1:B:52:ASN:HB2	2.45	0.52	
1:B:25:LYS:O	1:B:26:GLU:HG2	2.10	0.52	
1:B:21:ARG:HD3	1:B:61:TYR:CE1	2.45	0.51	
1:D:63:ASN:O	1:D:67:MET:HG3	2.10	0.51	
1:B:63:ASN:O	1:B:67:MET:HG3	2.11	0.51	
2:A:203:SO4:O4	1:B:2:PRO:HD2	2.11	0.50	
1:D:67:MET:HE2	3:D:89:HOH:O	2.13	0.49	
1:D:55:LYS:O	1:D:59:ARG:HD3	2.12	0.49	
1:A:59:ARG:HH11	1:A:59:ARG:HB2	1.77	0.49	
1:A:59:ARG:NH1	1:A:59:ARG:HB2	2.28	0.49	
1:A:36:ASN:ND2	3:A:225:HOH:O	2.46	0.48	
1:C:14:TYR:CG	1:C:52:ASN:HB2	2.49	0.47	
1:D:43:THR:OG1	1:D:47:ALA:HB3	2.15	0.46	
1:B:67:MET:O	1:B:68:SER:C	2.53	0.46	
1:B:27:TYR:HA	1:B:40:VAL:O	2.16	0.46	
1:D:67:MET:O	1:D:68:SER:HB2	2.15	0.45	
1:A:21:ARG:HG3	1:A:21:ARG:NH1	2.31	0.45	
1:B:27:TYR:CZ	1:B:62:ILE:HG23	2.51	0.45	
1:C:37:PRO:HA	3:C:234:HOH:O	2.16	0.45	
1:C:13:ALA:HB2	3:C:238:HOH:O	2.16	0.44	
1:A:15:ILE:HD12	1:A:15:ILE:HA	1.88	0.44	
1:A:24:ILE:HD12	1:A:41:PHE:HB3	1.98	0.44	
1:D:67:MET:CE	3:D:89:HOH:O	2.65	0.44	
1:B:23:HIS:O	1:B:43:THR:HA	2.18	0.43	
1:B:9:PRO:HG3	3:B:78:HOH:O	2.19	0.42	
1:B:52:ASN:HA	1:B:53:PRO:HD2	1.93	0.42	
1:C:27:TYR:CD1	1:C:27:TYR:C	2.93	0.42	
1:B:14:TYR:HE1	1:B:38:ALA:HB2	1.85	0.42	
1:D:39:VAL:HG23	1:D:53:PRO:HB3	2.02	0.42	
1:C:15:ILE:HD13	1:C:16:ALA:H	1.85	0.42	
1:A:38:ALA:HA	1:A:53:PRO:CG	2.49	0.42	
1:C:52:ASN:HA	1:C:53:PRO:HD2	1.94	0.42	

Continued on next page...



I 'omtamalod	trom	mmonia	maaa
Continued	11 0116	DICUIUUS	Daue
	.,	10	1

Atom-1	Atom-2	Interatomic	$\operatorname{Clash}$	
Atom-1	Atom-2	${\rm distance}  (\rm \AA)$	overlap (Å)	
1:C:38:ALA:HA	1:C:53:PRO:HD3	2.02	0.41	
1:C:23:HIS:O	1:C:43:THR:HA	2.19	0.41	
1:B:8:THR:HA	1:B:9:PRO:HD3	1.95	0.41	
1:B:58:VAL:O	1:B:62:ILE:HG13	2.21	0.41	
1:C:27:TYR:CE2	1:C:62:ILE:HG23	2.56	0.41	
1:C:56:LYS:HD2	2:C:202:SO4:O1	2.20	0.40	

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

## 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	58/68 (85%)	57 (98%)	1 (2%)	0	100	100
1	В	66/68 (97%)	62 (94%)	3 (4%)	1 (2%)	10	8
1	С	$62/68 \; (91\%)$	60 (97%)	1 (2%)	1 (2%)	9	7
1	D	66/68~(97%)	65 (98%)	1 (2%)	0	100	100
All	All	252/272~(93%)	244 (97%)	6 (2%)	2 (1%)	19	19

#### All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	23	HIS
1	С	16	ALA

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the	number o	of residues	for	which	the	${\rm sidechain}$	conformation	was
analysed, and the total number of	residues.							

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	52/59~(88%)	51 (98%)	1 (2%)	57 71
1	В	55/59~(93%)	54 (98%)	1 (2%)	59 72
1	$\mathbf{C}$	53/59 (90%)	51 (96%)	2 (4%)	33 42
1	D	55/59~(93%)	55 (100%)	0	100 100
All	All	215/236 (91%)	211 (98%)	4 (2%)	57 71

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	ARG
1	В	27	TYR
1	С	15	ILE
1	С	19	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	ASN
1	A	48	GLN
1	A	63	ASN
1	В	63	ASN
1	С	23	HIS
1	С	63	ASN

## 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Cha		Chain	Chain Res	Res Link	Bond lengths			Bond angles		
Mol Type Ch	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
2	SO4	С	202	-	4,4,4	0.20	0	6,6,6	0.15	0
2	SO4	A	201	-	4,4,4	0.23	0	6,6,6	0.09	0
2	SO4	A	203	-	4,4,4	0.25	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	202	SO4	1	0
2	A	203	SO4	1	0

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:



Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	42:VAL	С	43:THR	N	1.63



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

## 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

