

wwPDB X-ray Structure Validation Summary Report (i)

Aug 15, 2023 – 07:00 PM EDT

| PDB ID | : | 1XB0 |
|--------------|---|--|
| Title | : | Structure of the BIR domain of IAP-like protein 2 |
| Authors | : | Shin, H.; Renatus, M.; Eckelman, B.P.; Nunes, V.A.; Sampaio, C.A.M.; |
| | | Salvesen, G.S. |
| Deposited on | : | 2004-08-27 |
| Resolution | : | 2.20 Å(reported) |

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


| MolProbity | : | 4.02b-467 |
|--------------------------------|---|--|
| Xtriage (Phenix) | : | NOT EXECUTED |
| EDS | : | NOT EXECUTED |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.35 |

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | $egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$ | ${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$ |
|-----------------------|--|---|
| Clashscore | 141614 | 5594 (2.20-2.20) |
| Ramachandran outliers | 138981 | 5503 (2.20-2.20) |
| Sidechain outliers | 138945 | 5504 (2.20-2.20) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain | | | | | | | |
|-----|-------|--------|------------------|-----|-------|------|--|--|--|--|
| 1 | А | 108 | | 68% | 15% • | 15% | | | | |
| 1 | В | 108 | | 72% | 19% | • 6% | | | | |
| 1 | С | 108 | | 67% | 16% • | 16% | | | | |
| 1 | D | 108 | | 67% | 16% • | 16% | | | | |
| 1 | Е | 108 | | 72% | 11% • | 14% | | | | |
| 1 | F | 108 | | 73% | 19% | • 5% | | | | |
| 2 | G | 7 | 14% | 43% | 43% | | | | | |
| 2 | Н | 7 | 29% | 29% | 43% | | | | | |



| Mol | Chain | | | Quality of cha | ain |
|-----|-------|---|-----|----------------|-----|
| 2 | Ι | 7 | 14% | 43% | 43% |
| 2 | J | 7 | 29% | 43% | 29% |
| 2 | K | 7 | 43% | 14% | 43% |
| 2 | L | 7 | | 57% | 43% |



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

| Mol | Chain | Residues | | At | oms | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 1 | А | 92 | Total | С | Ν | 0 | S | 3 | 0 | 0 |
| | Л | 92 | 751 | 482 | 130 | 134 | 5 | 5 | 0 | 0 |
| 1 | В | 101 | Total | С | Ν | Ο | S | 34 | 0 | 0 |
| | D | 101 | 819 | 524 | 142 | 148 | 5 | 54 | 0 | 0 |
| 1 | С | 91 | Total | С | Ν | Ο | S | 18 | 0 | 0 |
| | | 91 | 746 | 479 | 129 | 133 | 5 | 10 | 0 | 0 |
| 1 | D | 91 | Total | С | Ν | 0 | S | 25 | 0 | 0 |
| | D | 91 | 746 | 479 | 129 | 133 | 5 | 20 | 0 | U |
| 1 | Е | 93 | Total | С | Ν | Ο | S | 33 | 0 | 0 |
| | Ľ | 90 | 757 | 485 | 131 | 136 | 5 | | 0 | 0 |
| 1 | F | 103 | Total | С | Ν | 0 | S | 47 | 0 | 0 |
| | | 105 | 832 | 531 | 145 | 151 | 5 | 41 | | 0 |

• Molecule 1 is a protein called Baculoviral IAP repeat-containing protein 8.

There are 78 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| А | 249 | GLY | - | cloning artifact | UNP Q96P09 |
| А | 250 | SER | - | cloning artifact | UNP Q96P09 |
| А | 251 | HIS | - | cloning artifact | UNP Q96P09 |
| А | 252 | MET | - | cloning artifact | UNP Q96P09 |
| А | 253 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| А | 254 | THR | - | SEE REMARK 999 | UNP Q96P09 |
| А | 255 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| А | 256 | LEU | - | SEE REMARK 999 | UNP Q96P09 |
| А | 257 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| А | 258 | ARG | - | SEE REMARK 999 | UNP Q96P09 |
| А | 259 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| А | 260 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| А | 261 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| В | 249 | GLY | - | cloning artifact | UNP Q96P09 |
| В | 250 | SER | - | cloning artifact | UNP Q96P09 |
| В | 251 | HIS | - | cloning artifact | UNP Q96P09 |
| В | 252 | MET | - | cloning artifact | UNP Q96P09 |



| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| В | 253 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| В | 254 | THR | - | SEE REMARK 999 | UNP Q96P09 |
| В | 255 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| В | 256 | LEU | - | SEE REMARK 999 | UNP Q96P09 |
| В | 257 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| В | 258 | ARG | - | SEE REMARK 999 | UNP Q96P09 |
| В | 259 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| В | 260 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| В | 261 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| С | 249 | GLY | - | cloning artifact | UNP Q96P09 |
| С | 250 | SER | - | cloning artifact | UNP Q96P09 |
| С | 251 | HIS | - | cloning artifact | UNP Q96P09 |
| С | 252 | MET | _ | cloning artifact | UNP Q96P09 |
| С | 253 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| С | 254 | THR | _ | SEE REMARK 999 | UNP Q96P09 |
| С | 255 | ASN | _ | SEE REMARK 999 | UNP Q96P09 |
| С | 256 | LEU | _ | SEE REMARK 999 | UNP Q96P09 |
| С | 257 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| С | 258 | ARG | - | SEE REMARK 999 | UNP Q96P09 |
| С | 259 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| С | 260 | PRO | _ | SEE REMARK 999 | UNP Q96P09 |
| С | 261 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| D | 249 | GLY | - | cloning artifact | UNP Q96P09 |
| D | 250 | SER | _ | cloning artifact | UNP Q96P09 |
| D | 251 | HIS | - | cloning artifact | UNP Q96P09 |
| D | 252 | MET | - | cloning artifact | UNP Q96P09 |
| D | 253 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| D | 254 | THR | - | SEE REMARK 999 | UNP Q96P09 |
| D | 255 | ASN | _ | SEE REMARK 999 | UNP Q96P09 |
| D | 256 | LEU | - | SEE REMARK 999 | UNP Q96P09 |
| D | 257 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| D | 258 | ARG | - | SEE REMARK 999 | UNP Q96P09 |
| D | 259 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| D | 260 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| D | 261 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 249 | GLY | - | cloning artifact | UNP Q96P09 |
| Е | 250 | SER | - | cloning artifact | UNP Q96P09 |
| Е | 251 | HIS | - | cloning artifact | UNP Q96P09 |
| Е | 252 | MET | - | cloning artifact | UNP Q96P09 |
| Е | 253 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 254 | THR | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 255 | ASN | _ | SEE REMARK 999 | UNP Q96P09 |



| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| E | 256 | LEU | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 257 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 258 | ARG | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 259 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 260 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| Е | 261 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| F | 249 | GLY | - | cloning artifact | UNP Q96P09 |
| F | 250 | SER | - | cloning artifact | UNP Q96P09 |
| F | 251 | HIS | - | cloning artifact | UNP Q96P09 |
| F | 252 | MET | - | cloning artifact | UNP Q96P09 |
| F | 253 | SER | - | SEE REMARK 999 | UNP Q96P09 |
| F | 254 | THR | - | SEE REMARK 999 | UNP Q96P09 |
| F | 255 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| F | 256 | LEU | - | SEE REMARK 999 | UNP Q96P09 |
| F | 257 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| F | 258 | ARG | - | SEE REMARK 999 | UNP Q96P09 |
| F | 259 | ASN | - | SEE REMARK 999 | UNP Q96P09 |
| F | 260 | PRO | - | SEE REMARK 999 | UNP Q96P09 |
| F | 261 | SER | - | SEE REMARK 999 | UNP Q96P09 |

• Molecule 2 is a protein called Diablo homolog, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------------|---------|--------|--------|---------|---------|-------|
| 2 | G | 4 | Total 27 | C 19 | N 4 | 0 4 | 0 | 0 | 0 |
| 2 | Н | 4 | Total 27 | C 19 | N 4 | 0 4 | 0 | 0 | 0 |
| 2 | Ι | 4 | Total 27 | C 19 | N 4 | 0 4 | 0 | 0 | 0 |
| 2 | J | 5 | Total 32 | C 22 | N 5 | O 5 | 0 | 0 | 0 |
| 2 | К | 4 | Total 27 | C 19 | N 4 | 0 4 | 0 | 0 | 0 |
| 2 | L | 4 | Total 27 | C 19 | N 4 | 0 4 | 0 | 0 | 0 |

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

| Mo | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|----|-------|----------|------------|--------|---------|---------|
| 3 | А | 4 | Total 4 | Zn 4 | 0 | 0 |



Continued from previous page...

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 3 | В | 4 | Total Zn 4 4 | 0 | 0 |
| 3 | С | 4 | Total Zn 4 4 | 0 | 0 |
| 3 | D | 4 | Total Zn 4 4 | 0 | 0 |
| 3 | Ε | 4 | Total Zn 4 4 | 0 | 0 |
| 3 | F | 4 | Total Zn 4 4 | 0 | 0 |

• Molecule 4 is water.

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|---|---------|---------|
| 4 | А | 51 | Total O 51 51 | 0 | 0 |
| 4 | G | 1 | Total O 1 1 | 0 | 0 |
| 4 | В | 58 | Total O 58 58 | 0 | 0 |
| 4 | Н | 4 | Total O 4 4 | 0 | 0 |
| 4 | С | 43 | TotalO4343 | 0 | 0 |
| 4 | D | 48 | Total O 48 48 | 0 | 0 |
| 4 | J | 2 | Total O 2 2 | 0 | 0 |
| 4 | Е | 44 | Total O 44 44 | 0 | 0 |
| 4 | К | 4 | Total O 4 4 | 0 | 0 |
| 4 | F | 57 | Total O 57 57 | 0 | 0 |
| 4 | L | 4 | Total O 4 4 | 0 | 0 |




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Baculoviral IAP repeat-containing protein 8



• Molecule 1: Baculoviral IAP repeat-containing protein 8



| Chain F: | | 73% | 19% · | 5% |
|---|--|--|--|----|
| GLY SER HIS MET SER T254 N255 | N259 R266 1270 1270 1270 1280 8281 7280 7283 7283 7283 7283 7283 7283 | L307 W310 E318 E318 A321 Y324 K328 X328 | 1330 1332 1332 1335 1345 1345 1345 1345 1345 | |
| • Molecule | 2: Diablo homolog, | mitochondrial | | |
| Chain G: | 14% | 43% | 43% | _ |
| A901 V902 1903 1904 GLN LYS | | | | |
| • Molecule | 2: Diablo homolog, | nitochondrial | | |
| Chain H: | 29% | 29% | 43% | _ |
| A901 V902 P903 1904 Ala GlN LYS | | | | |
| • Molecule | 2: Diablo homolog, | nitochondrial | | |
| Chain I: | 14% 4 | 3% | 43% | - |
| A901 V902 P903 1904 ALA GLN LYS | | | | |
| • Molecule | 2: Diablo homolog, | nitochondrial | | |
| Chain J: | 29% | 43% | 29% | _ |
| A901 V902 P903 1904 A905 GLN LYS | | | | |
| • Molecule | 2: Diablo homolog, | mitochondrial | | |
| Chain K: | 43% | 14% | 43% | _ |
| A901 1904 ALA GLN LYS | | | | |
| • Molecule | 2: Diablo homolog, | mitochondrial | | |
| Chain L: | 57% | | 43% | _ |
| A901 1904 ALA GLN LYS | | | | |



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

| Property | Value | Source |
|--|---|-----------|
| Space group | P 41 21 2 | Depositor |
| Cell constants | 86.40Å 86.40Å 226.77Å | Depositor |
| a, b, c, α , β , γ | 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 50.00 - 2.20 | Depositor |
| % Data completeness | 98.7 (50.00-2.20) | Depositor |
| (in resolution range) | 36.1 (50.00-2.20) | Depositor |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | 0.07 | Depositor |
| Refinement program | CNS 1.0 | Depositor |
| R, R_{free} | 0.223 , 0.244 | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| Total number of atoms | 5158 | wwPDB-VP |
| Average B, all atoms $(Å^2)$ | 29.0 | wwPDB-VP |



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond | lengths | Bo | nd angles |
|-----|-------|------|----------|------|---------------|
| | Chain | RMSZ | # Z > 5 | RMSZ | # Z > 5 |
| 1 | А | 0.36 | 0/776 | 0.55 | 0/1051 |
| 1 | В | 0.33 | 0/844 | 0.57 | 0/1143 |
| 1 | С | 0.36 | 0/771 | 0.49 | 0/1044 |
| 1 | D | 0.34 | 0/771 | 0.50 | 0/1044 |
| 1 | Ε | 0.33 | 0/782 | 0.51 | 0/1059 |
| 1 | F | 0.35 | 0/857 | 0.85 | 3/1161~(0.3%) |
| 2 | G | 0.24 | 0/27 | 0.52 | 0/37 |
| 2 | Н | 0.40 | 0/27 | 0.67 | 0/37 |
| 2 | Ι | 0.38 | 0/27 | 0.56 | 0/37 |
| 2 | J | 0.40 | 0/32 | 0.62 | 0/44 |
| 2 | Κ | 0.33 | 0/27 | 0.73 | 0/37 |
| 2 | L | 0.37 | 0/27 | 0.51 | 0/37 |
| All | All | 0.35 | 0/4968 | 0.60 | 3/6731~(0.0%) |

There are no bond length outliers.

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | $Observed(^{o})$ | $Ideal(^{o})$ |
|-----|-------|-----|------|-----------|--------|------------------|---------------|
| 1 | F | 268 | ARG | NE-CZ-NH1 | -15.52 | 112.54 | 120.30 |
| 1 | F | 268 | ARG | NE-CZ-NH2 | 14.82 | 127.71 | 120.30 |
| 1 | F | 268 | ARG | CD-NE-CZ | 6.81 | 133.14 | 123.60 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | А | 751 | 0 | 703 | 13 | 0 |
| 1 | В | 819 | 0 | 780 | 16 | 0 |
| 1 | С | 746 | 0 | 701 | 13 | 0 |
| 1 | D | 746 | 0 | 701 | 12 | 1 |
| 1 | Е | 757 | 0 | 708 | 10 | 0 |
| 1 | F | 832 | 0 | 788 | 16 | 0 |
| 2 | G | 27 | 0 | 31 | 4 | 0 |
| 2 | Н | 27 | 0 | 31 | 2 | 0 |
| 2 | Ι | 27 | 0 | 31 | 3 | 0 |
| 2 | J | 32 | 0 | 36 | 3 | 0 |
| 2 | Κ | 27 | 0 | 31 | 2 | 0 |
| 2 | L | 27 | 0 | 31 | 0 | 0 |
| 3 | А | 4 | 0 | 0 | 0 | 0 |
| 3 | В | 4 | 0 | 0 | 0 | 0 |
| 3 | С | 4 | 0 | 0 | 0 | 0 |
| 3 | D | 4 | 0 | 0 | 0 | 0 |
| 3 | Е | 4 | 0 | 0 | 0 | 0 |
| 3 | F | 4 | 0 | 0 | 0 | 0 |
| 4 | А | 51 | 0 | 0 | 0 | 0 |
| 4 | В | 58 | 0 | 0 | 1 | 0 |
| 4 | С | 43 | 0 | 0 | 0 | 0 |
| 4 | D | 48 | 0 | 0 | 0 | 0 |
| 4 | Е | 44 | 0 | 0 | 0 | 1 |
| 4 | F | 57 | 0 | 0 | 0 | 1 |
| 4 | G | 1 | 0 | 0 | 0 | 0 |
| 4 | Н | 4 | 0 | 0 | 0 | 0 |
| 4 | J | 2 | 0 | 0 | 0 | 0 |
| 4 | K | 4 | 0 | 0 | 0 | 0 |
| 4 | L | 4 | 0 | 0 | 0 | 0 |
| All | All | 5158 | 0 | 4572 | 84 | 2 |

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|-----------------|-----------------------------|----------------------|
| 1:A:311:LYS:HB2 | 1:A:314:GLU:HG3 | 1.54 | 0.88 |
| 1:A:292:ILE:HD11 | 1:A:297:LYS:HD3 | 1.52 | 0.88 |



| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|-----------------------------|----------------------|
| 1:F:280:ASN:HD21 | 1:F:282:GLU:HB2 | 1.42 | 0.83 |
| 1:F:336:HIS:CE1 | 1:F:340:ASN:HD21 | 2.03 | 0.76 |
| 1:F:345:THR:HG21 | 1:F:356:THR:HG21 | 1.65 | 0.76 |

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|----------------------|-----------------------------|----------------------|
| 4:E:839:HOH:O | 4:E:839:HOH:O[7_555] | 2.03 | 0.17 |
| 1:D:333:GLU:OE2 | 4:F:937:HOH:O[7_555] | 2.17 | 0.03 |

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percent | iles |
|-----|-------|---------------|-----------|---------|----------|---------|------|
| 1 | А | 90/108~(83%) | 86~(96%) | 3~(3%) | 1 (1%) | 14 1 | 12 |
| 1 | В | 99/108~(92%) | 94~(95%) | 5 (5%) | 0 | 100 1 | 100 |
| 1 | С | 89/108~(82%) | 85~(96%) | 4 (4%) | 0 | 100 1 | 100 |
| 1 | D | 89/108~(82%) | 86~(97%) | 3~(3%) | 0 | 100 1 | 100 |
| 1 | Е | 91/108~(84%) | 82 (90%) | 8 (9%) | 1 (1%) | 14 1 | 12 |
| 1 | F | 101/108~(94%) | 95~(94%) | 5 (5%) | 1 (1%) | 15 1 | 14 |
| 2 | G | 2/7~(29%) | 2(100%) | 0 | 0 | 100 1 | 100 |
| 2 | Н | 2/7~(29%) | 2(100%) | 0 | 0 | 100 1 | 100 |
| 2 | Ι | 2/7~(29%) | 2(100%) | 0 | 0 | 100 1 | 100 |
| 2 | J | 3/7~(43%) | 3~(100%) | 0 | 0 | 100 1 | 100 |
| 2 | Κ | 2/7~(29%) | 2(100%) | 0 | 0 | 100 1 | 100 |
| 2 | L | 2/7~(29%) | 2 (100%) | 0 | 0 | 100 1 | L00 |
| All | All | 572/690~(83%) | 541 (95%) | 28~(5%) | 3~(0%) | 29 3 | 31 |



1 XB0

All (3) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | А | 255 | ASN |
| 1 | Е | 273 | GLY |
| 1 | F | 255 | ASN |

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentile | \mathbf{s} |
|-----|--------------|---------------|-----------|----------|------------|--------------|
| 1 | А | 76/90~(84%) | 71~(93%) | 5(7%) | 16 19 | |
| 1 | В | 84/90~(93%) | 81~(96%) | 3~(4%) | 35 45 | |
| 1 | \mathbf{C} | 76/90~(84%) | 72~(95%) | 4(5%) | 22 27 | |
| 1 | D | 76/90~(84%) | 72~(95%) | 4(5%) | 22 27 | |
| 1 | Ε | 77/90~(86%) | 73~(95%) | 4(5%) | 23 28 | |
| 1 | F | 85/90~(94%) | 82~(96%) | 3~(4%) | 36 46 | |
| 2 | G | 3/5~(60%) | 3~(100%) | 0 | 100 100 | |
| 2 | Η | 3/5~(60%) | 3~(100%) | 0 | 100 100 | |
| 2 | Ι | 3/5~(60%) | 3~(100%) | 0 | 100 100 | |
| 2 | J | 3/5~(60%) | 3~(100%) | 0 | 100 100 | |
| 2 | Κ | 3/5~(60%) | 3~(100%) | 0 | 100 100 | |
| 2 | L | 3/5~(60%) | 3~(100%) | 0 | 100 100 | |
| All | All | 492/570~(86%) | 469~(95%) | 23~(5%) | 26 33 | |

5 of 23 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | D | 269 | LEU |
| 1 | Е | 259 | ASN |
| 1 | Е | 258 | ARG |
| 1 | Е | 269 | LEU |
| 1 | В | 269 | LEU |



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | Е | 259 | ASN |
| 1 | Е | 336 | HIS |
| 1 | Е | 309 | ASN |
| 1 | F | 259 | ASN |
| 1 | С | 259 | ASN |

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 24 ligands modelled in this entry, 24 are monoatomic - leaving 0 for Mogul analysis. There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

