



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 07:16 AM EDT

PDB ID : 1XD9
Title : Crystal Structure of the Nitrogenase Fe protein Asp39Asn with MgADP bound
Authors : Jang, S.B.; Jeong, M.S.; Seefeldt, L.C.; Peters, J.W.
Deposited on : 2004-09-05
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	
1	B	289	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrogenase iron protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	289	2187	1364	370	432	21	0	0	0
1	B	289	2187	1364	370	432	21	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASN	ASP	engineered mutation	UNP P00459
B	39	ASN	ASP	engineered mutation	UNP P00459

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
2	A	1	1	1	0	0
2	B	1	1	1	0	0

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe S	0	0
			8	4 4		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitrogenase iron protein 1



- Molecule 1: Nitrogenase iron protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.05Å 92.21Å 61.38Å 90.00° 101.23° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 19.71 – 2.70	Depositor EDS
% Data completeness (in resolution range)	85.2 (20.00-2.80) 89.7 (19.71-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.71Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.229 , 0.299 0.246 , 0.309	Depositor DCC
R_{free} test set	738 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	62.6	Xtrriage
Anisotropy	0.407	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4438	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/2211	1.72	21/2977 (0.7%)
1	B	0.70	0/2211	1.75	21/2977 (0.7%)
All	All	0.70	0/4422	1.73	42/5954 (0.7%)

There are no bond length outliers.

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	245	ARG	NE-CZ-NH1	23.02	131.81	120.30
1	B	100	ARG	NE-CZ-NH1	-22.02	109.29	120.30
1	B	245	ARG	NE-CZ-NH2	21.55	131.08	120.30
1	B	100	ARG	NE-CZ-NH2	21.54	131.07	120.30
1	A	191	ARG	NE-CZ-NH2	21.32	130.96	120.30
1	A	191	ARG	NE-CZ-NH1	-21.00	109.80	120.30
1	B	191	ARG	NE-CZ-NH2	-20.86	109.87	120.30
1	A	100	ARG	NE-CZ-NH2	-20.29	110.15	120.30
1	B	191	ARG	NE-CZ-NH1	20.19	130.40	120.30
1	B	245	ARG	NE-CZ-NH1	-19.35	110.62	120.30
1	A	100	ARG	NE-CZ-NH1	19.01	129.80	120.30
1	A	245	ARG	NE-CZ-NH2	-18.50	111.05	120.30
1	A	245	ARG	CD-NE-CZ	18.13	148.98	123.60
1	B	245	ARG	CD-NE-CZ	17.24	147.73	123.60
1	B	191	ARG	CD-NE-CZ	16.92	147.29	123.60
1	B	100	ARG	CD-NE-CZ	16.39	146.55	123.60
1	A	191	ARG	CD-NE-CZ	15.31	145.04	123.60
1	A	100	ARG	CD-NE-CZ	15.02	144.62	123.60
1	A	187	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	A	187	ARG	NE-CZ-NH2	-6.72	116.94	120.30
1	A	213	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	A	118	ASP	CA-C-N	-6.48	102.94	117.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	95	VAL	CA-CB-CG2	-6.34	101.39	110.90
1	B	213	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	B	187	ARG	NE-CZ-NH1	6.31	123.45	120.30
1	B	118	ASP	O-C-N	6.29	132.76	122.70
1	B	118	ASP	CA-C-N	-6.25	103.45	117.20
1	B	187	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	B	46	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	A	118	ASP	O-C-N	5.85	132.07	122.70
1	A	95	VAL	CA-CB-CG2	-5.77	102.25	110.90
1	B	22	LEU	CA-CB-CG	5.54	128.05	115.30
1	A	124	TYR	CB-CG-CD2	-5.39	117.77	121.00
1	A	22	LEU	CA-CB-CG	5.37	127.65	115.30
1	B	285	THR	CA-CB-CG2	5.29	119.81	112.40
1	A	190	ASP	N-CA-C	5.28	125.26	111.00
1	A	118	ASP	CB-CG-OD1	5.28	123.05	118.30
1	B	190	ASP	N-CA-C	5.23	125.13	111.00
1	B	100	ARG	CB-CG-CD	-5.23	98.01	111.60
1	A	285	THR	CA-CB-CG2	5.18	119.65	112.40
1	A	46	ARG	NE-CZ-NH1	5.05	122.82	120.30
1	B	4	GLN	CA-C-N	-5.01	106.18	117.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2187	0	2199	48	0
1	B	2187	0	2199	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	0	0	0
4	A	27	0	12	3	0
4	B	27	0	12	1	0
All	All	4438	0	4422	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:CYS:HG	1:A:148:TYR:HE2	1.30	0.77
1:B:76:LEU:HD21	1:B:84:LYS:HB3	1.70	0.72
1:B:165:SER:HA	1:B:168:ILE:HD12	1.78	0.65
1:A:165:SER:HA	1:A:168:ILE:HD12	1.78	0.65
1:A:57:ILE:HD11	1:A:75:VAL:HG13	1.79	0.64
1:A:76:LEU:HD21	1:A:84:LYS:HB3	1.79	0.63
1:B:20:GLN:HE22	1:B:47:LEU:H	1.48	0.61
1:B:57:ILE:HD11	1:B:75:VAL:HG13	1.82	0.61
1:B:284:LYS:HD3	1:B:288:GLU:HB3	1.83	0.60
1:A:284:LYS:HD3	1:A:288:GLU:HB3	1.83	0.60
1:B:255:ILE:HD12	1:B:255:ILE:H	1.66	0.60
1:B:268:LEU:HB3	1:B:274:MET:HB2	1.83	0.60
1:A:92:GLU:HB3	1:A:95:VAL:HG12	1.85	0.59
1:B:213:ARG:HD2	4:B:292:ADP:C2	2.36	0.59
1:A:268:LEU:HB3	1:A:274:MET:HB2	1.85	0.59
1:A:213:ARG:HD2	4:A:291:ADP:C2	2.38	0.58
1:A:255:ILE:H	1:A:255:ILE:HD12	1.68	0.58
1:A:20:GLN:HE22	1:A:47:LEU:H	1.50	0.58
1:B:187:ARG:HA	1:B:213:ARG:NH1	2.20	0.56
1:B:92:GLU:HB3	1:B:95:VAL:HG12	1.88	0.56
1:B:238:ASP:HA	1:B:241:ARG:HB2	1.88	0.55
1:A:187:ARG:HA	1:A:213:ARG:NH1	2.22	0.55
1:A:238:ASP:HA	1:A:241:ARG:HB2	1.88	0.55
1:A:33:VAL:HG22	1:A:121:PHE:HB2	1.89	0.54
1:B:13:ILE:HD12	1:B:151:CYS:HA	1.90	0.54
1:B:279:GLU:O	1:B:282:VAL:HG22	2.08	0.53
1:B:27:ALA:HA	1:B:31:LYS:H	1.74	0.52
1:B:56:THR:HA	1:B:87:GLU:O	2.10	0.52
1:A:279:GLU:O	1:A:282:VAL:HG22	2.09	0.52
1:A:13:ILE:HD12	1:A:151:CYS:HA	1.92	0.51
1:B:105:ALA:O	1:B:109:LEU:HD23	2.11	0.50
1:B:33:VAL:HG22	1:B:121:PHE:HB2	1.94	0.50
1:A:49:LEU:HD13	1:A:85:CYS:SG	2.53	0.49
1:B:3:ARG:HH12	1:B:248:VAL:HA	1.78	0.49
1:A:56:THR:HA	1:A:87:GLU:O	2.13	0.49
1:A:27:ALA:HA	1:A:31:LYS:H	1.78	0.48
1:A:149:ILE:HG21	1:A:161:ALA:HA	1.93	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281:ILE:HG22	1:A:289:VAL:HG21	1.95	0.48
1:B:211:VAL:HA	1:B:212:PRO:HD3	1.78	0.48
1:B:260:THR:HB	1:B:263:GLU:H	1.79	0.48
1:B:5:CYS:SG	1:B:146:GLU:HB2	2.53	0.48
1:B:281:ILE:HG22	1:B:289:VAL:HG21	1.95	0.47
1:B:148:TYR:CE1	1:B:180:GLY:HA3	2.50	0.47
1:A:284:LYS:O	1:B:223:ARG:HD3	2.14	0.46
1:A:260:THR:HB	1:A:263:GLU:H	1.80	0.46
1:A:105:ALA:O	1:A:109:LEU:HD23	2.16	0.46
1:B:155:MET:SD	1:B:264:LEU:HD21	2.55	0.46
1:A:95:VAL:HG23	1:B:171:TYR:OH	2.16	0.46
1:A:168:ILE:HD13	1:A:179:LEU:HB2	1.99	0.45
1:B:102:VAL:O	1:B:106:ILE:HG12	2.16	0.45
1:A:3:ARG:HH12	1:A:248:VAL:HA	1.82	0.45
1:B:49:LEU:HD13	1:B:85:CYS:SG	2.57	0.45
1:B:235:LYS:O	1:B:239:GLU:HG2	2.16	0.45
1:A:148:TYR:CE1	1:A:180:GLY:HA3	2.52	0.45
1:B:149:ILE:HG21	1:B:161:ALA:HA	1.97	0.45
1:A:182:LEU:O	1:A:208:ILE:HG22	2.17	0.44
1:A:49:LEU:CD1	1:A:85:CYS:SG	3.05	0.44
1:A:5:CYS:SG	1:A:146:GLU:HB2	2.57	0.44
1:A:155:MET:SD	1:A:264:LEU:HD21	2.58	0.44
1:B:187:ARG:HA	1:B:213:ARG:HH12	1.81	0.44
1:B:193:ASP:O	1:B:197:ILE:HG13	2.17	0.44
1:B:4:GLN:HB2	1:B:145:GLN:HE21	1.83	0.44
1:A:10:LYS:O	1:A:13:ILE:HG12	2.18	0.43
1:A:187:ARG:HA	1:A:213:ARG:HH12	1.82	0.43
1:B:168:ILE:HD13	1:B:179:LEU:HB2	2.00	0.43
1:B:162:ASN:O	1:B:165:SER:HB2	2.18	0.43
1:B:5:CYS:SG	1:B:148:TYR:HE2	2.41	0.43
1:A:235:LYS:HE3	1:A:235:LYS:HB3	1.87	0.43
1:A:162:ASN:HD21	1:A:259:ILE:H	1.67	0.43
1:B:66:THR:HG23	1:B:68:GLU:H	1.83	0.42
1:B:10:LYS:O	1:B:13:ILE:HG12	2.19	0.42
1:A:4:GLN:HB2	1:A:145:GLN:HE21	1.84	0.42
1:A:66:THR:HG23	1:A:68:GLU:H	1.84	0.42
1:A:193:ASP:O	1:A:197:ILE:HG13	2.20	0.42
1:B:103:ILE:HD12	1:B:103:ILE:HA	1.88	0.41
1:A:235:LYS:O	1:A:239:GLU:HG2	2.20	0.41
1:A:223:ARG:NE	1:B:281:ILE:HB	2.34	0.41
1:B:61:ALA:HA	1:B:70:LEU:HD11	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ILE:HG22	1:A:207:MET:HG3	2.02	0.41
1:A:236:GLN:HA	1:A:236:GLN:NE2	2.36	0.41
1:A:103:ILE:HD12	1:A:103:ILE:HA	1.91	0.41
1:B:139:ILE:HG22	1:B:171:TYR:CD2	2.56	0.41
1:A:14:GLY:HA2	4:A:291:ADP:O2A	2.21	0.41
1:A:159:TYR:O	1:A:162:ASN:HB3	2.21	0.40
1:A:14:GLY:HA2	4:A:291:ADP:PA	2.61	0.40
1:B:235:LYS:HE3	1:B:235:LYS:HB3	1.87	0.40
1:A:54:GLN:HE22	1:A:86:VAL:HA	1.87	0.40
1:A:61:ALA:HA	1:A:70:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/289 (99%)	262 (91%)	23 (8%)	2 (1%)	22	53
1	B	287/289 (99%)	262 (91%)	23 (8%)	2 (1%)	22	53
All	All	574/578 (99%)	524 (91%)	46 (8%)	4 (1%)	22	53

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	190	ASP
1	B	190	ASP
1	A	285	THR
1	B	285	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/233 (100%)	214 (92%)	19 (8%)	11	33
1	B	233/233 (100%)	211 (91%)	22 (9%)	8	26
All	All	466/466 (100%)	425 (91%)	41 (9%)	10	29

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	66	THR
1	A	73	GLU
1	A	86	VAL
1	A	92	GLU
1	A	95	VAL
1	A	120	ASP
1	A	156	MET
1	A	178	ARG
1	A	187	ARG
1	A	189	THR
1	A	190	ASP
1	A	201	ASN
1	A	218	GLN
1	A	226	THR
1	A	235	LYS
1	A	238	ASP
1	A	241	ARG
1	A	255	ILE
1	B	22	LEU
1	B	66	THR
1	B	73	GLU
1	B	86	VAL
1	B	92	GLU
1	B	95	VAL
1	B	120	ASP
1	B	156	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	178	ARG
1	B	187	ARG
1	B	189	THR
1	B	190	ASP
1	B	191	ARG
1	B	201	ASN
1	B	218	GLN
1	B	226	THR
1	B	235	LYS
1	B	238	ASP
1	B	241	ARG
1	B	245	ARG
1	B	255	ILE
1	B	260	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	54	GLN
1	A	145	GLN
1	A	162	ASN
1	A	185	ASN
1	B	20	GLN
1	B	50	HIS
1	B	54	GLN
1	B	55	ASN
1	B	107	ASN
1	B	145	GLN
1	B	162	ASN
1	B	173	ASN
1	B	185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SF4	A	290	1	0,12,12	-	-	-		
4	ADP	B	292	-	24,29,29	2.20	4 (16%)	29,45,45	1.88	9 (31%)
4	ADP	A	291	-	24,29,29	1.57	3 (12%)	29,45,45	1.68	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	B	292	-	-	1/12/32/32	0/3/3/3
3	SF4	A	290	1	-	-	0/6/5/5
4	ADP	A	291	-	-	1/12/32/32	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	292	ADP	PB-O1B	8.31	1.77	1.50
4	A	291	ADP	PB-O3B	5.21	1.74	1.54
4	B	292	ADP	O4'-C1'	4.05	1.46	1.41
4	A	291	ADP	O4'-C1'	3.41	1.45	1.41
4	B	292	ADP	C3'-C4'	2.92	1.60	1.53
4	A	291	ADP	C8-N7	-2.16	1.30	1.34
4	B	292	ADP	C8-N7	-2.09	1.31	1.34

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	292	ADP	O2B-PB-O3A	3.93	117.82	104.64
4	A	291	ADP	C3'-C2'-C1'	-3.56	95.62	100.98
4	B	292	ADP	O3B-PB-O3A	3.39	116.00	104.64
4	A	291	ADP	O4'-C1'-C2'	-3.38	101.98	106.93
4	B	292	ADP	C2'-C3'-C4'	-3.25	96.32	102.64
4	A	291	ADP	O2B-PB-O3A	3.24	115.50	104.64
4	B	292	ADP	C3'-C2'-C1'	-3.12	96.29	100.98
4	A	291	ADP	C2'-C3'-C4'	-2.92	96.96	102.64
4	A	291	ADP	O3B-PB-O1B	-2.85	99.51	110.68
4	B	292	ADP	O4'-C1'-C2'	-2.79	102.85	106.93
4	B	292	ADP	PA-O3A-PB	-2.73	123.47	132.83
4	B	292	ADP	O3A-PB-O1B	-2.60	96.79	111.19
4	B	292	ADP	O2B-PB-O1B	-2.43	101.17	110.68
4	A	291	ADP	PA-O3A-PB	-2.38	124.66	132.83
4	B	292	ADP	O5'-C5'-C4'	2.08	116.16	108.99

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	292	ADP	PA-O3A-PB-O1B
4	A	291	ADP	C5'-O5'-PA-O1A

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	292	ADP	1	0
4	A	291	ADP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/289 (100%)	-0.03	2 (0%) 87 84	37, 82, 99, 100	0
1	B	289/289 (100%)	0.03	7 (2%) 59 49	41, 85, 100, 100	0
All	All	578/578 (100%)	0.00	9 (1%) 72 66	37, 83, 100, 100	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	226	THR	3.2
1	B	213	ARG	2.8
1	B	1	ALA	2.7
1	B	233	LYS	2.6
1	B	82	GLY	2.5
1	B	116	GLU	2.4
1	B	65	GLY	2.4
1	A	1	ALA	2.3
1	A	85	CYS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	B	294	1/1	0.80	0.11	57,57,57,57	0
4	ADP	B	292	27/27	0.94	0.15	77,85,89,91	0
4	ADP	A	291	27/27	0.95	0.12	66,78,88,90	0
2	MG	A	293	1/1	0.95	0.09	49,49,49,49	0
3	SF4	A	290	8/8	0.99	0.08	34,37,39,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.