



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 4, 2024 – 07:28 PM EST

PDB ID : 1XDI  
Title : Crystal structure of LpdA (Rv3303c) from Mycobacterium tuberculosis  
Authors : Argyrou, A.; Vetting, M.W.; Blanchard, J.S.  
Deposited on : 2004-09-06  
Resolution : 2.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4060 (2.84-2.80)
Ramachandran outliers	138981	3978 (2.84-2.80)
Sidechain outliers	138945	3980 (2.84-2.80)
RSRZ outliers	127900	3552 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	499	
1	B	499	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rv3303c-lpdA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	459	3356	2101	607	635	13	0	0	0
1	B	459	3356	2101	607	635	13	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	494	HIS	-	expression tag	UNP O53355
A	495	HIS	-	expression tag	UNP O53355
A	496	HIS	-	expression tag	UNP O53355
A	497	HIS	-	expression tag	UNP O53355
A	498	HIS	-	expression tag	UNP O53355
A	499	HIS	-	expression tag	UNP O53355
B	494	HIS	-	expression tag	UNP O53355
B	495	HIS	-	expression tag	UNP O53355
B	496	HIS	-	expression tag	UNP O53355
B	497	HIS	-	expression tag	UNP O53355
B	498	HIS	-	expression tag	UNP O53355
B	499	HIS	-	expression tag	UNP O53355

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	53	27	9	15	2	0	0
2	B	1	53	27	9	15	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	57	57	57	0	0
3	B	50	50	50	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Rv3303c-lpdA



#### • Molecule 1: Rv3303c-lpdA



H338
A339
L340
G341
E342
P346
I347
R348
L349
R350
T355
V356
F357
T358
V365
G366
V367
P368
Q369
S370
V371
I372
D373
S376
R380
T381
I382
M383
L384
R387
A390
R391
A392
S395
E396
M397
R398
F401
V402
K403
I404
F405
S409
T410
G411
V412
V413
I414
G415
S424
E425
L426
I427
L428
P429
V432
A433
N436
R437
I438
T439
V440
N441
E442
L443
A444
Q445
T446
L447
P451
S452
L453
S454
G455
S456
I457
T458
E459
A460
A461
R462
R463
L464
M465
A466
HIS
ASP
ASP
LEU
ASP
ASP
CYS
THR
ALA
ALA
ALA
GLN
ALA
ALA
ALA
ALA
GLU
GLN
LEU
ALA
LEU
VAL
PRO
HIS
HIS
LEU
PHO
THR
SER
ASN
HIS
HIS
HIS
HIS
HIS
HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	244.20Å 244.20Å 104.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.81 35.25 – 2.81	Depositor EDS
% Data completeness (in resolution range)	98.9 (50.00-2.81) 99.7 (35.25-2.81)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.31 (at 2.81Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.194 , 0.276 0.191 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.4	Xtrriage
Anisotropy	0.100	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6925	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.86	3/3404 (0.1%)	1.06	4/4637 (0.1%)
1	B	0.87	0/3404	1.06	8/4637 (0.2%)
All	All	0.86	3/6808 (0.0%)	1.06	12/9274 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	306	ARG	CB-CG	-5.17	1.38	1.52
1	A	434	VAL	CB-CG1	-5.05	1.42	1.52
1	A	201	TYR	CD2-CE2	5.04	1.47	1.39

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	306	ARG	CG-CD-NE	-6.36	98.45	111.80
1	B	183	ASP	CB-CG-OD1	-5.74	113.13	118.30
1	B	262	ASP	CB-CA-C	-5.72	98.96	110.40
1	B	414	ILE	CG1-CB-CG2	-5.59	99.11	111.40
1	B	128	ARG	NE-CZ-NH1	-5.44	117.58	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	220	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3356	0	3454	187	0
1	B	3356	0	3454	217	0
2	A	53	0	31	5	0
2	B	53	0	31	2	0
3	A	57	0	0	4	0
3	B	50	0	0	4	0
All	All	6925	0	6970	385	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 385 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:275:ILE:HG22	1:B:276:GLY:H	1.03	1.14
1:A:275:ILE:HG22	1:A:276:GLY:H	1.00	1.13
1:B:308:LEU:H	1:B:308:LEU:HD12	1.16	1.11
1:A:70:LEU:HD13	1:B:60:LEU:HD11	1.28	1.06
1:A:308:LEU:HD12	1:A:308:LEU:H	1.14	1.06

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/499 (91%)	418 (92%)	30 (7%)	7 (2%)	10	31
1	B	455/499 (91%)	413 (91%)	34 (8%)	8 (2%)	8	26
All	All	910/998 (91%)	831 (91%)	64 (7%)	15 (2%)	9	29

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	63	GLU
1	A	64	LEU
1	A	253	GLY
1	A	261	THR
1	A	275	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/386 (91%)	306 (87%)	46 (13%)	4	12
1	B	352/386 (91%)	301 (86%)	51 (14%)	3	9
All	All	704/772 (91%)	607 (86%)	97 (14%)	3	10

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	85	GLN
1	B	239	ARG
1	B	92	THR
1	B	161	SER
1	B	262	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	338	HIS
1	B	441	ASN
1	B	69	HIS
1	B	104	GLN
1	B	270	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	B	999	-	53,58,58	2.45	14 (26%)	68,89,89	1.47	13 (19%)
2	FAD	A	999	-	53,58,58	2.20	14 (26%)	68,89,89	1.43	11 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	999	-	-	6/30/50/50	0/6/6/6
2	FAD	A	999	-	-	7/30/50/50	0/6/6/6

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	999	FAD	C4A-N3A	8.50	1.47	1.35
2	B	999	FAD	C4X-N5	7.23	1.44	1.30
2	A	999	FAD	C4X-N5	6.21	1.42	1.30
2	A	999	FAD	C4A-N3A	5.91	1.43	1.35
2	B	999	FAD	C9A-N10	5.11	1.50	1.41

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	999	FAD	O4'-C4'-C5'	-3.67	101.67	109.92
2	A	999	FAD	C3B-C2B-C1B	3.27	105.90	100.98
2	B	999	FAD	N3A-C2A-N1A	-3.13	123.78	128.68
2	A	999	FAD	O4'-C4'-C5'	-3.08	102.98	109.92
2	A	999	FAD	C5'-C4'-C3'	3.07	118.14	112.20

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	999	FAD	C5B-O5B-PA-O1A
2	A	999	FAD	C5B-O5B-PA-O3P
2	B	999	FAD	C5B-O5B-PA-O3P
2	A	999	FAD	P-O3P-PA-O1A
2	A	999	FAD	PA-O3P-P-O5'


There are no ring outliers.


2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	999	FAD	2	0
2	A	999	FAD	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	459/499 (91%)	-0.74	1 (0%) 95   94	1, 12, 34, 61	0
1	B	459/499 (91%)	-0.68	0 100   100	1, 12, 37, 51	0
All	All	918/998 (91%)	-0.71	1 (0%) 95   95	1, 12, 36, 61	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	80	LYS	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FAD	A	999	53/53	0.97	0.13	1,6,10,12	0
2	FAD	B	999	53/53	0.98	0.12	1,8,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.