



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 06:10 am BST

PDB ID : 1XSQ  
Title : Crystal structure of ureidoglycolate hydrolase from E.coli. Northeast Structural Genomics Consortium target ET81.  
Authors : Kuzin, A.P.; Vorobiev, S.M.; Abashidze, M.; Acton, T.B.; Ma, L.-C.; Xiao, R.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2004-10-19  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	 8% 77% 13% • 8%
1	B	168	 11% 77% 14% • 8%

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 2740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ureidoglycolate hydrolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	154	1243	793	213	232	3	2	0	0	0
1	B	155	1251	799	214	233	3	2	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P77731
A	84	MSE	MET	MODIFIED RESIDUE	UNP P77731
A	161	LEU	-	CLONING ARTIFACT	UNP P77731
A	162	GLU	-	CLONING ARTIFACT	UNP P77731
A	163	HIS	-	CLONING ARTIFACT	UNP P77731
A	164	HIS	-	CLONING ARTIFACT	UNP P77731
A	165	HIS	-	CLONING ARTIFACT	UNP P77731
A	166	HIS	-	CLONING ARTIFACT	UNP P77731
A	167	HIS	-	CLONING ARTIFACT	UNP P77731
A	168	HIS	-	CLONING ARTIFACT	UNP P77731
B	1	MSE	MET	MODIFIED RESIDUE	UNP P77731
B	84	MSE	MET	MODIFIED RESIDUE	UNP P77731
B	161	LEU	-	CLONING ARTIFACT	UNP P77731
B	162	GLU	-	CLONING ARTIFACT	UNP P77731
B	163	HIS	-	CLONING ARTIFACT	UNP P77731
B	164	HIS	-	CLONING ARTIFACT	UNP P77731
B	165	HIS	-	CLONING ARTIFACT	UNP P77731
B	166	HIS	-	CLONING ARTIFACT	UNP P77731
B	167	HIS	-	CLONING ARTIFACT	UNP P77731
B	168	HIS	-	CLONING ARTIFACT	UNP P77731

- Molecule 2 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	A	122	Total 122	O 122	0	0
2	B	124	Total 124	O 124	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.19Å 45.27Å 51.80Å 100.70° 99.84° 117.19°	Depositor
Resolution (Å)	24.34 – 1.60 24.34 – 1.60	Depositor EDS
% Data completeness (in resolution range)	85.0 (24.34-1.60) 94.7 (24.34-1.60)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.11 (at 1.60Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.227 , 0.257 0.236 , 0.267	Depositor DCC
$R_{free}$ test set	3967 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.0	Xtrriage
Anisotropy	0.588	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2740	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/1270	0.62	0/1725
1	B	0.33	0/1278	0.63	0/1736
All	All	0.34	0/2548	0.63	0/3461

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1243	0	1207	26	0
1	B	1251	0	1218	28	0
2	A	122	0	0	2	0
2	B	124	0	0	3	0
All	All	2740	0	2425	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:LEU:HD23	1:B:137:LEU:HD23	1.41	1.00

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:59:ALA:HB2	1:B:127:LEU:HD21	1.60	0.81
1:A:137:LEU:CD2	1:B:137:LEU:HD23	2.17	0.72
1:A:30:ILE:HB	1:A:35:VAL:HB	1.73	0.71
1:A:137:LEU:HD23	1:B:137:LEU:CD2	2.20	0.68
1:B:141:ARG:NH1	1:B:146:ASN:H	1.92	0.66
1:B:141:ARG:HH12	1:B:146:ASN:H	1.45	0.62
1:B:160:ALA:O	1:B:161:LEU:HB2	2.00	0.62
1:A:66:LEU:HD13	1:A:157:LEU:HD22	1.82	0.61
1:A:127:LEU:HD13	1:B:35:VAL:HG21	1.81	0.61
1:A:54:ILE:HG12	1:A:139:ILE:HD12	1.83	0.61
1:B:1:MSE:HE2	1:B:155:GLN:HE22	1.65	0.60
1:A:54:ILE:HG12	1:A:139:ILE:CD1	2.33	0.59
1:B:104:THR:HG22	2:B:272:HOH:O	2.02	0.58
1:B:1:MSE:HE2	1:B:155:GLN:NE2	2.21	0.55
1:A:82:ILE:HD13	1:B:137:LEU:HD21	1.87	0.55
1:B:47:LEU:HD12	1:B:48:GLU:H	1.71	0.54
1:A:3:LEU:HB2	1:A:155:GLN:HE21	1.74	0.53
1:A:153:PRO:O	1:A:155:GLN:HG2	2.10	0.52
1:A:157:LEU:HD12	1:A:157:LEU:H	1.74	0.52
1:A:68:ILE:HG23	1:A:157:LEU:HD11	1.92	0.51
1:A:17:GLY:HA2	1:A:119:HIS:CD2	2.46	0.50
1:B:11:GLU:HG3	2:B:244:HOH:O	2.10	0.50
1:A:137:LEU:HD21	1:B:82:ILE:HD13	1.95	0.48
1:A:127:LEU:CD1	1:B:35:VAL:HG21	2.42	0.48
1:A:86:GLY:HA3	1:B:85:LYS:HE2	1.96	0.47
1:B:48:GLU:HB3	1:B:50:ASP:OD2	2.15	0.46
1:A:157:LEU:N	1:A:157:LEU:HD12	2.31	0.46
1:B:141:ARG:HH12	1:B:146:ASN:N	2.14	0.45
2:A:192:HOH:O	1:B:37:ARG:HD2	2.16	0.45
1:A:40:ASP:OD1	1:B:53:LEU:HD13	2.16	0.45
1:A:54:ILE:HB	1:B:42:ALA:HB3	1.99	0.45
1:A:52:THR:HB	1:B:44:VAL:HB	1.97	0.45
1:B:47:LEU:HD12	1:B:48:GLU:N	2.32	0.44
1:A:131:GLN:CB	1:A:132:ARG:HD2	2.48	0.44
1:A:17:GLY:HA2	1:A:119:HIS:CG	2.53	0.43
1:B:1:MSE:HE3	2:B:246:HOH:O	2.19	0.42
1:A:138:THR:O	1:A:139:ILE:HD13	2.19	0.42
1:A:85:LYS:HB2	1:A:87:GLU:OE2	2.20	0.42
1:B:73:ARG:O	1:B:147:CYS:HB2	2.19	0.41
1:B:17:GLY:HA2	1:B:119:HIS:CD2	2.54	0.41
1:B:73:ARG:O	1:B:75:PRO:HD3	2.19	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:ASP:HA	2:A:268:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	148/168 (88%)	142 (96%)	6 (4%)	0	100	100
1	B	149/168 (89%)	141 (95%)	8 (5%)	0	100	100
All	All	297/336 (88%)	283 (95%)	14 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/147 (93%)	135 (98%)	2 (2%)	65	44
1	B	138/147 (94%)	136 (99%)	2 (1%)	67	47
All	All	275/294 (94%)	271 (98%)	4 (2%)	65	44

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	127	LEU
1	A	132	ARG
1	B	47	LEU
1	B	127	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	23	GLN
1	A	60	GLN
1	A	79	GLN
1	A	117	ASN
1	A	146	ASN
1	A	155	GLN
1	B	4	GLN
1	B	24	GLN
1	B	49	GLN
1	B	79	GLN
1	B	117	ASN
1	B	125	HIS
1	B	155	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	152/168 (90%)	0.58	13 (8%) 10   9	10, 18, 35, 50	0
1	B	153/168 (91%)	0.72	19 (12%) 4   3	10, 19, 37, 48	0
All	All	305/336 (90%)	0.65	32 (10%) 6   5	10, 19, 37, 50	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	142	GLY	7.6
1	B	47	LEU	6.7
1	A	160	ALA	5.8
1	B	161	LEU	5.0
1	A	145	ASP	4.6
1	A	146	ASN	4.2
1	B	63	ASN	3.8
1	B	146	ASN	3.6
1	B	99	LYS	3.2
1	B	50	ASP	3.2
1	B	141	ARG	3.2
1	B	145	ASP	3.1
1	A	50	ASP	2.9
1	A	64	LEU	2.9
1	A	147	CYS	2.9
1	B	49	GLN	2.9
1	B	128	PHE	2.8
1	B	127	LEU	2.7
1	B	149	VAL	2.7
1	B	48	GLU	2.6
1	A	127	LEU	2.5
1	B	98	ASP	2.4
1	A	132	ARG	2.4
1	A	154	GLU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	2	LYS	2.4
1	B	23	GLN	2.3
1	A	149	VAL	2.3
1	B	64	LEU	2.2
1	A	99	LYS	2.1
1	B	120	ARG	2.1
1	B	97	ASP	2.1
1	A	139	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.