

Full wwPDB X-ray Structure Validation Report (i)

Oct 16, 2023 – 10:29 AM EDT

PDB ID : 1YI6

Title : C-term tail segment of human tyrosine kinase (258-533)

Authors: Fleury, D.; Sarubbi, E.; Courjaud, A.; Guitton, J.D.; Ducruix, A.

Deposited on : 2005-01-11

Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	A	276	89%	8% •
1	В	276	83%	11% 5% •



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
1	A	276		C 1422		_			0	0	0
1	В	276	Total 2230	C 1422		_	P 2	S 16	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	335	PTR	TYR	modified residue	UNP P12931
A	416	PTR	TYR	modified residue	UNP P12931
В	335	PTR	TYR	modified residue	UNP P12931
В	416	PTR	TYR	modified residue	UNP P12931

• Molecule 2 is water.

ľ	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	2	A	304	Total O 304 304	0	0
	2	В	231	Total O 231 231	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Proto-oncogene tyrosine-protein kinase Src





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	66.98Å 63.42Å 76.69Å	Depositor	
a, b, c, α , β , γ	90.00° 109.54° 90.00°	Depositor	
Resolution (Å)	19.00 - 2.00	Depositor	
% Data completeness	98.0 (19.00-2.00)	Depositor	
(in resolution range)	30.0 (13.00 2.00)	Depositor	
R_{merge}	0.08	Depositor	
R_{sym}	0.07	Depositor	
Refinement program	CNS 1.0	Depositor	
R, R_{free}	0.216 , 0.252	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	4995	wwPDB-VP	
Average B, all atoms (Å ²)	17.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.33	0/2248	0.60	0/3041
1	В	0.34	0/2248	0.72	7/3041 (0.2%)
All	All	0.33	0/4496	0.66	7/6082 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$Observed(^o)$	$Ideal(^{o})$
1	В	277	CYS	N-CA-C	7.27	130.63	111.00
1	В	300	GLY	N-CA-C	6.57	129.52	113.10
1	В	414	ASN	N-CA-C	6.38	128.22	111.00
1	В	301	THR	N-CA-C	5.83	126.75	111.00
1	В	278	PHE	N-CA-C	-5.82	95.28	111.00
1	В	299	PRO	N-CA-C	5.78	127.12	112.10
1	В	415	GLU	N-CA-C	5.36	125.47	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2230	0	2198	13	0
1	В	2230	0	2198	32	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	304	0	0	0	0
2	В	231	0	0	1	0
All	All	4995	0	4396	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

1:B:308:LEU:HD21 1 1:B:418:ALA:HB3 1 1:B:275:GLN:O 1 1:B:525:PRO:HB2 1	1:A:423:LYS:H 1:A:335:PTR:HB3 1:B:334:ILE:HD11 :B:420:GLN:HE21 1:B:275:GLN:HG3	1.53 1.71 1.77 1.62	0.74 0.71 0.66
1:A:329:VAL:HB 1 1:B:308:LEU:HD21 1 1:B:418:ALA:HB3 1 1:B:275:GLN:O 1 1:B:525:PRO:HB2 1	1:A:335:PTR:HB3 1:B:334:ILE:HD11 :B:420:GLN:HE21	1.71 1.77	0.71
1:B:308:LEU:HD21 1 1:B:418:ALA:HB3 1 1:B:275:GLN:O 1 1:B:525:PRO:HB2 1	1:B:334:ILE:HD11 :B:420:GLN:HE21	1.77	
1:B:418:ALA:HB3 1 1:B:275:GLN:O 1 1:B:525:PRO:HB2 1	:B:420:GLN:HE21		LLAN
1:B:275:GLN:O 1 1:B:525:PRO:HB2 1		1.62	
1:B:525:PRO:HB2 1	1:B:275:GLN:HG3		0.64
	D voc GI II II Dod	1.99	0.62
	:B:526:GLN:HE21	1.66	0.61
	1:B:298:LYS:HZ2	1.65	0.61
	1:A:322:LEU:HG	1.82	0.60
1:B:418:ALA:C	1:B:420:GLN:H	2.05	0.60
	1:B:526:GLN:NE2	2.22	0.54
	:A:301:THR:HG21	2.07	0.54
1:A:353:GLU:O	1:A:356:LYS:HG2	2.09	0.52
1:A:438:ARG:HD3 1	1:B:397:ASN:OD1	2.10	0.52
1:B:424:PHE:CD2	1:B:424:PHE:O	2.63	0.51
1:A:413:ASP:HB3 1	1:A:416:PTR:HB2	1.92	0.51
1:B:381:ASN:HB3	2:B:642:HOH:O	2.12	0.50
1:B:268:ARG:HH11 1	1:B:268:ARG:HB3	1.78	0.49
1:B:308:LEU:CD2 1	1:B:334:ILE:HD11	2.43	0.48
1:B:416:PTR:CG	1:B:417:THR:N	2.76	0.48
1:B:426:ILE:HD11 1	:B:472:LEU:HD13	1.95	0.48
1:A:500:ARG:O 1	1:A:506:ARG:HD2	2.14	0.48
1:A:423:LYS:H	1:A:423:LYS:CE	2.23	0.47
1:B:323:VAL:HG21 1	:B:393:LEU:HD12	1.95	0.47
1:B:418:ALA:C	1:B:420:GLN:N	2.67	0.47
1:B:329:VAL:O 1	1:B:334:ILE:HG13	2.13	0.47
1:A:334:ILE:O 1	1:A:335:PTR:HD1	2.15	0.47
1:B:477:ARG:O 1	1:B:477:ARG:HG3	2.14	0.47
1:B:297:LEU:HB2	1:B:307:PHE:CE1	2.50	0.46
1:B:280:GLU:HG3	1:B:298:LYS:NZ	2.31	0.46
1:B:275:GLN:OE1	1:B:277:CYS:SG	2.75	0.45
1:B:311:ALA:O	1:B:315:LYS:HG3	2.17	0.45



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
1100111 1	1100111 2	$\operatorname{distance}\left(\mathrm{\AA}\right)$	overlap (Å)
1:B:419:ARG:HH11	1:B:419:ARG:HG3	1.82	0.44
1:B:278:PHE:O	1:B:297:LEU:HD12	2.18	0.43
1:B:423:LYS:HE3	1:B:423:LYS:H	1.83	0.43
1:B:424:PHE:O	1:B:426:ILE:N	2.52	0.42
1:B:375:ALA:O	1:B:379:ARG:HG3	2.19	0.42
1:A:346:LEU:HD22	1:A:350:LEU:HD22	2.01	0.41
1:B:298:LYS:HA	1:B:299:PRO:HD3	1.80	0.41
1:A:272:LYS:HG2	1:A:282:TRP:CE2	2.55	0.41
1:B:275:GLN:NE2	1:B:280:GLU:OE1	2.54	0.41
1:B:362:GLN:O	1:B:366:MET:HG3	2.21	0.41
1:A:375:ALA:O	1:A:379:ARG:HG3	2.21	0.41
1:B:360:LEU:HD22	1:B:364:VAL:HG23	2.03	0.41
1:B:361:PRO:HA	1:B:520:PHE:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	272/276~(99%)	264 (97%)	7 (3%)	1 (0%)	34 30
1	В	272/276 (99%)	252 (93%)	15 (6%)	5 (2%)	8 3
All	All	544/552 (99%)	516 (95%)	22 (4%)	6 (1%)	14 8

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	277	CYS
1	В	299	PRO
1	В	425	PRO
1	В	418	ALA
1	A	404	ASP



Continued from previous page...

Mol	Chain	Res	Type
1	В	417	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	236/236 (100%)	225 (95%)	11 (5%)	26	22	
1	В	236/236 (100%)	221 (94%)	15 (6%)	17	13	
All	All	472/472 (100%)	446 (94%)	26 (6%)	21	17	

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	265	GLU
1	A	272	LYS
1	A	297	LEU
1	A	346	LEU
1	A	350	LEU
1	A	398	LEU
1	A	414	ASN
1	A	423	LYS
1	A	426	ILE
1	A	491	LEU
1	A	533	LEU
1	В	268	ARG
1	В	275	GLN
1	В	278	PHE
1	В	280	GLU
1	В	299	PRO
1	В	301	THR
1	В	304	PRO
1	В	346	LEU
1	В	350	LEU
1	В	360	LEU
1	В	423	LYS



Continued from previous page...

Mol	Chain	Res	Type
1	В	424	PHE
1	В	425	PRO
1	В	426	ILE
1	В	477	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	287	ASN
1	A	468	ASN
1	A	526	GLN
1	В	381	ASN
1	В	420	GLN
1	В	526	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trimo	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Des	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	eles
MIOI	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2													
1	PTR	В	335	1	15,16,17	1.16	1 (6%)	19,22,24	1.49	3 (15%)													
1	PTR	В	416	1	15,16,17	1.05	1 (6%)	19,22,24	1.14	1 (5%)													
1	PTR	A	416	1	15,16,17	1.36	2 (13%)	19,22,24	1.07	1 (5%)													
1	PTR	A	335	1	15,16,17	1.02	1 (6%)	19,22,24	1.20	2 (10%)													

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	В	335	1	-	2/10/11/13	0/1/1/1
1	PTR	В	416	1	-	1/10/11/13	0/1/1/1
1	PTR	A	416	1	-	0/10/11/13	0/1/1/1
1	PTR	A	335	1	-	4/10/11/13	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$Ideal(\AA)$
1	A	416	PTR	P-OH	3.15	1.64	1.59
1	В	335	PTR	P-OH	2.42	1.63	1.59
1	В	416	PTR	P-OH	2.17	1.62	1.59
1	A	416	PTR	OH-CZ	2.14	1.45	1.40
1	A	335	PTR	P-OH	2.03	1.62	1.59

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
1	В	335	PTR	CB-CA-C	-3.17	105.53	111.47
1	A	335	PTR	CD2-CE2-CZ	-2.48	116.70	119.73
1	В	335	PTR	CD2-CE2-CZ	-2.48	116.71	119.73
1	В	416	PTR	CD2-CE2-CZ	-2.42	116.78	119.73
1	A	335	PTR	CB-CA-C	-2.13	107.48	111.47
1	A	416	PTR	CD2-CE2-CZ	-2.11	117.15	119.73
1	В	335	PTR	CG-CB-CA	2.09	118.34	114.10

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	335	PTR	CA-CB-CG-CD1
1	A	335	PTR	CA-CB-CG-CD2
1	A	335	PTR	N-CA-CB-CG
1	В	335	PTR	CA-CB-CG-CD1
1	В	335	PTR	CA-CB-CG-CD2
1	A	335	PTR	C-CA-CB-CG
1	В	416	PTR	C-CA-CB-CG

There are no ring outliers.

3 monomers are involved in 4 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	В	416	PTR	1	0
1	A	416	PTR	1	0
1	A	335	PTR	2	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

