

wwPDB X-ray Structure Validation Summary Report (i)

Apr 10, 2023 – 08:03 PM EDT

PDB ID	:	1Z1C
Title	:	Structural Determinants of Tissue Tropism and In Vivo Pathogenicity for the
		Parvovirus Minute virus of Mice
Authors	:	Kontou, M.; Govindasamy, L.; Nam, H.J.; Bryant, N.; Llamas-Saiz, A.L.;
		Foces-Foces, C.; Hernando, E.; Rubio, M.P.; McKenna, R.; Almendral, J.M.;
		Agbandje-McKenna, M.
Deposited on	:	2005-03-03
Resolution	:	3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)		
Validation Pipeline (wwPDB-VP)	:	2.32.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Q	uality of chain	
1	В	12		92%	8%
2	С	9		100%	
3	А	587	39%	42%	11% • 6%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a DNA chain called 5'-D(*AP*TP*CP*CP*TP*CP*TP*AP*TP*CP*AP*C)-3'.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace
1	В	12	Total 235	C 115	N 38	0 71	Р 11	0	0	0

• Molecule 2 is a DNA chain called 5'-D(*AP*CP*AP*CP*CP*AP*AP*AP*A)-3'.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	С	0	Total	С	Ν	Ο	Р	0	0	0
	U	9	180	87	39	46	8	0	0	0

• Molecule 3 is a protein called Coat protein VP2.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
3	А	549	Total 4328	C 2727	N 751	O 830	S 20	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
А	366	MET	VAL	SEE REMARK 999	UNP P07302
А	455	THR	ALA	SEE REMARK 999	UNP P07302

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	2	Total Ca 2 2	0	0

• Molecule 5 is 2'-DEOXYADENOSINE-5'-MONOPHOSPHATE (three-letter code: D5M) (formula: $C_{10}H_{14}N_5O_6P$).





Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
5	В	1	Total	С	Ν	Ο	0	0
0	D	L	18	10	5	3	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	5	Total O 5 5	0	0
6	С	11	Total O 11 11	0	0
6	А	85	Total O 85 85	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: 5'-D(*AP*TP*CP*CP*TP*CP*TP*AP*TP*CP*AP*C)-3'

Chain B:	92%		8%
A3 T4 C5 C6 C6 T7 T1 T11 C12 C12 C12 C12 C12			
• Molecule 2: $5'-D(3)$	*AP*CP*AP*CP*CP*	AP*AP*AP*A)-3'	
Chain C:	10	0%	
A15 C16 A17 C18 C19 C19 A21 A22 A22 A23			
• Molecule 3: Coat	protein VP2		
Chain A:	39%	42%	11% • 6%
MET SER SER SER GLY CLN PRO CLN PRO CLN ASP ASN ASN ASN ASN ASN ASN	SER ALA ALA ALA ARC ARC ARC ARC ARC ARC ARC CLY CLY CLY CLY CLY CLY CLY CLY CLY CL	01.Y 01.Y 01.Y 01.Y 01.Y 01.Y 01.Y 01.Y	048 049 049 0450 0451 151 155 055 059 059 059 059 059 059 163
A65 L66 L66 A67 A67 L68 R69 L70 L70 L70 L73 L73 M75 M75 K77 S78 S78	N8/0 184 184 184 188 188 188 188 188 188 188	M98 499 8100 8100 8100 8112 8112 8112 8112 1114 1114 1114 1114	A116 A116 A116 A118 V120 V121 V122 C123 C123 C123 D127 D127 V128 V129
M 35 1138 1138 1140 1144 1143 1144 1143 1144 1143 1148 1148	VII55 VII56 QII59 QII59 AI64 TI67 NI66 NI70 NI70	A179 V180 V180 S182 S182 N183 N183 N183 T185 T185 P189 P199 P199 P199	M195 M195 M195 F197 F197 F200 F200 F200 F200 F200 F200 F200 F20
A208 8209 7210 7211 7211 7215 7215 7215 7216 7216 7216 7216 7216 7219 7220 7220 7220 7220	Y225 Y225 P226 P228 P228 Q28 P229 H228 N235 N235 N235 N235 R223 P240 T239	2946 1260 1260 1265 1265 1265 1265 1265 1265 1265 1265	228 N275 N275 P276 P276 1280 1280 1280 1280 1280 1280 1280 1280
q288 (2390 (2390 (2392) (2394 (2394 (2394 (2394 (2396 (2396 (2396 (2396 (2396 (2397) (2303) (1307 1307 1309 1309 1310 1325 1325 1325 1325 1325 1325 1325 1325	A334 A335 Q335 Q337 Q337 Q340 Q340 P344 P344 F345 F345 F345 F345	3.945 R3450 A3550 P3551 P355 P355 P355 P355 P355 C3555 A356 C3555 M366 B367 B367 B367
E369 18371 18371 18371 18371 1837 1838 1838	4000 4000 4000 4000 4000 4000 4000 8400 8400 8400 8400 8400 8400 8400 8400 8400 8400 8400 8400	1409 1410 1411 1414 1414 1415 1419 1419 1419 1420 1420 1421 1420 1422	1425 1426 1427 1429 1429 1433 1433 1433 1433 1433 1433 1433 143
F443 5444 5444 5449 5449 5449 1453 F456 F456 F463 F464	4460 0467 0467 1468 1468 1468 1473 1473 1473 1473 1473 1473 1473 1473	H482 1483 1484 1484 P485 P485 F487 K490 K490 P494 Q495 Q495	R500 R501 L501 L501 P503 P503 N512 N513 N513 A514 T515 A514 T515 S517 S517 R518 R518
		PROTEIN DATA BANK	

1519 1523 1524 1523 1524 1524 1524 1524 1525 1524 1531 1533 1533 1533 1533 1533 1533 1533 1533 1533 1533 1533 1533 1544 1545 1545 1546 1555 1565 1565 1565 1566 1567 1567 1566 1567 1578 1578 1578 1578 1578 1578 1578 1578 1578 1578 1578 1578 1578 1578



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	448.70Å 416.70Å 305.30Å	Depositor
a, b, c, α , β , γ	90.00° 95.80° 90.00°	Depositor
Resolution (Å)	20.00 - 3.50	Depositor
% Data completeness	(Not available) (20.00-3.50)	Depositor
(in resolution range)	(100 available) (20.00 0.00)	Depositor
R_{merge}	0.15	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.325 , 0.329	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4864	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, $\mathrm{D5M}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain Bor		nd lengths	Bond angles	
10101	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	В	0.74	0/261	0.84	0/399
2	С	0.74	0/203	0.90	0/310
3	А	1.14	8/4452~(0.2%)	1.08	31/6088~(0.5%)
All	All	1.11	8/4916~(0.2%)	1.06	31/6797~(0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1
3	А	0	3
All	All	0	4

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
3	А	586	THR	N-CA	41.36	2.29	1.46
3	А	418	PRO	N-CD	37.59	2.00	1.47
3	А	234	HIS	N-CA	26.68	1.99	1.46
3	А	227	ASN	N-CA	13.97	1.74	1.46
3	А	581	PRO	N-CD	11.42	1.63	1.47

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	585	ASN	C-N-CA	-21.02	69.14	121.70
3	А	418	PRO	N-CA-CB	18.11	125.03	103.30
3	А	418	PRO	CA-N-CD	-17.03	87.65	111.50

Continued on next page...



	Chain	1	1 5	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	233	GLU	C-N-CA	-14.68	85.01	121.70
3	А	586	THR	N-CA-CB	-14.07	83.57	110.30

Continued from previous page...

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	А	194	SER	Mainchain
3	А	363	THR	Peptide
3	А	417	ALA	Peptide
1	В	7	DT	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	235	0	138	29	0
2	С	180	0	101	18	0
3	А	4328	0	4124	383	0
4	В	2	0	0	0	0
5	В	18	0	12	3	0
6	А	85	0	0	1	0
6	В	5	0	0	0	0
6	С	11	0	0	0	0
All	All	4864	0	4375	419	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 46.

The worst 5 of 419 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:227:ASN:N	3:A:227:ASN:CA	1.74	1.49
3:A:380:LYS:NZ	3:A:386:TRP:O	1.56	1.34
3:A:234:HIS:N	3:A:234:HIS:CA	1.99	1.26

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:385:ASN:OD1	3:A:386:TRP:N	1.67	1.24
3:A:418:PRO:N	3:A:418:PRO:CD	2.00	1.23

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	А	547/587~(93%)	432 (79%)	74 (14%)	41 (8%)	1 11

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	А	218	ASP
3	А	226	GLU
3	А	233	GLU
3	А	380	LYS
3	А	381	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	А	472/492~(96%)	423 (90%)	49 (10%)	7 31

 $5~{\rm of}~49$ residues with a non-rotameric side chain are listed below:



Mol	Chain	Res	Type
3	А	385	ASN
3	А	440	ASP
3	А	386	TRP
3	А	411	ASP
3	А	476	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
3	А	381	GLN
3	А	442	HIS
3	А	573	GLN
3	А	433	ASN
3	А	448	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	oe Chain F		Link	Bond lengths			Bond angles		
	moi Type	Cham	Res		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
5	D5M	В	231	-	18,20,24	0.59	0	$17,\!29,\!36$	0.74	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	D5M	В	231	-	-	0/2/18/22	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	В	231	D5M	C5-C6-N6	2.15	123.61	120.35

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	В	231	D5M	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

