



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 17, 2023 – 03:48 AM EST

PDB ID : 2AML  
Title : Crystal structure of Lmo0035 protein (46906266) from LISTERIA MONOCYTOGENES 4b F2365 at 1.50 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2005-08-09  
Resolution : 1.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	373	91% (green), 7% (yellow), 2% (orange), 0% (red), 0% (grey)
1	B	373	93% (green), 5% (yellow), 2% (orange), 0% (red), 0% (grey)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	363	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6866 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SIS domain protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	366	2926	1878	472	565	2	9	0	14	0
1	B	361	2853	1831	453	558	2	9	0	7	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MSE	-	expression tag	UNP Q4EK90
A	-10	GLY	-	expression tag	UNP Q4EK90
A	-9	SER	-	expression tag	UNP Q4EK90
A	-8	ASP	-	expression tag	UNP Q4EK90
A	-7	LYS	-	expression tag	UNP Q4EK90
A	-6	ILE	-	expression tag	UNP Q4EK90
A	-5	HIS	-	expression tag	UNP Q4EK90
A	-4	HIS	-	expression tag	UNP Q4EK90
A	-3	HIS	-	expression tag	UNP Q4EK90
A	-2	HIS	-	expression tag	UNP Q4EK90
A	-1	HIS	-	expression tag	UNP Q4EK90
A	0	HIS	-	expression tag	UNP Q4EK90
A	1	MSE	MET	modified residue	UNP Q4EK90
A	5	MSE	MET	modified residue	UNP Q4EK90
A	6	MSE	MET	modified residue	UNP Q4EK90
A	15	MSE	MET	modified residue	UNP Q4EK90
A	159	MSE	MET	modified residue	UNP Q4EK90
A	251	MSE	MET	modified residue	UNP Q4EK90
A	320	MSE	MET	modified residue	UNP Q4EK90
A	351	MSE	MET	modified residue	UNP Q4EK90
B	-11	MSE	-	expression tag	UNP Q4EK90
B	-10	GLY	-	expression tag	UNP Q4EK90
B	-9	SER	-	expression tag	UNP Q4EK90
B	-8	ASP	-	expression tag	UNP Q4EK90
B	-7	LYS	-	expression tag	UNP Q4EK90

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	ILE	-	expression tag	UNP Q4EK90
B	-5	HIS	-	expression tag	UNP Q4EK90
B	-4	HIS	-	expression tag	UNP Q4EK90
B	-3	HIS	-	expression tag	UNP Q4EK90
B	-2	HIS	-	expression tag	UNP Q4EK90
B	-1	HIS	-	expression tag	UNP Q4EK90
B	0	HIS	-	expression tag	UNP Q4EK90
B	1	MSE	MET	modified residue	UNP Q4EK90
B	5	MSE	MET	modified residue	UNP Q4EK90
B	6	MSE	MET	modified residue	UNP Q4EK90
B	15	MSE	MET	modified residue	UNP Q4EK90
B	159	MSE	MET	modified residue	UNP Q4EK90
B	251	MSE	MET	modified residue	UNP Q4EK90
B	320	MSE	MET	modified residue	UNP Q4EK90
B	351	MSE	MET	modified residue	UNP Q4EK90

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	557	Total 558	O 558	0	1
5	B	502	Total 503	O 503	0	1


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SIS domain protein



- Molecule 1: SIS domain protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.31Å 101.40Å 120.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.45 – 1.50 29.46 – 1.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.45-1.50) 100.0 (29.46-1.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.56 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.2.0013	Depositor
R, $R_{free}$	0.140 , 0.168 0.147 , 0.172	Depositor DCC
$R_{free}$ test set	5535 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.5	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 41.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6866	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/3052	0.80	3/4126 (0.1%)
1	B	0.79	0/2938	0.84	2/3974 (0.1%)
All	All	0.78	0/5990	0.82	5/8100 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	ARG	NE-CZ-NH2	-6.45	117.07	120.30
1	B	67	ARG	NE-CZ-NH1	5.65	123.13	120.30
1	B	303	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	173	ASP	CB-CG-OD1	5.26	123.03	118.30
1	A	201	PHE	CB-CG-CD1	-5.18	117.18	120.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2926	0	2847	19	0
1	B	2853	0	2793	7	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	0	8	5	0
3	B	6	0	8	1	0
4	B	4	0	3	0	0
5	A	558	0	0	4	0
5	B	503	0	0	2	0
All	All	6866	0	5659	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-1[B]:HIS:ND1	3:A:363:GOL:C1	2.29	0.96
1:A:-1[B]:HIS:ND1	3:A:363:GOL:H11	1.84	0.91
1:A:-1[B]:HIS:CE1	3:A:363:GOL:H11	2.16	0.81
1:A:20:LEU:HD21	1:A:185[B]:SER:OG	1.86	0.76
1:A:-1[B]:HIS:ND1	3:A:363:GOL:H12	2.00	0.75
1:B:272:ALA:HB3	3:B:364:GOL:H32	1.73	0.69
1:A:119[B]:LEU:HD23	1:A:135:LEU:HB2	1.76	0.66
1:A:214:LYS:HD3	5:A:725:HOH:O	1.97	0.65
1:B:25:THR:HG23	5:B:813:HOH:O	2.00	0.61
1:B:106:GLU:O	1:B:110[B]:LYS:HD3	2.03	0.58
1:A:33:LEU:HD11	1:A:133:ILE:HG12	1.86	0.58
1:A:186[B]:ARG:HD2	5:A:812:HOH:O	2.04	0.57
1:A:119[A]:LEU:HD11	1:A:157:THR:HG21	1.90	0.52
1:A:42:LEU:HB2	1:A:86[B]:LEU:HD21	1.91	0.52
1:A:20:LEU:CD2	1:A:185[B]:SER:OG	2.59	0.51
1:A:186[B]:ARG:NE	5:A:863:HOH:O	2.44	0.51
1:B:275[A]:GLU:HG3	1:B:276:ARG:N	2.33	0.44
1:B:57[B]:LYS:HE3	1:B:58:TYR:CZ	2.54	0.43
1:A:345:THR:HG22	5:A:804:HOH:O	2.18	0.42
1:B:273:VAL:HA	1:B:275[B]:GLU:OE1	2.20	0.42
1:A:2:LYS:HD3	1:A:361:ALA:OXT	2.20	0.42
1:A:86[A]:LEU:HD21	1:A:89:VAL:CG2	2.50	0.41
1:A:352:LYS:CD	5:B:796:HOH:O	2.68	0.41
1:A:119[A]:LEU:HG	1:A:137:ILE:CG2	2.50	0.41
1:A:271[B]:SER:OG	1:A:272:ALA:N	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	378/373 (101%)	368 (97%)	9 (2%)	1 (0%)	41	18
1	B	366/373 (98%)	358 (98%)	8 (2%)	0	100	100
All	All	744/746 (100%)	726 (98%)	17 (2%)	1 (0%)	51	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	170	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/312 (102%)	316 (99%)	3 (1%)	78	61
1	B	308/312 (99%)	304 (99%)	4 (1%)	69	44
All	All	627/624 (100%)	620 (99%)	7 (1%)	71	53

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	ARG
1	A	233	LYS
1	A	330	HIS
1	B	107	ARG
1	B	233	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	248	GLU
1	B	330	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ACT	B	362	-	3,3,3	0.86	0	3,3,3	1.74	2 (66%)
3	GOL	A	363	-	5,5,5	0.66	0	5,5,5	1.39	1 (20%)
3	GOL	B	364	-	5,5,5	0.81	0	5,5,5	0.57	0
2	PO4	B	363	-	4,4,4	0.38	0	6,6,6	1.21	1 (16%)
2	PO4	A	362	-	4,4,4	0.83	0	6,6,6	2.62	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	364	-	-	4/4/4/4	-
3	GOL	A	363	-	-	4/4/4/4	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	362	PO4	O4-P-O3	-5.11	91.56	107.97
3	A	363	GOL	O1-C1-C2	2.65	122.92	110.20
2	A	362	PO4	O3-P-O1	2.56	120.27	110.89
2	B	363	PO4	O4-P-O1	-2.38	102.20	110.89
4	B	362	ACT	OXT-C-CH3	2.17	124.16	115.18
4	B	362	ACT	OXT-C-O	-2.09	114.36	122.05

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	363	GOL	O1-C1-C2-C3
3	B	364	GOL	O1-C1-C2-C3
3	B	364	GOL	C1-C2-C3-O3
3	B	364	GOL	O2-C2-C3-O3
3	A	363	GOL	C1-C2-C3-O3
3	A	363	GOL	O1-C1-C2-O2
3	A	363	GOL	O2-C2-C3-O3
3	B	364	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	363	GOL	5	0
3	B	364	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.