



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 28, 2024 – 07:43 pm GMT

PDB ID : 2C3V  
Title : Structure of iodinated CBM25 from Bacillus halodurans amylase  
Authors : Boraston, A.B.; Healey, M.; Klassen, J.; Ficko-Blean, E.; Lammerts van Bueren, A.; Law, V.  
Deposited on : 2005-10-12  
Resolution : 1.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2247 (1.40-1.40)
Clashscore	180529	2446 (1.40-1.40)
Ramachandran outliers	177936	2398 (1.40-1.40)
Sidechain outliers	177891	2397 (1.40-1.40)
RSRZ outliers	164620	2246 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	102	
2	B	102	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IOD	B	2[B]	-	-	X	-
3	IOD	B	3[A]	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1695 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-AMYLASE G-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	I	N	O			
1	A	94	751	471	4	126	150	0	2	0

- Molecule 2 is a protein called ALPHA-AMYLASE G-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	94	724	453	125	144	2	0	1	0

- Molecule 3 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	I	0	3
			8	8		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	133	Total	O	0	2
			133	133		
4	B	79	Total	O	0	2
			79	79		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ALPHA-AMYLASE G-6



- Molecule 2: ALPHA-AMYLASE G-6



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	30.20Å 41.82Å 68.18Å 90.00° 97.18° 90.00°	Depositor
Resolution (Å)	67.42 – 1.39 67.42 – 1.39	Depositor EDS
% Data completeness (in resolution range)	88.0 (67.42-1.39) 88.0 (67.42-1.39)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 1.39Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.201 , 0.250 0.205 , 0.252	Depositor DCC
$R_{free}$ test set	1542 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.9	Xtrriage
Anisotropy	0.137	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1695	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, TYI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.97	0/734	0.96	0/998
2	B	0.87	0/746	1.08	4/1016 (0.4%)
All	All	0.92	0/1480	1.02	4/2014 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	170	ASP	CB-CG-OD2	7.05	124.64	118.30
2	B	160	ASP	CB-CG-OD2	5.33	123.09	118.30
2	B	85	ASP	CB-CG-OD2	5.17	122.96	118.30
2	B	88	ASP	CB-CG-OD2	5.12	122.91	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	113	THR	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	751	0	680	9	0
2	B	724	0	669	22	0
3	B	8	0	0	8	2
4	A	133	0	0	0	2
4	B	79	0	0	4	0
All	All	1695	0	1349	33	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:2[B]:IOD:I	4:B:2038:HOH:O	2.43	1.06
1:A:141:ARG:HD2	1:A:160:ASP:OD1	1.74	0.88
1:A:145:ASN:HD22	1:A:147:GLY:H	1.22	0.86
3:B:2[C]:IOD:I	4:B:2079:HOH:O	2.74	0.75
2:B:145:ASN:HD22	2:B:147:GLY:H	1.37	0.70
2:B:98:THR:HG22	2:B:99:HIS:CD2	2.28	0.68
2:B:92:TYR:HE1	3:B:2[B]:IOD:I	2.51	0.64
2:B:127:TYR:HD1	3:B:3[A]:IOD:I	2.53	0.61
2:B:154:ASN:HD21	2:B:157:ARG:NH2	1.98	0.60
1:A:145:ASN:ND2	1:A:147:GLY:H	1.98	0.59
2:B:92:TYR:CE1	3:B:2[B]:IOD:I	3.27	0.58
1:A:141:ARG:CD	1:A:160:ASP:OD1	2.47	0.58
2:B:155:GLN:NE2	4:B:2063:HOH:O	2.36	0.58
1:A:87:THR:HG22	1:A:134:ALA:O	2.03	0.58
2:B:122:SER:C	4:B:2038:HOH:O	2.41	0.58
1:A:87:THR:CG2	1:A:134:ALA:O	2.56	0.54
2:B:146:ASN:HD22	2:B:146:ASN:H	1.56	0.53
2:B:114:LEU:O	2:B:115:PRO:C	2.47	0.53
2:B:122:SER:HB2	3:B:2[B]:IOD:I	2.80	0.51
2:B:132:ILE:HD13	2:B:140:LEU:HD21	1.94	0.50
2:B:127:TYR:CD1	3:B:3[A]:IOD:I	3.34	0.48
2:B:92:TYR:HA	2:B:128:VAL:O	2.14	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ASN:ND2	1:A:148:SER:H	2.12	0.47
2:B:146:ASN:HD22	2:B:146:ASN:N	2.13	0.47
1:A:146:ASN:ND2	1:A:150:GLN:H	2.12	0.46
2:B:146:ASN:ND2	2:B:150:GLN:H	2.14	0.46
2:B:102:ILE:HB	2:B:119:LEU:HD21	1.97	0.45
2:B:92:TYR:OH	3:B:2[A]:IOD:I	3.02	0.45
2:B:145:ASN:HD22	2:B:147:GLY:N	2.11	0.44
2:B:146:ASN:HD21	2:B:150:GLN:H	1.66	0.44
2:B:97:TRP:CE2	2:B:146:ASN:HB3	2.53	0.44
1:A:94:LYS:HB2	1:A:127[A]:TYI:CE2	2.48	0.43
2:B:98:THR:CG2	2:B:99:HIS:CD2	3.01	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1[C]:IOD:I	4:A:2022:HOH:O[1_565]	1.23	0.97
3:B:1[B]:IOD:I	4:A:2020:HOH:O[1_565]	1.75	0.45

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	91/102 (89%)	88 (97%)	3 (3%)	0	100	100
2	B	90/102 (88%)	82 (91%)	8 (9%)	0	100	100
All	All	181/204 (89%)	170 (94%)	11 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/80 (95%)	75 (99%)	1 (1%)	65	38
2	B	76/82 (93%)	69 (91%)	7 (9%)	7	0
All	All	152/162 (94%)	144 (95%)	8 (5%)	19	2

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	ASN
2	B	94	LYS
2	B	114	LEU
2	B	119	LEU
2	B	140	LEU
2	B	146	ASN
2	B	150	GLN
2	B	175	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	GLN
1	A	145	ASN
1	A	146	ASN
2	B	99	HIS
2	B	145	ASN
2	B	146	ASN
2	B	150	GLN
2	B	155	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	TYI	A	124	1	13,14,15	1.41	2 (15%)	16,19,21	1.21	2 (12%)
1	TYI	A	127[B]	-	12,13,15	2.11	2 (16%)	14,17,21	1.44	3 (21%)
1	TYI	A	127[A]	-	12,13,15	2.07	2 (16%)	14,17,21	1.75	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TYI	A	124	1	-	0/5/6/8	0/1/1/1
1	TYI	A	127[B]	-	-	0/5/6/8	0/1/1/1
1	TYI	A	127[A]	-	-	0/5/6/8	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	127[B]	TYI	CE2-CZ	5.24	1.48	1.39
1	A	127[A]	TYI	CE2-CZ	4.81	1.48	1.39
1	A	127[B]	TYI	CZ-CE1	4.58	1.49	1.39
1	A	127[A]	TYI	CZ-CE1	4.50	1.48	1.39
1	A	124	TYI	CZ-CE2	3.90	1.48	1.40
1	A	124	TYI	CZ-CE1	2.14	1.45	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	127[A]	TYI	CZ-CE1-I1	4.35	124.12	119.81
1	A	124	TYI	CZ-CE1-I1	-3.23	114.45	119.42
1	A	124	TYI	CD1-CE1-I1	2.71	123.60	118.61

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	127[B]	TYI	CB-CA-C	-2.29	107.17	111.47
1	A	127[A]	TYI	CE2-CZ-CE1	-2.21	117.18	119.27
1	A	127[B]	TYI	CD2-CG-CD1	2.17	121.58	118.54
1	A	127[B]	TYI	CZ-CE1-I1	2.16	121.95	119.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	127[A]	TYI	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	92/102 (90%)	0.11	3 (3%) 49 50	12, 17, 24, 34	1 (1%)
2	B	94/102 (92%)	1.64	33 (35%) 1 1	14, 28, 38, 40	1 (1%)
All	All	186/204 (91%)	0.88	36 (19%) 4 3	12, 21, 37, 40	2 (1%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	114	LEU	7.2
2	B	117	VAL	4.1
2	B	86	ALA	3.8
2	B	174	LEU	3.7
2	B	128	VAL	3.6
2	B	127	TYR	3.6
2	B	115	PRO	3.4
2	B	82	ALA	3.2
2	B	99	HIS	3.1
2	B	157	ARG	3.0
2	B	167	THR	3.0
2	B	92	TYR	2.8
2	B	136	GLU	2.7
2	B	148	SER	2.6
1	A	86	ALA	2.5
2	B	177	THR	2.5
2	B	120	THR	2.5
2	B	119	LEU	2.5
2	B	156	GLY	2.4
2	B	147	GLY	2.4
1	A	85	ASP	2.4
2	B	97	TRP	2.4
2	B	170	ASP	2.4
2	B	98	THR	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	165	VAL	2.3
2	B	118	PRO	2.3
2	B	116	GLY	2.2
2	B	137	GLY	2.2
2	B	173	ILE	2.2
2	B	95	THR	2.2
2	B	81[A]	MET	2.2
2	B	101	HIS	2.2
2	B	171	GLY	2.1
2	B	169	ALA	2.1
2	B	113	THR	2.1
1	A	87	THR	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TYI	A	127[A]	13/15	0.84	0.23	15,48,97,97	10
1	TYI	A	127[B]	13/15	0.84	0.23	15,53,100,103	10
1	TYI	A	124	14/15	1.00	0.03	16,18,20,23	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IOD	B	3[A]	1/1	0.74	0.17	48,48,48,48	1
3	IOD	B	3[B]	1/1	0.74	0.17	36,36,36,36	1
3	IOD	B	1[C]	1/1	0.97	0.08	27,27,27,27	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	IOD	B	1[A]	1/1	0.97	0.08	35,35,35,35	1
3	IOD	B	1[B]	1/1	0.97	0.08	33,33,33,33	1
3	IOD	B	2[C]	1/1	0.98	0.06	32,32,32,32	1
3	IOD	B	2[A]	1/1	0.98	0.06	27,27,27,27	1
3	IOD	B	2[B]	1/1	0.98	0.06	29,29,29,29	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.