



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 4, 2023 – 06:57 am GMT

PDB ID : 2CIX
Title : chloroperoxidase complexed with cyclopentanedione
Authors : Kuhnel, K.; Blankenfeldt, W.; Terner, J.; Schlichting, I.
Deposited on : 2006-03-26
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			%	
1	A	299	91%	9%
2	B	3	100%	
3	C	2	50%	50%
4	D	2	100%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BMA	B	3	-	-	-	X

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 2975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHLOROPEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	299	2359	1494	391	467	7	0	7	0

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	39	22	2	15	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	2	Total	C	O	0	0	0
			22	12	10			

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mn	0	0
			1	1		

- Molecule 6 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
7	A	1	14	8	1	5	0	0

- Molecule 8 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	A	1	11	6	5	0	0
8	A	1	11	6	5	0	0
8	A	1	11	6	5	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			11	6	5		
8	A	1	Total	C	O	0	0
			11	6	5		
8	A	1	Total	C	O	0	0
			11	6	5		
8	A	1	Total	C	O	0	0
			11	6	5		
8	A	1	Total	C	O	0	0
			11	6	5		
8	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 9 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 10 is CYCLOPENTANE-1,3-DIONE (three-letter code: CEJ) (formula: C₅H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total C O 7 5 2	0	0

- Molecule 11 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	3	Total Br 3 3	0	0

- Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	345	Total O 345 345	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CHLOROPEROXIDASE

Chain A:  %




- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  50% 50%



- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose

Chain D:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	57.44Å 150.35Å 99.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.73 – 1.80 19.72 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.73-1.80) 98.9 (19.72-1.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.178 , 0.217 0.176 , 0.214	Depositor DCC
R_{free} test set	1998 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	19.9	Xtrriage
Anisotropy	0.540	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2975	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MN, BR, CEJ, MAN, HEM, PCA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/2435	0.65	0/3338

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2359	0	2173	20	0
2	B	39	0	34	0	0
3	C	28	0	25	2	0
4	D	22	0	19	0	0
5	A	1	0	0	0	0
6	A	43	0	30	5	0
7	A	14	0	13	0	0
8	A	110	0	100	2	0
9	A	4	0	6	0	0
10	A	7	0	6	1	0
11	A	3	0	0	0	0
12	A	345	0	0	5	0
All	All	2975	0	2406	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176[A]:SER:OG	12:A:2212:HOH:O	2.07	0.72
1:A:216:ASN:HB3	1:A:294:LEU:HD11	1.70	0.71
1:A:18:VAL:O	1:A:43[A]:HIS:HE1	1.77	0.68
1:A:1:GLU:H	1:A:297:ASN:ND2	1.93	0.66
6:A:1300:HEM:HMB2	6:A:1300:HEM:HBB2	1.84	0.58
1:A:118:VAL:HB	1:A:123:ASP:HB2	1.92	0.51
1:A:179:ILE:HG23	10:A:1320:CEJ:H102	1.92	0.51
1:A:159:GLN:NE2	8:A:1312:MAN:O4	2.44	0.51
6:A:1300:HEM:HMC1	6:A:1300:HEM:HBC2	1.94	0.50
1:A:211[A]:LYS:HG2	3:C:1:NAG:H81	1.93	0.50
6:A:1300:HEM:HBB2	6:A:1300:HEM:CMB	2.42	0.49
1:A:79:CYS:SG	1:A:87[B]:CYS:SG	3.05	0.48
1:A:284:ASN:H	1:A:284:ASN:HD22	1.60	0.47
1:A:57:PHE:CE1	6:A:1300:HEM:HMC2	2.49	0.47
1:A:162[B]:SER:O	1:A:166:GLU:HG3	2.14	0.46
1:A:43[B]:HIS:HE1	12:A:2070:HOH:O	1.97	0.46
6:A:1300:HEM:HBC2	6:A:1300:HEM:CMC	2.46	0.45
1:A:51:GLU:CD	12:A:2083:HOH:O	2.55	0.44
1:A:78:VAL:HG21	1:A:103:PHE:HB2	2.00	0.43
12:A:2309:HOH:O	3:C:2:NAG:H82	2.20	0.42
1:A:155:GLU:HG3	8:A:1311:MAN:O4	2.20	0.41
1:A:284:ASN:H	1:A:284:ASN:ND2	2.17	0.41
1:A:147:HIS:HA	1:A:205:VAL:O	2.21	0.41
1:A:294:LEU:HB3	1:A:298:ASP:HB2	2.03	0.41
1:A:13[B]:ASN:ND2	12:A:2034:HOH:O	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PCA	A	0	1	7,8,9	1.74	1 (14%)	9,10,12	1.66	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	0	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	0	PCA	CD-N	4.32	1.45	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	0	PCA	CA-N-CD	-2.90	103.64	113.58
1	A	0	PCA	OE-CD-CG	-2.31	122.74	126.76
1	A	0	PCA	CB-CA-N	2.21	109.65	103.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	0.66	0	17,19,21	1.28	3 (17%)
2	NAG	B	2	2	14,14,15	0.47	0	17,19,21	1.07	2 (11%)
2	BMA	B	3	2	11,11,12	0.61	0	15,15,17	1.22	3 (20%)
3	NAG	C	1	3,1	14,14,15	0.71	0	17,19,21	0.99	0
3	NAG	C	2	3	14,14,15	0.62	0	17,19,21	1.59	4 (23%)
4	MAN	D	1	4,1	11,11,12	0.60	0	15,15,17	1.03	1 (6%)
4	MAN	D	2	4	11,11,12	0.67	0	15,15,17	1.19	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
4	MAN	D	1	4,1	-	2/2/19/22	0/1/1/1
4	MAN	D	2	4	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C2-N2-C7	-4.18	116.95	122.90
4	D	2	MAN	C1-O5-C5	2.85	116.05	112.19
3	C	2	NAG	C4-C3-C2	2.55	114.76	111.02
3	C	2	NAG	C1-O5-C5	-2.55	108.74	112.19
2	B	2	NAG	C1-O5-C5	-2.53	108.77	112.19
2	B	3	BMA	O5-C5-C6	2.35	110.89	107.20
2	B	3	BMA	C1-C2-C3	2.31	112.51	109.67
2	B	1	NAG	C1-O5-C5	2.30	115.30	112.19
3	C	2	NAG	O5-C5-C6	2.22	110.69	107.20
2	B	3	BMA	C2-C3-C4	2.19	114.69	110.89
2	B	1	NAG	C1-C2-N2	2.15	114.16	110.49
2	B	2	NAG	O5-C5-C6	2.12	110.53	107.20
2	B	1	NAG	O7-C7-C8	-2.09	118.17	122.06
4	D	1	MAN	O2-C2-C3	-2.02	106.09	110.14

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	2	MAN	C4-C5-C6-O6
4	D	2	MAN	O5-C5-C6-O6
4	D	1	MAN	O5-C5-C6-O6
4	D	1	MAN	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2	NAG	1	0
3	C	1	NAG	1	0

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	A	1301	1	14,14,15	0.46	0	17,19,21	1.28	2 (11%)
8	MAN	A	1314	1	11,11,12	0.55	0	15,15,17	1.73	1 (6%)
6	HEM	A	1300	5,1	41,50,50	1.86	7 (17%)	45,82,82	1.85	12 (26%)
8	MAN	A	1315	1	11,11,12	0.69	0	15,15,17	1.34	2 (13%)
8	MAN	A	1311	1	11,11,12	0.63	0	15,15,17	1.14	2 (13%)
8	MAN	A	1313	1	11,11,12	0.51	0	15,15,17	1.38	1 (6%)
9	EDO	A	1319	-	3,3,3	0.48	0	2,2,2	0.45	0
8	MAN	A	1310	1	11,11,12	0.67	0	15,15,17	1.32	2 (13%)
8	MAN	A	1308	1	11,11,12	0.57	0	15,15,17	0.87	1 (6%)
8	MAN	A	1318	1	11,11,12	0.69	0	15,15,17	1.19	1 (6%)
8	MAN	A	1309	1	11,11,12	0.64	0	15,15,17	0.87	0
10	CEJ	A	1320	-	5,7,7	0.93	0	9,9,9	4.76	7 (77%)
8	MAN	A	1307	1	11,11,12	0.54	0	15,15,17	0.91	1 (6%)
8	MAN	A	1312	1	11,11,12	0.52	0	15,15,17	1.53	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	1301	1	-	0/6/23/26	0/1/1/1
8	MAN	A	1314	1	-	0/2/19/22	0/1/1/1
6	HEM	A	1300	5,1	-	3/12/54/54	-
8	MAN	A	1315	1	-	2/2/19/22	0/1/1/1
8	MAN	A	1311	1	-	2/2/19/22	0/1/1/1
8	MAN	A	1313	1	-	2/2/19/22	0/1/1/1
9	EDO	A	1319	-	-	0/1/1/1	-
8	MAN	A	1310	1	-	0/2/19/22	0/1/1/1
8	MAN	A	1308	1	-	0/2/19/22	0/1/1/1
8	MAN	A	1318	1	-	1/2/19/22	0/1/1/1
8	MAN	A	1309	1	-	0/2/19/22	0/1/1/1
10	CEJ	A	1320	-	-	-	0/1/1/1
8	MAN	A	1307	1	-	0/2/19/22	0/1/1/1
8	MAN	A	1312	1	-	0/2/19/22	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1300	HEM	C3D-C2D	6.64	1.50	1.36
6	A	1300	HEM	C3C-CAC	4.32	1.56	1.47
6	A	1300	HEM	C3C-C2C	-3.58	1.35	1.40
6	A	1300	HEM	CAB-C3B	2.78	1.55	1.47
6	A	1300	HEM	CAA-C2A	2.48	1.55	1.52
6	A	1300	HEM	C1A-NA	2.31	1.40	1.36
6	A	1300	HEM	CMB-C2B	2.20	1.55	1.50

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	1320	CEJ	C1-C2-C10	8.54	114.98	109.36
10	A	1320	CEJ	C10-C7-C5	8.43	114.91	109.36
8	A	1314	MAN	C1-O5-C5	5.98	120.29	112.19
6	A	1300	HEM	C4D-ND-C1D	5.45	110.71	105.07
8	A	1313	MAN	C1-O5-C5	4.96	118.92	112.19
6	A	1300	HEM	C1B-NB-C4B	4.63	109.85	105.07
8	A	1312	MAN	C1-O5-C5	4.17	117.84	112.19
10	A	1320	CEJ	C1-C5-C7	-3.91	104.27	106.55
10	A	1320	CEJ	C5-C1-C2	-3.70	104.39	106.55
6	A	1300	HEM	CHB-C1B-NB	3.69	128.95	124.38
8	A	1310	MAN	C1-O5-C5	3.53	116.98	112.19
10	A	1320	CEJ	O3-C2-C10	-3.24	122.49	125.40
10	A	1320	CEJ	O8-C7-C10	-3.21	122.52	125.40
6	A	1300	HEM	CHC-C4B-NB	3.20	127.91	124.43
7	A	1301	NAG	O5-C5-C6	3.18	112.19	107.20
10	A	1320	CEJ	C2-C10-C7	-2.99	101.43	105.92
6	A	1300	HEM	C4C-CHD-C1D	2.94	126.43	122.56
8	A	1315	MAN	O5-C5-C6	2.88	111.72	107.20
6	A	1300	HEM	CMA-C3A-C4A	-2.84	124.10	128.46
8	A	1311	MAN	O5-C5-C6	2.62	111.31	107.20
8	A	1311	MAN	C1-O5-C5	2.58	115.69	112.19
8	A	1315	MAN	C1-O5-C5	2.58	115.68	112.19
8	A	1310	MAN	C2-C3-C4	2.52	115.26	110.89
8	A	1312	MAN	C1-C2-C3	2.43	112.65	109.67
6	A	1300	HEM	C2B-C1B-NB	-2.38	107.02	109.84
6	A	1300	HEM	CHD-C1D-ND	2.35	126.98	124.43
8	A	1318	MAN	C3-C4-C5	2.28	114.30	110.24
7	A	1301	NAG	C1-O5-C5	2.26	115.25	112.19
6	A	1300	HEM	C4A-C3A-C2A	2.20	108.53	107.00
6	A	1300	HEM	CAD-C3D-C4D	2.12	128.36	124.66
8	A	1307	MAN	C1-O5-C5	2.10	115.03	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	1308	MAN	C1-C2-C3	2.07	112.21	109.67
6	A	1300	HEM	CAA-CBA-CGA	-2.03	108.07	113.76
6	A	1300	HEM	C4B-CHC-C1C	2.02	125.23	122.56

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	A	1315	MAN	C4-C5-C6-O6
8	A	1311	MAN	O5-C5-C6-O6
8	A	1318	MAN	O5-C5-C6-O6
8	A	1315	MAN	O5-C5-C6-O6
8	A	1313	MAN	O5-C5-C6-O6
8	A	1311	MAN	C4-C5-C6-O6
8	A	1313	MAN	C4-C5-C6-O6
6	A	1300	HEM	CAA-CBA-CGA-O2A
6	A	1300	HEM	CAA-CBA-CGA-O1A
6	A	1300	HEM	C2A-CAA-CBA-CGA

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1300	HEM	5	0
8	A	1311	MAN	1	0
10	A	1320	CEJ	1	0
8	A	1312	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/299 (99%)	-0.27	3 (1%) 82 80	12, 19, 33, 42	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	275	THR	4.3
1	A	276	MET	2.4
1	A	291	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	0	8/9	0.96	0.10	17,18,19,19	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	B	3	11/12	0.64	0.41	48,53,56,57	0
3	NAG	C	2	14/15	0.74	0.37	37,40,43,44	0
4	MAN	D	2	11/12	0.81	0.29	42,44,46,48	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	B	2	14/15	0.85	0.13	19,28,36,43	0
4	MAN	D	1	11/12	0.91	0.17	34,40,43,45	0
3	NAG	C	1	14/15	0.97	0.06	16,18,22,28	0
2	NAG	B	1	14/15	0.97	0.06	18,22,25,25	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	MAN	A	1311	11/12	0.71	0.22	43,46,48,49	0
8	MAN	A	1314	11/12	0.72	0.30	47,51,52,52	0
8	MAN	A	1318	11/12	0.73	0.28	39,46,49,52	0
8	MAN	A	1310	11/12	0.78	0.26	35,39,42,42	0
8	MAN	A	1312	11/12	0.80	0.19	41,42,45,48	0
8	MAN	A	1315	11/12	0.81	0.27	43,48,50,53	0
8	MAN	A	1313	11/12	0.88	0.10	35,37,41,43	0
8	MAN	A	1307	11/12	0.92	0.11	29,34,36,37	0
7	NAG	A	1301	14/15	0.93	0.14	27,29,31,33	0
10	CEJ	A	1320	7/7	0.94	0.16	26,28,30,31	0
8	MAN	A	1308	11/12	0.95	0.08	21,23,25,26	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	EDO	A	1319	4/4	0.96	0.09	24,26,26,26	0
6	HEM	A	1300	43/43	0.98	0.07	10,12,14,16	0
5	MN	A	1299	1/1	0.98	0.10	20,20,20,20	0
8	MAN	A	1309	11/12	0.98	0.08	15,17,18,19	0
11	BR	A	1321	1/1	0.98	0.04	28,28,28,28	1
11	BR	A	1322	1/1	0.98	0.08	34,34,34,34	1
11	BR	A	1323	1/1	0.99	0.07	25,25,25,25	1

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.