



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 19, 2023 – 11:13 PM EDT

PDB ID : 2FXK
Title : Crystal structure of the macro-domain of human core histone variant macroH2A1.1 (form A)
Authors : Kustatscher, G.; Hothorn, M.; Pugieux, C.; Scheffzek, K.; Ladurner, A.G.
Deposited on : 2006-02-06
Resolution : 2.54 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	211	
1	B	211	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H2A histone family, member Y isoform 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	187	1405	901	230	269	5	0	0	0
1	B	187	1405	901	230	269	5	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	159	GLY	-	cloning artifact	GB 20336746
A	160	ALA	-	cloning artifact	GB 20336746
A	161	MET	-	cloning artifact	GB 20336746
B	159	GLY	-	cloning artifact	GB 20336746
B	160	ALA	-	cloning artifact	GB 20336746
B	161	MET	-	cloning artifact	GB 20336746

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	21	Total	O	0	0
			21	21		
2	B	22	Total	O	0	0
			22	22		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H2A histone family, member Y isoform 1

Chain A: 



- Molecule 1: H2A histone family, member Y isoform 1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	90.77Å 98.27Å 42.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	90.91 – 2.54 98.27 – 2.54	Depositor EDS
% Data completeness (in resolution range)	99.0 (90.91-2.54) 99.0 (98.27-2.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 2.55Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.207 , 0.270 0.290 , (Not available)	Depositor DCC
R_{free} test set	37 reflections (0.29%)	wwPDB-VP
Wilson B-factor (Å ²)	61.5	Xtrriage
Anisotropy	0.333	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2853	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/1429	0.79	0/1929
1	B	0.61	0/1429	0.71	1/1929 (0.1%)
All	All	0.67	0/2858	0.75	1/3858 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	352	ASP	CB-CG-OD1	-5.06	113.75	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1405	0	1435	25	0
1	B	1405	0	1435	21	0
2	A	21	0	0	0	0
2	B	22	0	0	6	0
All	All	2853	0	2870	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:THR:HG22	1:B:221:TYR:O	1.99	0.63
1:A:325:GLN:CG	1:A:362:GLU:OE1	2.48	0.62
1:B:368:ALA:HB1	2:B:42:HOH:O	1.99	0.61
1:A:183:PHE:C	1:A:183:PHE:CD2	2.75	0.60
1:A:210:ASP:OD2	1:A:304:LYS:HG2	2.02	0.59
1:A:183:PHE:C	1:A:183:PHE:HD2	2.05	0.59
1:B:214:HIS:CE1	1:B:216:THR:HG23	2.38	0.59
1:A:214:HIS:CE1	1:A:216:THR:HG23	2.37	0.58
1:A:216:THR:CG2	1:A:221:TYR:O	2.51	0.58
1:A:216:THR:HG22	1:A:221:TYR:O	2.02	0.58
1:B:202:ALA:HB3	1:B:207:ILE:HD11	1.84	0.58
1:B:314:GLY:HA3	2:B:43:HOH:O	2.04	0.57
1:A:202:ALA:CB	1:A:207:ILE:HD11	2.35	0.57
1:A:321:GLN:OE1	1:A:325:GLN:NE2	2.37	0.56
1:B:202:ALA:CB	1:B:207:ILE:HD11	2.36	0.56
1:A:325:GLN:HG3	1:A:362:GLU:OE1	2.06	0.56
1:B:351:PHE:CE2	2:B:3:HOH:O	2.59	0.55
1:A:223:GLY:O	1:A:227:GLY:HA3	2.07	0.55
1:A:206:SER:HB3	1:A:233:LYS:HZ1	1.73	0.54
1:A:216:THR:HG21	1:A:220:PHE:HA	1.89	0.53
1:A:352:ASP:OD1	1:A:355:SER:OG	2.22	0.52
1:B:197:LEU:HD12	1:B:346:VAL:HG13	1.91	0.52
1:B:216:THR:CG2	1:B:221:TYR:O	2.59	0.50
1:A:202:ALA:HB3	1:A:207:ILE:HD11	1.94	0.49
1:B:182:GLY:N	2:B:16:HOH:O	2.44	0.49
1:A:254:VAL:O	1:A:255:ALA:HB3	2.13	0.48
1:B:328:LEU:HG	1:B:366:LEU:HD11	1.97	0.47
1:A:218:THR:HG22	1:A:274:ASN:ND2	2.31	0.46
1:B:216:THR:HG21	1:B:220:PHE:HA	1.98	0.46
1:A:245:LEU:HD22	1:A:260:SER:HB3	1.97	0.45
1:A:257:ALA:HA	1:A:272:HIS:O	2.16	0.45
1:B:316:ASN:N	1:B:316:ASN:HD22	2.13	0.45
1:A:281:ASP:O	1:A:282:LYS:C	2.54	0.44
1:B:255:ALA:O	1:B:293:ASN:ND2	2.46	0.44
1:A:242:VAL:HG22	1:A:272:HIS:CE1	2.52	0.43
1:B:252:LEU:HD23	1:B:274:ASN:HB2	2.01	0.43
1:B:201:GLN:O	1:B:202:ALA:HB2	2.19	0.43
1:A:201:GLN:HG2	1:A:350:LEU:O	2.18	0.42
1:A:206:SER:HB3	1:A:233:LYS:NZ	2.35	0.42
1:B:245:LEU:HD22	1:B:260:SER:HB3	2.02	0.42
1:A:223:GLY:O	1:A:227:GLY:CA	2.68	0.41
1:A:192:PHE:CZ	1:A:367:ASP:HB3	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:351:PHE:CZ	2:B:3:HOH:O	2.57	0.41
1:B:316:ASN:ND2	2:B:23:HOH:O	2.53	0.41
1:B:200:VAL:HG12	1:B:201:GLN:N	2.35	0.40
1:B:219:ASP:O	1:B:220:PHE:C	2.58	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/211 (88%)	175 (95%)	9 (5%)	1 (0%)	29	40
1	B	185/211 (88%)	172 (93%)	12 (6%)	1 (0%)	29	40
All	All	370/422 (88%)	347 (94%)	21 (6%)	2 (0%)	29	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	208	ASP
1	B	314	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/169 (90%)	136 (89%)	17 (11%)	6	6
1	B	153/169 (90%)	135 (88%)	18 (12%)	5	5
All	All	306/338 (90%)	271 (89%)	35 (11%)	5	5

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	183	PHE
1	A	201	GLN
1	A	216	THR
1	A	225	GLU
1	A	229	THR
1	A	240	GLU
1	A	243	LEU
1	A	253	GLU
1	A	275	SER
1	A	281	ASP
1	A	284	GLU
1	A	287	LEU
1	A	289	LYS
1	A	297	LEU
1	A	304	LYS
1	A	313	SER
1	A	344	LYS
1	B	187	SER
1	B	193	LEU
1	B	201	GLN
1	B	204	ILE
1	B	216	THR
1	B	219	ASP
1	B	231	GLU
1	B	253	GLU
1	B	278	TRP
1	B	287	LEU
1	B	304	LYS
1	B	315	ARG
1	B	316	ASN
1	B	321	GLN
1	B	328	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	336	VAL
1	B	337	SER
1	B	346	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	325	GLN
1	B	316	ASN
1	B	361	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.