



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 19, 2023 – 09:25 PM EDT

PDB ID : 2GL5
Title : Crystal Structure of Putative Dehydratase from Salmonella Thyphimurium
Authors : Patskovsky, Y.; Sauder, J.M.; Dickey, M.; Adams, J.M.; Ozyurt, S.; Wasserman, S.R.; Gerlt, J.; Almo, S.C.; Burley, S.K.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2006-04-04
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	410	
1	B	410	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative dehydratase protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3166	2013	534	603	16	0	7	0
1	B	401	3173	2020	535	602	16	0	8	0


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	cloning artifact	UNP Q8ZNH1
A	0	SER	-	cloning artifact	UNP Q8ZNH1
A	1	LEU	-	cloning artifact	UNP Q8ZNH1
A	143	ASN	ARG	engineered mutation	UNP Q8ZNH1
A	401	GLU	-	cloning artifact	UNP Q8ZNH1
A	402	GLY	-	cloning artifact	UNP Q8ZNH1
A	403	HIS	-	expression tag	UNP Q8ZNH1
A	404	HIS	-	expression tag	UNP Q8ZNH1
A	405	HIS	-	expression tag	UNP Q8ZNH1
A	406	HIS	-	expression tag	UNP Q8ZNH1
A	407	HIS	-	expression tag	UNP Q8ZNH1
A	408	HIS	-	expression tag	UNP Q8ZNH1
B	-1	MET	-	cloning artifact	UNP Q8ZNH1
B	0	SER	-	cloning artifact	UNP Q8ZNH1
B	1	LEU	-	cloning artifact	UNP Q8ZNH1
B	143	ASN	ARG	engineered mutation	UNP Q8ZNH1
B	401	GLU	-	cloning artifact	UNP Q8ZNH1
B	402	GLY	-	cloning artifact	UNP Q8ZNH1
B	403	HIS	-	expression tag	UNP Q8ZNH1
B	404	HIS	-	expression tag	UNP Q8ZNH1
B	405	HIS	-	expression tag	UNP Q8ZNH1
B	406	HIS	-	expression tag	UNP Q8ZNH1
B	407	HIS	-	expression tag	UNP Q8ZNH1
B	408	HIS	-	expression tag	UNP Q8ZNH1

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	308	Total O 308 308	0	0
4	B	316	Total O 316 316	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: putative dehydratase protein

Chain A:  93% 5%



- Molecule 1: putative dehydratase protein

Chain B:  91% 7%



4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	116.97Å 116.97Å 116.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.60 21.35 – 1.52	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.60) 99.6 (21.35-1.52)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.04 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.185 , 0.218 0.184 , 0.218	Depositor DCC
R_{free} test set	3482 reflections (2.92%)	wwPDB-VP
Wilson B-factor (Å ²)	9.9	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 30.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.075 for l,-k,h 0.076 for -l,-k,-h 0.070 for -h,-l,-k 0.075 for -h,l,k 0.256 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6971	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/3253	0.61	0/4407
1	B	0.46	0/3263	0.61	0/4418
All	All	0.46	0/6516	0.61	0/8825

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3166	0	3137	12	2
1	B	3173	0	3157	18	2
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	6	0	8	0	0
4	A	308	0	0	0	0
4	B	316	0	0	0	0
All	All	6971	0	6302	29	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:176:ARG:HG3	1:B:176:ARG:HH21	1.31	0.94
1:A:194:LEU:H	1:A:234:GLN:HE22	1.14	0.92
1:B:194:LEU:H	1:B:234:GLN:HE22	1.29	0.77
1:B:176:ARG:HG3	1:B:176:ARG:NH2	1.92	0.72
1:B:351:HIS:HD2	1:B:353[B]:ASN:H	1.38	0.71
1:B:351:HIS:HD2	1:B:353[A]:ASN:H	1.39	0.71
1:B:176:ARG:HH21	1:B:176:ARG:CG	2.05	0.69
1:B:127[B]:LYS:HG2	1:B:373:TYR:HB3	1.82	0.62
1:A:90:GLY:HA3	1:B:90:GLY:HA3	1.85	0.58
1:B:351:HIS:CD2	1:B:353[B]:ASN:H	2.22	0.57
1:B:351:HIS:CD2	1:B:353[A]:ASN:H	2.22	0.57
1:B:272:THR:HG22	1:B:291:ILE:HG21	1.91	0.53
1:A:194:LEU:N	1:A:234:GLN:HE22	1.97	0.51
1:A:351:HIS:CD2	1:A:353[B]:ASN:H	2.29	0.51
1:A:351:HIS:CD2	1:A:353[A]:ASN:H	2.30	0.50
1:A:351:HIS:HD2	1:A:353[B]:ASN:H	1.60	0.49
1:A:194:LEU:H	1:A:234:GLN:NE2	1.95	0.49
1:A:351:HIS:HD2	1:A:353[A]:ASN:H	1.62	0.48
1:B:1:LEU:HD22	1:B:34:SER:HB3	1.95	0.48
1:B:12:GLU:HG2	1:B:24:ASN:HD21	1.79	0.47
1:A:164:TYR:HH	1:A:350[A]:HIS:CD2	2.31	0.47
1:A:95:ALA:HB2	1:A:300:LEU:HD23	1.97	0.46
1:B:388:ASP:HA	1:B:391:VAL:HG22	2.00	0.44
1:B:295:GLN:HA	1:B:322:GLN:O	2.19	0.43
1:B:194:LEU:H	1:B:234:GLN:NE2	2.07	0.43
1:B:39:ILE:HG21	1:B:386:LEU:HG	2.03	0.41
1:B:170:ASP:HA	1:B:222:GLU:O	2.20	0.41
1:A:133:SER:HB3	1:A:351:HIS:HA	2.03	0.41
1:A:272:THR:HG22	1:A:291:ILE:HG21	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:317[A]:TYR:OH	1:B:287:GLU:OE2[3_555]	2.15	0.05
1:A:287:GLU:OE2	1:B:317[A]:TYR:OH[3_555]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	406/410 (99%)	395 (97%)	10 (2%)	1 (0%)	47	26
1	B	407/410 (99%)	393 (97%)	12 (3%)	2 (0%)	29	11
All	All	813/820 (99%)	788 (97%)	22 (3%)	3 (0%)	34	15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	49	GLY
1	A	49	GLY
1	B	134	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/338 (100%)	331 (98%)	6 (2%)	59	36
1	B	338/338 (100%)	333 (98%)	5 (2%)	65	44
All	All	675/676 (100%)	664 (98%)	11 (2%)	65	41

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	19	THR
1	A	176	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	241[A]	LYS
1	A	241[B]	LYS
1	A	374	TYR
1	B	22	SER
1	B	23	TYR
1	B	176	ARG
1	B	322	GLN
1	B	374	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	24	ASN
1	A	118	GLN
1	A	234	GLN
1	A	289	GLN
1	A	351	HIS
1	A	365	HIS
1	B	24	ASN
1	B	234	GLN
1	B	351	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	701	-	5,5,5	0.38	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	701	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	701	GOL	O1-C1-C2-O2
3	B	701	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.