



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2023 – 10:21 AM EDT

PDB ID : 2IBG
Title : Crystal Structure of Hedgehog Bound to the FNIII Domains of Ihog
Authors : McLellan, J.S.; Leahy, D.J.
Deposited on : 2006-09-11
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 86% 13%
1	B	214	 83% 15%
1	C	214	 87% 12%
1	D	214	 90% 9%
2	E	150	 76% 17% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	150	
2	G	150	
2	H	150	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CG9211-PA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	214	Total 1715	C 1095	N 290	O 325	S 5	0	0	0
1	B	213	Total 1705	C 1089	N 288	O 323	S 5	0	0	0
1	C	213	Total 1705	C 1089	N 288	O 323	S 5	0	0	0
1	D	213	Total 1705	C 1089	N 288	O 323	S 5	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	464	SER	-	cloning artifact	UNP Q9VM64
A	465	THR	-	cloning artifact	UNP Q9VM64
B	464	SER	-	cloning artifact	UNP Q9VM64
B	465	THR	-	cloning artifact	UNP Q9VM64
C	464	SER	-	cloning artifact	UNP Q9VM64
C	465	THR	-	cloning artifact	UNP Q9VM64
D	464	SER	-	cloning artifact	UNP Q9VM64
D	465	THR	-	cloning artifact	UNP Q9VM64

- Molecule 2 is a protein called Protein hedgehog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	142	Total 1161	C 732	N 206	O 218	S 5	0	0	0
2	F	132	Total 1070	C 679	N 190	O 196	S 5	0	0	0
2	G	134	Total 1092	C 694	N 191	O 202	S 5	0	0	0
2	H	140	Total 1142	C 722	N 201	O 214	S 5	0	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0
3	E	1	Total O P 5 4 1	0	0
3	E	1	Total O P 5 4 1	0	0
3	E	1	Total O P 5 4 1	0	0
3	F	1	Total O P 5 4 1	0	0
3	H	1	Total O P 5 4 1	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	66	Total O 66 66	0	0
4	B	48	Total O 48 48	0	0
4	C	63	Total O 63 63	0	0
4	D	54	Total O 54 54	0	0
4	E	28	Total O 28 28	0	0
4	F	8	Total O 8 8	0	0
4	G	12	Total O 12 12	0	0
4	H	10	Total O 10 10	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CG9211-PA

Chain A: 




- Molecule 1: CG9211-PA

Chain B: 



- Molecule 1: CG9211-PA

Chain C: 





- Molecule 1: CG9211-PA

Chain D: 



- Molecule 2: Protein hedgehog

Chain E: 



Y235
D248

• Molecule 2: Protein hedgehog



L99 L102 V103 L104 L180 T107 I108 L111 Y114 Y115 M16 M17 S117 A118 S119 G120 P121 L122 E123 G124 V125 V126 E202 R127 R128 D129 P131 K132 F133 K134 D135 L136 V137 P138 M139 Y140 M141 R147 ASP GLU GLU GLY GLY GLY ALA D155 R156 L157 M158 K160 R161 C162 K163 E164 K165

V168 E176 G179 R181 T185 E186 S187 TRP ASP GLU ASP TYR HIS HIS GLY GLN E197 S198 Y201 E202 G203 R204 A205 V206 T207 Y218 G219 M220 L224 A228 D231 W232 Y235 R239 H240 V245 K246 S247 ASP


• Molecule 2: Protein hedgehog



L99 V103 K105 I108 P109 S112 A118 S119 Y114 L122 V125 ILE ARG ASP SER PRO LYS F133 K134 D135 M139 Y140 M141 R142 D143 R147 ASP GLU GLU GLY THR GLY ALA ASP R156 L157 K165 M174 R181 V184 W188 D191 Y192

H193 E197 H200 Y201 E202 S203 R204 V206 Y218 L221 A222 V226 F227 A228 D231 S234 Y235 V236 I241 K246 S247 ASP

• Molecule 2: Protein hedgehog



LEU Y100 L104 K105 T108 P109 Y114 T115 N116 S117 A118 L122 V125 I126 D129 K132 F133 K134 D135 M139 R142 F145 ARG ASP GLU GLU GLY THR GLY ALA ASP A170 M174 I180 T185 W188 Y192 H200 Y201 V206 T207

I208 R213 D214 Q215 Y218 D231 Y235 R238 R239 H240 I241 K246 S247 D248

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.35Å 70.00Å 155.69Å 90.00° 90.18° 90.00°	Depositor
Resolution (Å)	37.68 – 2.20 37.68 – 2.20	Depositor EDS
% Data completeness (in resolution range)	92.2 (37.68-2.20) 92.1 (37.68-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.20Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.198 , 0.246 0.200 , 0.247	Depositor DCC
R_{free} test set	4169 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtrriage
Anisotropy	0.412	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.074 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11644	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/1764	0.73	0/2397
1	B	0.64	0/1754	0.68	0/2386
1	C	0.60	0/1754	0.67	0/2386
1	D	0.64	0/1754	0.71	0/2386
2	E	0.51	0/1187	0.66	1/1603 (0.1%)
2	F	1.98	31/1090 (2.8%)	1.07	7/1470 (0.5%)
2	G	0.52	0/1116	0.60	0/1508
2	H	0.48	0/1168	0.61	0/1578
All	All	0.83	31/11587 (0.3%)	0.72	8/15714 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	F	0	1

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	117	SER	CB-OG	23.21	1.72	1.42
2	F	202	GLU	CD-OE2	20.15	1.47	1.25
2	F	161	ARG	CZ-NH1	18.89	1.57	1.33
2	F	123	GLU	CD-OE1	17.42	1.44	1.25
2	F	121	PRO	C-N	15.07	1.68	1.34

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	161	ARG	NE-CZ-NH2	-19.50	110.55	120.30
2	F	121	PRO	O-C-N	8.21	135.84	122.70
2	F	121	PRO	CA-C-N	-6.54	102.82	117.20
2	F	161	ARG	NE-CZ-NH1	5.83	123.21	120.30
2	F	161	ARG	NH1-CZ-NH2	5.30	125.23	119.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	F	202	GLU	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1715	0	1674	25	0
1	B	1705	0	1661	21	0
1	C	1705	0	1661	15	0
1	D	1705	0	1661	18	0
2	E	1161	0	1137	16	0
2	F	1070	0	1080	29	0
2	G	1092	0	1073	22	0
2	H	1142	0	1120	18	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	C	10	0	0	0	0
3	D	5	0	0	0	0
3	E	15	0	0	0	0
3	F	5	0	0	1	0
3	H	5	0	0	0	0
4	A	66	0	0	2	0
4	B	48	0	0	0	0
4	C	63	0	0	1	0
4	D	54	0	0	1	0
4	E	28	0	0	0	0
4	F	8	0	0	0	0
4	G	12	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	10	0	0	0	0
All	All	11644	0	11067	154	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:121:PRO:C	2:F:122:LEU:N	1.68	1.46
2:F:117:SER:CB	2:F:117:SER:OG	1.72	1.34
2:H:185:THR:OG1	2:H:207:THR:HG22	1.68	0.93
1:C:580:PRO:HG2	1:C:610:ILE:HD11	1.65	0.78
2:F:168:VAL:HG23	2:F:228:ALA:HB1	1.67	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/214 (99%)	208 (98%)	4 (2%)	0	100	100
1	B	211/214 (99%)	207 (98%)	2 (1%)	2 (1%)	17	16
1	C	211/214 (99%)	206 (98%)	5 (2%)	0	100	100
1	D	211/214 (99%)	206 (98%)	5 (2%)	0	100	100
2	E	138/150 (92%)	136 (99%)	2 (1%)	0	100	100
2	F	126/150 (84%)	118 (94%)	7 (6%)	1 (1%)	19	19
2	G	128/150 (85%)	122 (95%)	6 (5%)	0	100	100
2	H	136/150 (91%)	128 (94%)	7 (5%)	1 (1%)	22	22
All	All	1373/1456 (94%)	1331 (97%)	38 (3%)	4 (0%)	41	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	509	LYS
1	B	508	ARG
2	F	198	SER
2	H	118	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	186/186 (100%)	177 (95%)	9 (5%)	25 32
1	B	185/186 (100%)	182 (98%)	3 (2%)	62 76
1	C	185/186 (100%)	182 (98%)	3 (2%)	62 76
1	D	185/186 (100%)	181 (98%)	4 (2%)	52 65
2	E	128/133 (96%)	119 (93%)	9 (7%)	15 16
2	F	119/133 (90%)	110 (92%)	9 (8%)	13 14
2	G	120/133 (90%)	112 (93%)	8 (7%)	16 18
2	H	126/133 (95%)	118 (94%)	8 (6%)	18 20
All	All	1234/1276 (97%)	1181 (96%)	53 (4%)	29 36

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	99	LEU
2	F	181	ARG
2	H	135	ASP
2	F	107	THR
2	F	139	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	598	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	670	GLN
2	H	139	ASN
1	D	517	ASN
2	E	139	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	E	408	-	4,4,4	4.14	1 (25%)	6,6,6	1.26	0
3	PO4	C	402	-	4,4,4	3.76	1 (25%)	6,6,6	1.50	2 (33%)
3	PO4	B	401	-	4,4,4	3.60	1 (25%)	6,6,6	1.27	0
3	PO4	D	405	-	4,4,4	4.18	1 (25%)	6,6,6	1.19	0
3	PO4	E	412	-	4,4,4	4.16	1 (25%)	6,6,6	1.47	1 (16%)
3	PO4	B	406	-	4,4,4	4.16	1 (25%)	6,6,6	1.28	0
3	PO4	A	404	-	4,4,4	4.17	1 (25%)	6,6,6	1.24	0
3	PO4	C	407	-	4,4,4	4.12	1 (25%)	6,6,6	1.21	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	F	410	-	4,4,4	4.20	1 (25%)	6,6,6	1.20	0
3	PO4	A	403	-	4,4,4	3.97	1 (25%)	6,6,6	1.81	2 (33%)
3	PO4	E	411	-	4,4,4	4.14	1 (25%)	6,6,6	1.19	1 (16%)
3	PO4	H	409	-	4,4,4	4.15	1 (25%)	6,6,6	1.30	0

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	410	PO4	P-O4	8.28	1.79	1.54
3	D	405	PO4	P-O4	8.23	1.79	1.54
3	E	412	PO4	P-O4	8.23	1.79	1.54
3	H	409	PO4	P-O4	8.21	1.79	1.54
3	B	406	PO4	P-O4	8.21	1.79	1.54

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	412	PO4	O3-P-O2	2.58	116.26	107.97
3	A	403	PO4	O4-P-O1	-2.54	101.60	110.89
3	A	403	PO4	O4-P-O2	-2.18	100.98	107.97
3	E	411	PO4	O4-P-O1	-2.13	103.11	110.89
3	C	402	PO4	O4-P-O1	-2.12	103.15	110.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	410	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	F	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	F	121:PRO	C	122:LEU	N	1.68

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/214 (100%)	-0.34	1 (0%) 91 90	21, 30, 43, 54	0
1	B	213/214 (99%)	-0.38	3 (1%) 75 73	24, 32, 45, 61	0
1	C	213/214 (99%)	-0.32	3 (1%) 75 73	24, 32, 45, 58	0
1	D	213/214 (99%)	-0.35	1 (0%) 91 90	24, 33, 44, 58	0
2	E	142/150 (94%)	-0.06	2 (1%) 75 73	35, 43, 52, 62	0
2	F	132/150 (88%)	0.96	18 (13%) 3 2	39, 51, 60, 65	0
2	G	134/150 (89%)	0.36	5 (3%) 41 39	39, 49, 55, 58	0
2	H	140/150 (93%)	0.43	6 (4%) 35 33	44, 54, 62, 64	0
All	All	1401/1456 (96%)	-0.05	39 (2%) 53 51	21, 38, 56, 65	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	127	ARG	7.9
2	F	126	ILE	4.8
2	F	129	ASP	4.7
1	B	507	LYS	4.3
2	F	203	GLY	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	F	410	5/5	0.90	0.14	91,91,91,92	0
3	PO4	H	409	5/5	0.93	0.15	76,77,78,78	0
3	PO4	A	404	5/5	0.94	0.08	68,69,69,69	0
3	PO4	D	405	5/5	0.96	0.13	79,79,79,80	0
3	PO4	E	411	5/5	0.96	0.16	70,70,71,72	0
3	PO4	E	408	5/5	0.97	0.23	78,78,79,79	0
3	PO4	E	412	5/5	0.98	0.15	57,58,60,60	0
3	PO4	B	406	5/5	0.98	0.07	75,76,76,76	0
3	PO4	A	403	5/5	0.98	0.10	35,35,37,37	0
3	PO4	B	401	5/5	0.99	0.06	24,26,29,29	0
3	PO4	C	402	5/5	0.99	0.06	29,30,31,32	0
3	PO4	C	407	5/5	0.99	0.11	79,79,79,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.