



# wwPDB NMR Structure Validation Summary Report ⓘ

Jun 12, 2024 – 09:42 AM EDT

PDB ID : 2IFS  
Title : Structure of the N-WASP EVH1 domain in complex with an extended WIP peptide  
Authors : Volkman, B.F.; Peterson, F.C.; Deng, Q.  
Deposited on : 2006-09-21

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	169	

## 2 Ensemble composition and analysis i

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:455-A:478, A:534-A:637 (128)	1.18	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 6, 9, 12, 13, 16
2	7, 14, 17
3	4, 10, 20
4	2, 15
5	8, 19
Single-model clusters	3; 5; 11; 18

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2706 atoms, of which 1346 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Wiskott-Aldrich Syndrome Protein interacting protein and Neural Wiskott-Aldrich syndrome protein chimera.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	169	2706	860	1346	245	248	7	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	449	GLY	-	cloning artifact	UNP O08816
A	450	SER	-	cloning artifact	UNP O08816
A	516	GLY	-	SEE REMARK 999	UNP O08816
A	517	GLY	-	SEE REMARK 999	UNP O08816
A	518	LEU	-	SEE REMARK 999	UNP O08816
A	519	VAL	-	SEE REMARK 999	UNP O08816
A	520	PRO	-	SEE REMARK 999	UNP O08816
A	521	ARG	-	SEE REMARK 999	UNP O08816
A	522	GLY	-	SEE REMARK 999	UNP O08816
A	523	SER	-	SEE REMARK 999	UNP O08816
A	524	GLY	-	SEE REMARK 999	UNP O08816
A	525	GLY	-	SEE REMARK 999	UNP O08816

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Wiskott-Aldrich Syndrome Protein interacting protein and Neural Wiskott-Aldrich syndrome protein chimera



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

- Molecule 1: Wiskott-Aldrich Syndrome Protein interacting protein and Neural Wiskott-Aldrich syndrome protein chimera



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *AUTOMATED METHODS WERE USED FOR BACKBONE CHEMICAL SHIFT ASSIGNMENT AND ITERATIVE NOE REFINEMENT. FINAL STRUCTURES WERE OBTAINED BY MOLECULAR DYNAMICS IN EXPLICIT SOLVENT.*

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Xplor-NIH	refinement	2.9.3

No chemical shift data was provided.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1038	1017	1015	14±3
All	All	20760	20340	20300	280

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

5 of 131 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:563:CYS:HG	1:A:578:PHE:HZ	0.80	1.18	4	3
1:A:603:TYR:HA	1:A:618:PHE:CE2	0.72	2.19	15	1
1:A:597:TYR:CD1	1:A:628:ARG:HG3	0.67	2.24	14	12
1:A:577:ILE:HG12	1:A:587:TRP:CE3	0.67	2.25	16	11
1:A:582:ASP:OD2	1:A:584:LYS:HG2	0.63	1.93	15	4

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/169 (76%)	118±2 (92±2%)	7±2 (6±2%)	2±1 (2±1%)	12	54
All	All	2560/3380 (76%)	2365 (92%)	148 (6%)	47 (2%)	12	54

5 of 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	550	ARG	19
1	A	585	LEU	18
1	A	594	ASN	2
1	A	610	ASP	2
1	A	612	CYS	2

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/147 (78%)	109±2 (96±2%)	5±2 (4±2%)	34	82
All	All	2280/2940 (78%)	2186 (96%)	94 (4%)	34	82

5 of 40 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	588	GLU	11
1	A	589	GLN	9
1	A	459	ILE	7
1	A	566	LYS	7
1	A	556	LYS	4

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.



## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided