



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 10, 2023 – 11:37 pm GMT

PDB ID : 2IUO
Title : Site Directed Mutagenesis of Key Residues Involved in the Catalytic Mechanism of Cyanase
Authors : Guilloton, M.; Walsh, M.A.; Joachimiak, A.; Anderson, M.P.
Deposited on : 2006-06-06
Resolution : 1.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)



The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	156	83% 17%
1	B	156	83% 16% .
1	C	156	82% 16% .
1	D	156	86% 12% .
1	E	156	83% 16% .
1	F	156	85% 15%
1	G	156	85% 15% .
1	H	156	82% 17% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	I	156		84% 16%
1	J	156		83% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BR	B	1157	-	-	X	-
2	BR	D	1157	-	-	X	-
2	BR	E	1157	-	-	X	-
2	BR	F	1157	-	-	X	-
4	SO4	B	1160	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 14230 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYANATE HYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	156	1201	771	199	226	5	0	1	0
1	B	156	1201	771	199	226	5	0	1	0
1	C	156	1201	771	199	226	5	0	1	0
1	D	156	1206	776	199	226	5	0	2	0
1	E	156	1206	776	199	226	5	0	2	0
1	F	156	1211	781	199	226	5	0	3	0
1	G	156	1201	771	199	226	5	0	1	0
1	H	156	1206	776	199	226	5	0	2	0
1	I	156	1201	771	199	226	5	0	1	0
1	J	156	1206	776	199	226	5	0	2	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	GLY	SER	engineered mutation	UNP P00816
B	122	GLY	SER	engineered mutation	UNP P00816
C	122	GLY	SER	engineered mutation	UNP P00816
D	122	GLY	SER	engineered mutation	UNP P00816
E	122	GLY	SER	engineered mutation	UNP P00816
F	122	GLY	SER	engineered mutation	UNP P00816
G	122	GLY	SER	engineered mutation	UNP P00816
H	122	GLY	SER	engineered mutation	UNP P00816
I	122	GLY	SER	engineered mutation	UNP P00816

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	122	GLY	SER	engineered mutation	UNP P00816

- Molecule 2 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Br 1 1	0	0
2	B	1	Total Br 1 1	0	0
2	C	1	Total Br 1 1	0	0
2	D	1	Total Br 1 1	0	0
2	E	1	Total Br 1 1	0	0
2	F	1	Total Br 1 1	0	0
2	G	1	Total Br 1 1	0	0
2	H	1	Total Br 1 1	0	0
2	I	1	Total Br 1 1	0	0
2	J	1	Total Br 1 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	B	1	Total Cl 1 1	0	0
3	C	1	Total Cl 1 1	0	0
3	D	1	Total Cl 1 1	0	0
3	E	1	Total Cl 1 1	0	0
3	F	1	Total Cl 1 1	0	0
3	G	1	Total Cl 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	H	1	Total Cl 1 1	0	0
3	I	1	Total Cl 1 1	0	0
3	J	1	Total Cl 1 1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		
4	G	1	Total	O	S	0	0
			5	4	1		
4	G	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	I	1	Total	O	S	0	0
			5	4	1		
4	I	1	Total	O	S	0	0
			5	4	1		
4	J	1	Total	O	S	0	0
			5	4	1		
4	J	1	Total	O	S	0	0
			5	4	1		
4	J	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is AZIDE ION (three-letter code: AZI) (formula: N₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total N 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	201	Total O 201 201	0	0
6	B	224	Total O 224 224	0	0
6	C	205	Total O 205 205	0	0
6	D	208	Total O 208 208	0	0
6	E	217	Total O 217 217	0	0
6	F	215	Total O 215 215	0	0
6	G	187	Total O 187 187	0	0
6	H	192	Total O 192 192	0	0
6	I	186	Total O 186 186	0	0
6	J	217	Total O 217 217	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: CYANATE HYDRATASE

Chain A: 83% 17%




- Molecule 1: CYANATE HYDRATASE

Chain B: 83% 16%



- Molecule 1: CYANATE HYDRATASE

Chain C: 82% 16%



- Molecule 1: CYANATE HYDRATASE

Chain D: 86% 12%



- Molecule 1: CYANATE HYDRATASE

Chain E: 83% 16%



- Molecule 1: CYANATE HYDRATASE

Chain F: 85% 15%



- Molecule 1: CYANATE HYDRATASE

Chain G: 85% 15%



- Molecule 1: CYANATE HYDRATASE

Chain H: 82% 17%



- Molecule 1: CYANATE HYDRATASE

Chain I: 84% 16%



- Molecule 1: CYANATE HYDRATASE

Chain J: 83% 14%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	76.60Å 80.75Å 82.28Å 70.02° 71.81° 66.34°	Depositor
Resolution (Å)	76.70 – 1.90	Depositor
% Data completeness (in resolution range)	94.8 (76.70-1.90)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.149 , 0.208	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	14230	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CL, BR, SO4, AZI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	1/1221 (0.1%)	0.92	1/1650 (0.1%)
1	B	0.99	1/1221 (0.1%)	0.91	1/1650 (0.1%)
1	C	0.98	0/1221	0.89	1/1650 (0.1%)
1	D	1.00	1/1229 (0.1%)	1.00	4/1661 (0.2%)
1	E	0.98	0/1229	0.92	0/1661
1	F	0.99	1/1237 (0.1%)	0.92	2/1673 (0.1%)
1	G	0.90	0/1221	0.89	1/1650 (0.1%)
1	H	0.92	0/1229	0.92	2/1661 (0.1%)
1	I	0.96	0/1221	0.90	2/1649 (0.1%)
1	J	0.92	0/1229	0.90	1/1662 (0.1%)
All	All	0.96	4/12258 (0.0%)	0.92	15/16567 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	110	ALA	CA-CB	6.65	1.66	1.52
1	D	40	GLU	CB-CG	-6.09	1.40	1.52
1	B	69	GLU	CB-CG	-5.57	1.41	1.52
1	A	46	ALA	CA-CB	5.41	1.63	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	96	ARG	NE-CZ-NH1	7.37	123.98	120.30
1	I	96	ARG	NE-CZ-NH2	-6.34	117.13	120.30
1	D	81	ARG	NE-CZ-NH2	-6.17	117.22	120.30
1	J	1	MET	CG-SD-CE	6.15	110.04	100.20
1	F	96	ARG	NE-CZ-NH2	-5.91	117.35	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1201	0	1252	43	0
1	B	1201	0	1252	39	0
1	C	1201	0	1252	31	0
1	D	1206	0	1263	29	0
1	E	1206	0	1263	30	0
1	F	1211	0	1274	33	0
1	G	1201	0	1252	35	0
1	H	1206	0	1263	40	0
1	I	1201	0	1252	37	0
1	J	1206	0	1263	37	0
2	A	1	0	0	1	0
2	B	1	0	0	2	0
2	C	1	0	0	1	0
2	D	1	0	0	2	0
2	E	1	0	0	2	0
2	F	1	0	0	2	0
2	G	1	0	0	1	0
2	H	1	0	0	1	0
2	I	1	0	0	1	0
2	J	1	0	0	1	0
3	A	1	0	0	1	0
3	B	1	0	0	1	0
3	C	1	0	0	1	0
3	D	1	0	0	1	0
3	E	1	0	0	1	0
3	F	1	0	0	1	0
3	G	1	0	0	1	0
3	H	1	0	0	1	0
3	I	1	0	0	1	0
3	J	1	0	0	1	0
4	A	10	0	0	0	0
4	B	10	0	0	2	0
4	C	10	0	0	0	0
4	D	15	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	15	0	0	0	0
4	F	10	0	0	0	0
4	G	10	0	0	0	0
4	H	10	0	0	0	0
4	I	10	0	0	0	0
4	J	15	0	0	1	0
5	B	3	0	0	0	0
6	A	201	0	0	17	0
6	B	224	0	0	16	0
6	C	205	0	0	10	0
6	D	208	0	0	6	0
6	E	217	0	0	11	0
6	F	215	0	0	9	0
6	G	187	0	0	8	0
6	H	192	0	0	12	0
6	I	186	0	0	16	0
6	J	217	0	0	12	0
All	All	14230	0	12586	321	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 321 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:CE	1:B:2:ILE:HG22	1.49	1.39
1:G:1:MET:CE	1:G:2:ILE:HG22	1.57	1.33
1:J:25:LYS:HE2	6:J:2075:HOH:O	1.29	1.32
1:D:1:MET:CE	1:D:2:ILE:HG22	1.60	1.31
1:H:1:MET:CE	1:H:2:ILE:HG22	1.64	1.27

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/156 (99%)	152 (98%)	3 (2%)	0	100	100
1	B	155/156 (99%)	151 (97%)	4 (3%)	0	100	100
1	C	155/156 (99%)	152 (98%)	3 (2%)	0	100	100
1	D	156/156 (100%)	153 (98%)	3 (2%)	0	100	100
1	E	156/156 (100%)	152 (97%)	4 (3%)	0	100	100
1	F	157/156 (101%)	154 (98%)	3 (2%)	0	100	100
1	G	155/156 (99%)	152 (98%)	3 (2%)	0	100	100
1	H	156/156 (100%)	153 (98%)	3 (2%)	0	100	100
1	I	155/156 (99%)	152 (98%)	3 (2%)	0	100	100
1	J	156/156 (100%)	152 (97%)	4 (3%)	0	100	100
All	All	1556/1560 (100%)	1523 (98%)	33 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/126 (101%)	127 (100%)	0	100	100
1	B	127/126 (101%)	124 (98%)	3 (2%)	49	43
1	C	127/126 (101%)	122 (96%)	5 (4%)	32	23
1	D	128/126 (102%)	126 (98%)	2 (2%)	62	60
1	E	128/126 (102%)	124 (97%)	4 (3%)	40	32
1	F	129/126 (102%)	127 (98%)	2 (2%)	62	60
1	G	127/126 (101%)	122 (96%)	5 (4%)	32	23
1	H	128/126 (102%)	125 (98%)	3 (2%)	50	45
1	I	127/126 (101%)	126 (99%)	1 (1%)	81	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	128/126 (102%)	124 (97%)	4 (3%)	40	32
All	All	1276/1260 (101%)	1247 (98%)	29 (2%)	50	45

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	27	LEU
1	J	102	GLN
1	G	36	THR
1	I	127	LYS
1	G	27	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	76	GLN
1	H	102	GLN
1	H	51	GLN
1	I	51	GLN
1	C	102	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 44 ligands modelled in this entry, 20 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	G	1159	-	4,4,4	0.18	0	6,6,6	0.56	0
5	AZI	B	1161	-	0,2,2	-	-	0,1,1	-	-
4	SO4	H	1159	-	4,4,4	0.18	0	6,6,6	0.50	0
4	SO4	G	1160	-	4,4,4	0.12	0	6,6,6	0.39	0
4	SO4	E	1160	-	4,4,4	0.81	0	6,6,6	2.01	1 (16%)
4	SO4	F	1160	-	4,4,4	0.18	0	6,6,6	0.40	0
4	SO4	J	1161	-	4,4,4	0.22	0	6,6,6	0.78	0
4	SO4	E	1159	-	4,4,4	0.21	0	6,6,6	0.56	0
4	SO4	C	1160	-	4,4,4	0.64	0	6,6,6	2.39	2 (33%)
4	SO4	B	1159	-	4,4,4	0.09	0	6,6,6	0.24	0
4	SO4	I	1160	-	4,4,4	0.17	0	6,6,6	0.54	0
4	SO4	J	1160	-	4,4,4	0.71	0	6,6,6	1.87	3 (50%)
4	SO4	B	1160	-	4,4,4	0.70	0	6,6,6	2.06	2 (33%)
4	SO4	E	1161	-	4,4,4	0.21	0	6,6,6	0.45	0
4	SO4	A	1159	-	4,4,4	0.09	0	6,6,6	0.65	0
4	SO4	A	1160	-	4,4,4	0.17	0	6,6,6	0.27	0
4	SO4	H	1160	-	4,4,4	0.20	0	6,6,6	0.58	0
4	SO4	D	1161	-	4,4,4	0.15	0	6,6,6	0.31	0
4	SO4	I	1159	-	4,4,4	0.10	0	6,6,6	0.51	0
4	SO4	F	1159	-	4,4,4	0.19	0	6,6,6	0.64	0
4	SO4	J	1159	-	4,4,4	0.19	0	6,6,6	0.36	0
4	SO4	D	1159	-	4,4,4	0.24	0	6,6,6	0.72	0
4	SO4	D	1160	-	4,4,4	0.77	0	6,6,6	0.92	0
4	SO4	C	1159	-	4,4,4	0.22	0	6,6,6	0.45	0

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1160	SO4	O3-S-O2	-5.19	82.23	109.31
4	E	1160	SO4	O4-S-O2	-4.33	86.73	109.31
4	B	1160	SO4	O4-S-O2	-3.51	91.00	109.31
4	J	1160	SO4	O2-S-O1	-2.38	91.87	109.43
4	C	1160	SO4	O4-S-O1	2.37	121.70	109.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	J	1160	SO4	1	0
4	B	1160	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.