



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 06:09 am GMT

PDB ID : 2IY6
Title : 1-PYRROLINE-5-CARBOXYLATE DEHYDROGENASE FROM THERMUS WITH BOUND CITRATE
Authors : Inagaki, E.; Sakamoto, K.; Nishio, M.; Yokoyama, S.
Deposited on : 2006-07-13
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

PERCENTILES INFOmissingINFO

1 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 9428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-PYRROLINE-5-CARBOXYLATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	516	Total	C	N	O	S	0	8	0
			4064	2614	698	745	7			
1	B	516	Total	C	N	O	S	0	10	0
			4073	2618	700	748	7			

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0

- Molecule 4 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 8 6 2	0	0
4	A	1	Total C O 8 6 2	0	0
4	B	1	Total C O 8 6 2	0	0
4	B	1	Total C O 8 6 2	0	0
4	B	1	Total C O 8 6 2	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0
5	B	1	Total Na 1 1	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0
6	B	1	Total Cl 1 1	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	590	Total 590	O 590	0	0
7	B	567	Total 567	O 567	0	0

2 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1-PYRROLINE-5-CARBOXYLATE DEHYDROGENASE

Chain A: 



- Molecule 1: 1-PYRROLINE-5-CARBOXYLATE DEHYDROGENASE

Chain B: 



3 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	102.05Å 102.05Å 278.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.80 28.08 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-1.80) 99.1 (28.08-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	12.44 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.139 , 0.170 0.139 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	9.9	Xtrriage
Anisotropy	0.327	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.087 for -h-k,k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9428	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality

4.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, FLC, MRD, NA, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/4210	0.60	0/5710
1	B	0.53	0/4227	0.59	0/5733
All	All	0.54	0/8437	0.60	0/11443

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4064	0	4040	18	0
1	B	4073	0	4051	18	0
2	A	13	0	5	0	0
2	B	13	0	5	1	0
3	A	24	0	42	1	0
3	B	40	0	70	1	0
4	A	16	0	28	2	0
4	B	24	0	42	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	1	0	0	0	0
7	A	590	0	0	3	0
7	B	567	0	0	5	0
All	All	9428	0	8283	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286[B]:TYR:HD2	7:A:2554:HOH:O	1.15	1.27
1:B:286[B]:TYR:HD2	7:B:2527:HOH:O	1.15	1.23
1:A:61:PRO:HD2	1:A:395:GLU:HG2	1.36	1.04
1:B:286[B]:TYR:CD2	7:B:2527:HOH:O	1.98	0.91
1:A:61:PRO:HD2	1:A:395:GLU:CG	2.05	0.86
1:A:286[B]:TYR:CD2	7:A:2554:HOH:O	2.01	0.86
1:A:245:ALA:HB2	4:A:1533:MRD:H1C1	1.78	0.65
1:B:369:GLU:OE2	1:B:391:ARG:NE	2.34	0.60
1:A:286[B]:TYR:CZ	1:A:505:LEU:HB3	2.38	0.59
1:A:80:LEU:HA	1:A:205:ILE:HD11	1.87	0.56
1:B:286[B]:TYR:CZ	1:B:505:LEU:HB3	2.41	0.56
1:B:111[B]:ARG:NE	7:B:2187:HOH:O	2.42	0.53
1:B:80:LEU:HA	1:B:205:ILE:HD11	1.92	0.52
1:B:61:PRO:HD2	1:B:395:GLU:HG3	1.91	0.51
1:A:144:TYR:CE1	3:B:1530:MPD:H11	2.45	0.51
1:B:111[B]:ARG:NH2	7:B:2187:HOH:O	2.44	0.50
1:B:21:ARG:NH2	1:B:25:GLU:CD	2.66	0.49
1:B:185:PHE:CE1	2:B:1517:FLC:HG2	2.50	0.47
1:B:180[A]:ILE:HD13	4:B:1535:MRD:H5C2	1.97	0.47
1:B:309:GLY:HA3	1:B:450:TYR:OH	2.15	0.47
1:B:180[B]:ILE:HD13	1:B:207:LYS:HB3	1.97	0.46
1:A:367:GLU:HG3	1:A:371:LYS:HE3	1.97	0.45
1:A:126:LYS:HG2	1:A:317:PHE:CE1	2.52	0.45
1:A:200:VAL:HB	1:A:507:LEU:HD21	1.99	0.45
1:A:112:ARG:NH2	7:A:2210:HOH:O	2.43	0.44
1:B:21:ARG:NH2	1:B:25:GLU:OE1	2.48	0.44
1:A:180[A]:ILE:HD13	4:A:1533:MRD:H5C2	2.00	0.44
1:A:180[B]:ILE:HD13	1:A:207:LYS:HB3	1.99	0.43
1:B:1:MET:CE	1:B:103:LEU:HD11	2.49	0.43
1:A:286[B]:TYR:CE2	1:A:505:LEU:HD22	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:200:VAL:HB	1:B:507:LEU:HD21	2.00	0.42
1:A:138:ALA:HB2	1:A:188:ALA:HA	2.02	0.41
1:A:392:LEU:HD21	1:A:400:ALA:HB2	2.01	0.41
1:B:24:ARG:HD3	7:B:2052:HOH:O	2.19	0.41
1:B:140:ASP:HB3	1:B:480:VAL:HB	2.02	0.41
3:A:1531:MPD:HM1	3:A:1531:MPD:O4	2.21	0.41

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	522/516 (101%)	515 (99%)	7 (1%)	0	100	100
1	B	524/516 (102%)	515 (98%)	9 (2%)	0	100	100
All	All	1046/1032 (101%)	1030 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/407 (102%)	408 (98%)	7 (2%)	60	51
1	B	417/407 (102%)	410 (98%)	7 (2%)	60	51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	832/814 (102%)	818 (98%)	14 (2%)	65 51

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	LEU
1	A	286[A]	TYR
1	A	286[B]	TYR
1	A	288	GLU
1	A	340	GLU
1	A	419	PHE
1	A	464	PHE
1	B	29	ARG
1	B	101	LEU
1	B	286[A]	TYR
1	B	286[B]	TYR
1	B	288	GLU
1	B	419	PHE
1	B	464	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	381	ASN
1	A	384	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry

Of 19 ligands modelled in this entry, 4 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MPD	B	1531	-	7,7,7	0.36	0	9,10,10	0.47	0
3	MPD	B	1533	-	7,7,7	0.35	0	9,10,10	0.46	0
4	MRD	B	1534	-	7,7,7	0.32	0	9,10,10	0.62	0
4	MRD	B	1535	-	7,7,7	0.25	0	9,10,10	0.23	0
3	MPD	A	1531	-	7,7,7	0.34	0	9,10,10	1.03	1 (11%)
3	MPD	A	1530	-	7,7,7	0.38	0	9,10,10	0.67	0
3	MPD	A	1534	-	7,7,7	0.26	0	9,10,10	0.25	0
3	MPD	B	1532	-	7,7,7	0.26	0	9,10,10	0.23	0
4	MRD	A	1532	-	7,7,7	0.32	0	9,10,10	0.46	0
4	MRD	B	1537	-	7,7,7	0.38	0	9,10,10	0.50	0
3	MPD	B	1530	-	7,7,7	0.33	0	9,10,10	0.67	0
2	FLC	B	1517	-	12,12,12	1.11	0	17,17,17	1.30	2 (11%)
4	MRD	A	1533	-	7,7,7	0.28	0	9,10,10	0.27	0
2	FLC	A	1517	-	12,12,12	1.20	2 (16%)	17,17,17	1.38	2 (11%)
3	MPD	B	1536	-	7,7,7	0.24	0	9,10,10	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	B	1531	-	-	0/5/5/5	-
3	MPD	B	1533	-	-	0/5/5/5	-
4	MRD	B	1534	-	-	0/5/5/5	-
4	MRD	B	1535	-	-	0/5/5/5	-
3	MPD	A	1531	-	-	1/5/5/5	-
3	MPD	A	1530	-	-	1/5/5/5	-
3	MPD	A	1534	-	-	1/5/5/5	-
3	MPD	B	1532	-	-	1/5/5/5	-
4	MRD	A	1532	-	-	0/5/5/5	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MRD	B	1537	-	-	0/5/5/5	-
3	MPD	B	1530	-	-	1/5/5/5	-
2	FLC	B	1517	-	-	4/16/16/16	-
4	MRD	A	1533	-	-	0/5/5/5	-
2	FLC	A	1517	-	-	6/16/16/16	-
3	MPD	B	1536	-	-	3/5/5/5	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1517	FLC	OG2-CGC	-2.06	1.23	1.30
2	A	1517	FLC	CB-CBC	-2.01	1.51	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1517	FLC	OB2-CBC-CB	3.07	118.38	113.05
3	A	1531	MPD	CM-C2-C1	-2.88	104.58	110.57
2	B	1517	FLC	OB2-CBC-CB	2.82	117.94	113.05
2	B	1517	FLC	CB-CG-CGC	2.13	118.98	113.81
2	A	1517	FLC	OB1-CBC-CB	-2.12	119.25	122.25

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1530	MPD	C2-C3-C4-O4
3	B	1530	MPD	C2-C3-C4-O4
3	B	1532	MPD	O2-C2-C3-C4
3	B	1536	MPD	O2-C2-C3-C4
3	A	1534	MPD	C1-C2-C3-C4
3	B	1536	MPD	C1-C2-C3-C4
3	B	1536	MPD	CM-C2-C3-C4
2	A	1517	FLC	CA-CB-CBC-OB1
2	A	1517	FLC	CA-CB-CBC-OB2
2	A	1517	FLC	CG-CB-CBC-OB2
2	A	1517	FLC	CG-CB-CBC-OB1
2	B	1517	FLC	CA-CB-CBC-OB2
2	B	1517	FLC	CB-CG-CGC-OG2
2	B	1517	FLC	CA-CB-CBC-OB1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1517	FLC	CB-CG-CGC-OG1
2	B	1517	FLC	CB-CG-CGC-OG1
2	A	1517	FLC	CB-CG-CGC-OG2
3	A	1531	MPD	C2-C3-C4-C5

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1535	MRD	1	0
3	A	1531	MPD	1	0
3	B	1530	MPD	1	0
2	B	1517	FLC	1	0
4	A	1533	MRD	2	0

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	516/516 (100%)	-0.84	1 (0%) 95 93	3, 7, 17, 25	0
1	B	516/516 (100%)	-0.83	1 (0%) 95 93	4, 8, 19, 25	0
All	All	1032/1032 (100%)	-0.83	2 (0%) 95 93	3, 8, 18, 25	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	3.1
1	A	1	MET	3.0

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MRD	B	1537	8/8	0.75	0.23	27,30,30,31	0
3	MPD	A	1531	8/8	0.79	0.18	30,31,33,34	0
3	MPD	B	1533	8/8	0.84	0.15	30,32,33,34	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MPD	B	1532	8/8	0.87	0.13	26,27,29,29	0
3	MPD	B	1531	8/8	0.88	0.14	27,28,30,31	0
3	MPD	A	1534	8/8	0.88	0.13	35,36,37,37	0
3	MPD	B	1536	8/8	0.90	0.14	32,33,34,35	0
4	MRD	A	1533	8/8	0.93	0.11	22,23,24,25	0
4	MRD	B	1535	8/8	0.93	0.12	22,23,24,24	0
4	MRD	A	1532	8/8	0.93	0.09	20,21,22,22	0
3	MPD	A	1530	8/8	0.94	0.10	17,20,21,22	0
4	MRD	B	1534	8/8	0.94	0.13	17,19,20,21	0
3	MPD	B	1530	8/8	0.95	0.15	15,18,20,20	0
5	NA	A	1540	1/1	0.98	0.09	19,19,19,19	0
2	FLC	A	1517	13/13	0.99	0.06	6,9,11,11	0
2	FLC	B	1517	13/13	0.99	0.05	8,10,11,11	0
5	NA	B	1540	1/1	0.99	0.06	17,17,17,17	0
6	CL	A	1541	1/1	0.99	0.04	18,18,18,18	0
6	CL	B	1541	1/1	1.00	0.05	16,16,16,16	0

5.5 Other polymers [i](#)

There are no such residues in this entry.