



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 20, 2024 – 07:14 PM EDT

PDB ID : 2KQW
Title : SARS coronavirus-unique domain (SUD): Three-domain molecular architecture in solution and RNA binding. II: Structure of the SUD-C domain of SUD-MC
Authors : Johnson, M.A.; Chatterjee, A.; Wuthrich, K.; Joint Center for Structural Genomics (JCSG)
Deposited on : 2009-11-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	198	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:132-A:198 (67)	0.70	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	3, 6, 9, 12, 13, 16, 17, 20
2	4, 5, 10, 11, 14, 18, 19
3	1, 2, 7
4	8, 15

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1062 atoms, of which 523 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Non-structural protein 3.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	67	1062	343	523	88	108	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P0C6U8
A	2	SER	-	expression tag	UNP P0C6U8
A	3	HIS	-	expression tag	UNP P0C6U8
A	4	MET	-	expression tag	UNP P0C6U8

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Non-structural protein 3

Chain A:  30% • • 66%

GLY
SER
HIS
MET
GLY
THR
VAL
SER
TRP
ASN
LEU
ARG
GLU
MET
LEU
ALA
HIS
ALA
ALA
GLU
THR
LEU
ARG
LYS
MET
LEU
PRO
ILE
CYS
TYR
MET
ASP
VAL
ARG
ALA
ILE
MET
LEU
ALA
THR
ILE
GLN
ARG
LYS
TYR
LYS
ILE
LYS
ALA
GLN
GLU
GLY
ILE
VAL
SER
ASP
TYR
GLY
VAL
ARG
PHE
PHE

THR
THR
LYS
GLY
PRO
VAL
ALA
SER
ILE
ASN
LEU
THR
LYS
LEU
ASN
SER
LEU
SER
GLU
PRO
THR
LEU
VAL
THR
MET
PRO
ILE
GLY
TYR
VAL
THR
HIS
GLY
PHE
ASN
LEU
MET
GLU
ALA
THR
ALA
ILE
GLN
ARG
LYS
TYR
MET
SER
ARG
SER
LEU
LYS
ALA
PRO
ALA
VAL
VAL
SER
VAL
SER
SER
PRO
ASP
ALA
VAL
THR

THR
TYR
ASN
TYR
LEU
THR
SER
SER
SER
LYS
T132
R148
R156
R166
S177
P178
V179
L193
L196
L197
S198

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

- Molecule 1: Non-structural protein 3

Chain A:  29% • 66%

GLY
SER
HIS
MET
GLY
THR
VAL
SER
TRP
ASN
LEU
ARG
GLU
MET
LEU
ALA
HIS
ALA
ALA
GLU
THR
LEU
ARG
LYS
MET
LEU
PRO
ILE
CYS
TYR
MET
ASP
VAL
ARG
ALA
ILE
MET
LEU
ALA
THR
ILE
GLN
ARG
LYS
TYR
LYS
ILE
LYS
ALA
PRO
ALA
VAL
VAL
SER
VAL
SER
SER
PRO
ASP
ALA
VAL
THR

THR
THR
LYS
GLY
PRO
VAL
ALA
SER
ILE
ASN
LEU
THR
LYS
LEU
ASN
SER
LEU
SER
GLU
PRO
THR
LEU
VAL
THR
MET
PRO
ILE
GLY
TYR
VAL
THR
HIS
GLY
PHE
ASN
LEU
MET
GLU
ALA
THR
ALA
ILE
GLN
ARG
LYS
TYR
MET
SER
ARG
SER
LEU
LYS
ALA
PRO
ALA
VAL
VAL
SER
VAL
SER
SER
PRO
ASP
ALA
VAL
THR

THR
TYR
ASN
TYR
LEU
THR
SER
SER
SER
LYS
T132
E135
T140
R148
S151
R156
R166
S177
P178
V179
E180
S189
S198

5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 80 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
UNIO	refinement	
CANDID	refinement	
OPALp	refinement	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.62±0.01	0±0/550 (0.0± 0.0%)	1.10±0.05	1±1/742 (0.1± 0.1%)
All	All	0.62	0/11000 (0.0%)	1.10	16/14840 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.0±1.0
All	All	0	20

There are no bond-length outliers.

5 of 11 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	197	LEU	CB-CG-CD2	8.44	125.34	111.00	8	2
1	A	166	ARG	NE-CZ-NH2	-6.36	117.12	120.30	7	2
1	A	193	LEU	CD1-CG-CD2	6.13	128.89	110.50	16	3
1	A	197	LEU	CD1-CG-CD2	6.08	128.75	110.50	17	1
1	A	148	ARG	NE-CZ-NH2	-6.03	117.29	120.30	20	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	148	ARG	Sidechain	6
1	A	166	ARG	Sidechain	6
1	A	152	TYR	Sidechain	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	156	ARG	Sidechain	3
1	A	163	PHE	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	539	523	522	2±2
All	All	10780	10460	10440	46

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

5 of 24 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:183:LEU:HD13	1:A:184:ASP:HB3	0.64	1.67	12	1
1:A:193:LEU:CD1	1:A:197:LEU:HD21	0.57	2.29	20	3
1:A:182:HIS:CD2	1:A:187:VAL:HG22	0.57	2.35	12	2
1:A:141:VAL:HG11	1:A:164:LEU:HD11	0.56	1.77	4	1
1:A:174:THR:HG23	1:A:180:GLU:OE1	0.55	2.02	14	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	65/198 (33%)	55±2 (85±3%)	8±2 (12±3%)	2±1 (4±1%)	6	34
All	All	1300/3960 (33%)	1099 (85%)	154 (12%)	47 (4%)	6	34

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of

occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	177	SER	20
1	A	179	VAL	15
1	A	174	THR	4
1	A	148	ARG	2
1	A	153	SER	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	61/174 (35%)	53±3 (87±4%)	8±3 (13±4%)	7 49
All	All	1220/3480 (35%)	1063 (87%)	157 (13%)	7 49

5 of 37 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	177	SER	20
1	A	156	ARG	14
1	A	196	LEU	13
1	A	132	THR	8
1	A	166	ARG	8

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided