

wwPDB NMR Structure Validation Summary Report (i)

Jun 5, 2023 – 03:00 PM EDT

PDB ID : 2M26 BMRB ID : 18901

Title: NMR structure of the C-terminal domain of the protein HCFC1 from Mus

musculus

Authors: Serrano, P.; Geralt, M.; Dutta, S.K.; Wuthrich, K.; Joint Center for Structural

Genomics (JCSG); Partnership for Stem Cell Biology (STEMCELL)

Deposited on : 2012-12-14

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$rac{ m NMR~archive}{ m (\#Entries)}$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	126	71%	7%	22%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:6-A:46, A:62-A:118 (98)	1.41	4		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 16, 17, 18
2	13, 19
3	12, 14
Single-model clusters	9; 15; 20



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1706 atoms, of which 770 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called HCF C-terminal chain 1.

Mol	Chain	Residues	Atoms				Trace		
1	Λ	196	Total	С	Н	N	О	S	0
	A	126	1706	590	770	159	182	5	U

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q61191



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: HCF C-terminal chain 1



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 4. Colouring as in section 4.1 above.

• Molecule 1: HCF C-terminal chain 1





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: molecular dynamics.

Of the 80 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	
UNIO	structure solution	
OPALp	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1314
Number of shifts mapped to atoms	1180
Number of unparsed shifts	0
Number of shifts with mapping errors	134
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%



6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	ol Chain		ond lengths	Bond angles	
WIOI	Chain	RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.63 ± 0.01	$0\pm0/776~(~0.0\pm~0.0\%)$	1.06 ± 0.05	$1\pm 1/1060$ ($0.1\pm$ 0.1%)
All	All	0.63	0/15520 (0.0%)	1.06	18/21200 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	0.8 ± 0.7
All	All	0	17

There are no bond-length outliers.

5 of 15 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Chain Res 7		Atoma 7	7	Observed (0)	$\mathrm{Ideal}(^{o})$	Models	
IVIOI	Chain	nes	Type	Atoms	$oxed{\mathbf{Z} \left \; ext{Observed}(^o) \; \right }$		ideai()	Worst	Total
1	A	41	VAL	CA-CB-CG1	12.61	129.81	110.90	1	2
1	A	68	TYR	CB-CG-CD2	-7.12	116.72	121.00	1	1
1	A	67	VAL	CG1-CB-CG2	6.97	122.06	110.90	9	1
1	A	102	ARG	NE-CZ-NH2	-6.32	117.14	120.30	10	1
1	A	102	ARG	NE-CZ-NH1	5.90	123.25	120.30	1	1

There are no chirality outliers.

5 of 11 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	114	ARG	Sidechain	3
1	A	75	CYS	Peptide	2
1	A	89	TYR	Sidechain	2



Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	102	ARG	Sidechain	2
1	A	68	TYR	Sidechain	2

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	753	632	754	1±1
All	All	15060	12640	15080	25

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

5 of 16 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	$Distance(\mathring{A})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:41:VAL:HG12	1:A:68:TYR:HB2	0.72	1.60	9	2
1:A:116:LEU:H	1:A:116:LEU:HD22	0.69	1.47	7	4
1:A:92:LYS:HE3	1:A:116:LEU:HD21	0.53	1.81	9	1
1:A:100:ALA:HB3	1:A:107:TYR:CZ	0.52	2.40	12	1
1:A:102:ARG:HG2	1:A:107:TYR:CE1	0.51	2.40	15	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured		Outliers	Percentiles	
1	A	98/126 (78%)	81±3 (83±3%)	15±3 (15±3%)	2±2 (2±2%)	11	52
All	All	1960/2520 (78%)	1622 (83%)	298 (15%)	40 (2%)	11	52

5 of 22 unique Ramachandran outliers are listed below. They are sorted by the frequency of



occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	71	PRO	5
1	A	108	GLY	3
1	A	33	SER	3
1	A	46	GLN	3
1	A	31	VAL	3

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Percentiles		
1	A	82/104 (79%)	69±3 (84±4%)	13±3 (16±4%)	5	42	
All	All	1640/2080 (79%)	1383 (84%)	257 (16%)	5	42	

5 of 55 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	114	ARG	19
1	A	26	TRP	12
1	A	62	LEU	11
1	A	90	THR	11
1	A	102	ARG	10

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.



6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 82% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1314
Number of shifts mapped to atoms	1180
Number of unparsed shifts	0
Number of shifts with mapping errors	134
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

The following errors were found when reading this chemical shift list.

• Chemical shift has been reported more than once. All 3 occurrences are reported below.

Ligt ID	Chain	Pag	Trme	Type Atom		Shift Data		
LIST ID	Chain	nes	туре	Atom	Value	Uncertainty	Ambiguity	
1	A	43	LEU	HD11	0.691	0.000	•	
1	A	43	LEU	HD12	0.691	0.000	•	
1	A	43	LEU	HD13	0.691	0.000	•	

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

• No matching atom found in the structure. First 5 (of 134) occurrences are reported below.

List ID	Chain	Res Type	Atom	Shift Data			
LIST ID	Chain	nes	Type	Atom	Value	Uncertainty	Ambiguity
1	A	38	GLU	HB2	1.654	0.000	
1	A	38	GLU	HG2	1.878	0.000	•
1	A	39	TYR	HB2	3.159	0.000	
1	A	40	SER	HB2	3.467	0.000	
1	A	42	TYR	HB2	2.893	0.000	



 $Continued\ from\ previous\ page...$

	a from pr			A .		Shift Dat	a
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	43	LEU	HB2	1.319	0.000	
1	A	46	GLN	HB2	1.852	0.000	
1	A	47	SER	HB2	3.704	0.000	
1	A	49	GLN	HB2	2.075	0.000	
1	A	54	PRO	HD2	3.84	0.000	
1	A	55	LYS	HB2	1.741	0.000	
1	A	59	PRO	HB2	2.227	0.000	
1	A	61	GLN	HG2	2.293	0.000	
1	A	62	LEU	HB2	1.383	0.000	
1	A	64	PHE	HB2	2.683	0.000	
1	A	66	ARG	HB2	1.431	0.000	
1	A	66	ARG	HG2	1.251	0.000	
1	A	66	ARG	HD2	3.075	0.000	
1	A	68	TYR	HB2	3.491	0.000	
1	A	69	CYS	HB2	2.368	0.000	
1	A	76	LEU	HB2	1.577	0.000	
1	A	78	GLN	HB2	2.122	0.000	
1	A	78	GLN	HG2	2.519	0.000	
1	A	79	SER	HB2	3.882	0.000	
1	A	80	SER	HB2	3.888	0.000	
1	A	81	SER	HB2	3.78	0.000	
1	A	82	LEU	HB2	1.608	0.000	
1	A	83	SER	HB2	3.908	0.000	
1	A	84	ASN	HB2	2.636	0.000	
1	A	87	ILE	HG12	1.204	0.000	
1	A	88	ASP	HB2	2.368	0.000	
1	A	89	TYR	HB2	3.262	0.000	
1	A	93	PRO	HB2	1.698	0.000	
1	A	96	ILE	HG12	1.4	0.000	
1	A	97	PHE	HB2	2.534	0.000	
1	A	98	ARG	HB2	1.094	0.000	
1	A	98	ARG	HG2	1.16	0.000	
1	A	103	ASN	HB2	3.45	0.000	
1	A	104	GLU	HB2	2.023	0.000	
1	A	104	GLU	HG2	2.318	0.000	
1	A	105	LYS	HB2	1.658	0.000	
1	A	105	LYS	HG2	1.108	0.000	
1	A	107	TYR	HB2	3.009	0.000	
1	A	112	GLN	HB2	1.841	0.000	
1	A	114	ARG	HB2	1.708	0.000	
1	A	114	ARG	HG2	1.58	0.000	



 $Continued\ from\ previous\ page...$

Continue				A .		Shift Dat	a
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity
1	A	114	ARG	HD2	3.103	0.000	
1	A	6	PHE	HB2	2.753	0.000	
1	A	13	ILE	HG12	1.335	0.000	•
1	A	15	ILE	HG12	0.309	0.000	
1	A	16	SER	HB2	3.777	0.000	
1	A	17	LYS	HB2	2.064	0.000	
1	A	17	LYS	HG2	1.621	0.000	
1	A	17	LYS	HE2	3.021	0.000	
1	A	17	LYS	HD2	1.777	0.000	•
1	A	18	SER	HB2	3.98	0.000	•
1	A	20	ASP	HB2	2.736	0.000	
1	A	23	HIS	HB2	3.045	0.000	
1	A	24	LEU	HB2	1.441	0.000	
1	A	26	TRP	HB2	3.27	0.000	•
1	A	27	GLU	HB2	2.053	0.000	
1	A	27	GLU	HG2	2.265	0.000	•
1	A	35	LYS	HB2	1.603	0.000	
1	A	35	LYS	HG2	1.316	0.000	•
1	A	35	LYS	HE2	2.879	0.000	
1	A	36	ILE	HG12	1.595	0.000	
1	A	37	ILE	HG12	0.712	0.000	•
1	A	45	ILE	HG12	1.208	0.000	
1	A	46	GLN	HG2	2.304	0.000	•
1	A	53	GLU	HB2	2.026	0.000	•
1	A	61	GLN	HB2	2.041	0.000	
1	A	65	MET	HG2	2.537	0.000	•
1	A	86	HIS	HB2	2.592	0.000	
1	A	92	LYS	HB2	1.691	0.000	•
1	A	99	ILE	HG12	1.112	0.000	
1	A	102	ARG	HB2	0.632	0.000	
1	A	102	ARG	HD2	2.444	0.000	•
1	A	105	LYS	HE2	2.777	0.000	•
1	A	105	LYS	HD2	1.408	0.000	
1	A	112	GLN	HG2	2.108	0.000	•
1	A	115	TRP	HB2	3.533	0.000	
1	A	116	LEU	HB2	1.677	0.000	•
1	A	117	GLN	HB2	1.793	0.000	•
1	A	11	CYS	HB2	2.836	0.000	•
1	A	14	LYS	HB2	1.594	0.000	•
1	A	14	LYS	HG2	1.32	0.000	•
1	A	14	LYS	HE2	2.879	0.000	•



 $Continued\ from\ previous\ page...$

	$\frac{a \text{ from } pr}{a}$			A 4	Shift Data				
List ID	Chain	Res	Type	Atom	Value	Uncertainty	Ambiguity		
1	A	14	LYS	HD2	1.529	0.000			
1	A	19	PRO	HG2	1.965	0.000			
1	A	29	PRO	HD2	3.837	0.000			
1	A	30	SER	HB2	3.376	0.000			
1	A	35	LYS	HD2	1.581	0.000			
1	A	59	PRO	HD2	3.49	0.000			
1	A	59	PRO	HG2	2.012	0.000			
1	A	93	PRO	HD2	3.509	0.000			
1	A	93	PRO	HG2	1.476	0.000			
1	A	95	ILE	HG12	0.981	0.000			
1	A	7	PRO	HD2	3.652	0.000	•		
1	A	10	PRO	HD2	3.76	0.000	•		
1	A	10	PRO	HG2	1.622	0.000			
1	A	10	PRO	HB2	2.272	0.000	•		
1	A	92	LYS	HE2	2.948	0.000			
1	A	92	LYS	HG2	1.397	0.000			
1	A	71	PRO	HD2	3.963	0.000			
1	A	118	GLU	HG2	1.681	0.000			
1	A	118	GLU	HB2	1.525	0.000			
1	A	71	PRO	HG2	2.202	0.000			
1	A	71	PRO	HB2	2.358	0.000			
1	A	54	PRO	HG2	1.739	0.000			
1	A	54	PRO	HB2	2.234	0.000			
1	A	75	CYS	HB2	1.599	0.000	•		
1	A	28	PRO	HD2	3.658	0.000			
1	A	72	SER	HB2	3.86	0.000			
1	A	57	SER	HB2	3.817	0.000			
1	A	73	PRO	HB2	1.704	0.000			
1	A	73	PRO	HD2	3.494	0.000			
1	A	73	PRO	HG2	2.013	0.000			
1	A	51	SER	HB2	4.287	0.000			
1	A	55	LYS	HG2	1.374	0.000			
1	A	56	SER	HB2	4.366	0.000			
1	A	7	PRO	HG2	1.982	0.000			
1	A	19	PRO	HB2	1.841	0.000			
1	A	33	SER	HB2	4.192	0.000			
1	A	109	PRO	HD2	3.837	0.000			
1	A	109	PRO	HB2	2.189	0.000			
1	A	74	SER	HB2	3.374	0.000			
1	A	28	PRO	HG2	1.957	0.000			
1	A	28	PRO	HB2	1.857	0.000			



Continued from previous page...

List ID	Chain	Res	Type	Atom	Shift Data			
LIST ID	Chain	nes	туре	Atom	Value	Uncertainty	Ambiguity	
1	A	29	PRO	HB2	1.82	0.000	•	
1	A	29	PRO	HG2	2.203	0.000	•	
1	A	48	SER	HB2	3.872	0.000		
1	A	49	GLN	HG2	2.308	0.000	•	
1	A	53	GLU	HG2	2.246	0.000		
1	A	55	LYS	HE2	3.479	0.000		

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction} \pm {\rm precision}, ppm$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	113	0.00 ± 0.10	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	105	-0.07 ± 0.16	None needed ($< 0.5 \text{ ppm}$)
¹³ C′	85	-0.42 ± 0.22	None needed ($< 0.5 \text{ ppm}$)
^{15}N	102	1.06 ± 0.28	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1156 atoms were assigned a chemical shift out of a possible 1302. 0 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	448/478 (94%)	190/193 (98%)	170/196 (87%)	88/89 (99%)
Sidechain	624/711 (88%)	$420/469 \ (90\%)$	199/219 (91%)	5/23 (22%)
Aromatic	84/113 (74%)	41/55 (75%)	41/54 (76%)	2/4 (50%)
Overall	1156/1302 (89%)	651/717 (91%)	410/469 (87%)	95/116 (82%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	7	PRO	CA	4.19	55.85 - 70.84	-39.5
1	A	102	ARG	HB3	-0.15	0.43 - 3.11	-7.1
1	A	26	TRP	NE1	115.62	118.53 - 139.98	-6.4




Continued from previous page...

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	75	CYS	HB3	0.64	0.69 - 5.10	-5.1

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1525
Intra-residue ($ i-j =0$)	399
Sequential ($ i-j =1$)	431
Medium range ($ i-j >1$ and $ i-j <5$)	118
Long range (i-j ≥5)	577
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	449
Number of restraints per residue	12.1
Number of long range restraints per residue ¹	4.6

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	17.8	0.2
0.2-0.5 (Medium)	28.4	0.5
>0.5 (Large)	61.5	4.22



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Doodnointe tour	Count	% ¹	Vi	olated	3	Consis	tentl	\mathbf{y} Violated 4
Restraints type	Count	%0°	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	399	26.2	58	14.5	3.8	0	0.0	0.0
Backbone-Backbone	3	0.2	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	319	20.9	49	15.4	3.2	0	0.0	0.0
Sidechain-Sidechain	77	5.0	9	11.7	0.6	0	0.0	0.0
Sequential (i-j =1)	431	28.3	72	16.7	4.7	2	0.5	0.1
Backbone-Backbone	120	7.9	3	2.5	0.2	0	0.0	0.0
Backbone-Sidechain	278	18.2	59	21.2	3.9	2	0.7	0.1
Sidechain-Sidechain	33	2.2	10	30.3	0.7	0	0.0	0.0
Medium range ($ i-j >1 & i-j <5$)	118	7.7	32	27.1	2.1	1	0.8	0.1
Backbone-Backbone	22	1.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	41	2.7	12	29.3	0.8	0	0.0	0.0
Sidechain-Sidechain	55	3.6	20	36.4	1.3	1	1.8	0.1
Long range ($ i-j \ge 5$)	577	37.8	151	26.2	9.9	3	0.5	0.2
Backbone-Backbone	71	4.7	3	4.2	0.2	0	0.0	0.0
Backbone-Sidechain	218	14.3	50	22.9	3.3	0	0.0	0.0
Sidechain-Sidechain	288	18.9	98	34.0	6.4	3	1.0	0.2
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1525	100.0	313	20.5	20.5	6	0.4	0.4
Backbone-Backbone	216	14.2	6	2.8	0.4	0	0.0	0.0
Backbone-Sidechain	856	56.1	170	19.9	11.1	2	0.2	0.1
Sidechain-Sidechain	453	29.7	137	30.2	9.0	4	0.9	0.3

 $^{^1}$ percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	tions	,	Mean (Å)	N/1 (Å)	${ m SD}^6$ (Å)	Madian (8)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (Å)	$SD^*(A)$	Median (Å)
1	22	31	13	47	0	113	0.84	2.74	0.62	0.67
2	24	28	16	50	0	118	0.79	2.72	0.61	0.66
3	19	28	10	38	0	95	0.68	2.66	0.55	0.53
4	23	29	8	43	0	103	0.72	2.56	0.56	0.55
5	16	31	10	44	0	101	0.69	2.52	0.57	0.53
6	21	27	11	50	0	109	0.85	3.97	0.68	0.67
7	22	28	10	52	0	112	0.84	2.58	0.63	0.69
8	26	32	11	45	0	114	0.71	2.96	0.6	0.59
9	17	29	15	46	0	107	0.82	2.63	0.58	0.68
10	18	33	14	54	0	119	0.78	4.05	0.64	0.67
11	19	35	11	45	0	110	0.79	2.7	0.64	0.58




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	19	30	10	39	0	98	0.74	2.76	0.63	0.58
13	26	35	15	43	0	119	0.74	2.66	0.54	0.59
14	25	32	11	57	0	125	0.77	2.62	0.6	0.61
15	17	26	13	50	0	106	0.84	2.85	0.64	0.69
16	21	29	13	48	0	111	0.71	2.75	0.58	0.54
17	16	33	8	39	0	96	0.73	2.56	0.57	0.58
18	21	20	9	38	0	88	0.77	4.22	0.64	0.66
19	15	28	8	44	0	95	0.69	2.67	0.58	0.5
20	21	31	13	49	0	114	0.74	2.77	0.58	0.6

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1212(IR:341, SQ:359, MR:86, LR:426, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	restra	aints	Fraction of the ensemble		
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%	
12	13	4	38	0	67	1	5.0	
6	6	6	16	0	34	2	10.0	
4	3	1	14	0	22	3	15.0	
3	3	2	11	0	19	4	20.0	
8	4	3	13	0	28	5	25.0	
2	4	1	8	0	15	6	30.0	
2	3	3	2	0	10	7	35.0	
1	7	2	11	0	21	8	40.0	
2	2	0	4	0	8	9	45.0	
2	3	2	6	0	13	10	50.0	
0	2	1	1	0	4	11	55.0	
1	3	0	4	0	8	12	60.0	
1	1	0	2	0	4	13	65.0	
5	3	3	0	0	11	14	70.0	
3	0	1	1	0	5	15	75.0	
0	2	0	5	0	7	16	80.0	
1	4	0	3	0	8	17	85.0	
4	6	2	8	0	20	18	90.0	
1	1	0	1	0	3	19	95.0	
0	2	1	3	0	6	20	100.0	

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	\mathbf{SD}^1 (Å)	Median (Å)
(1,931)	1:A:41:VAL:HG11	1:A:68:TYR:HE1	20	2.52	0.83	2.57
(1,931)	1:A:41:VAL:HG12	1:A:68:TYR:HE1	20	2.52	0.83	2.57
(1,931)	1:A:41:VAL:HG13	1:A:68:TYR:HE1	20	2.52	0.83	2.57
(1,428)	1:A:42:TYR:HB3	1:A:98:ARG:HG3	20	1.14	0.69	1.02
(1,707)	1:A:35:LYS:HD3	1:A:104:GLU:HB3	20	1.05	0.56	1.04
(1,661)	1:A:22:ALA:HA	1:A:23:HIS:HB3	20	1.01	0.2	1.07
(1,1309)	1:A:10:PRO:HB3	1:A:13:ILE:HD11	20	0.91	0.23	0.89
(1,1309)	1:A:10:PRO:HB3	1:A:13:ILE:HD12	20	0.91	0.23	0.89
(1,1309)	1:A:10:PRO:HB3	1:A:13:ILE:HD13	20	0.91	0.23	0.89
(1,360)	1:A:23:HIS:HA	1:A:24:LEU:HB3	20	0.84	0.14	0.87
(1,427)	1:A:42:TYR:HB3	1:A:98:ARG:H	19	0.78	0.29	0.71
(1,854)	1:A:92:LYS:HG3	1:A:93:PRO:HD3	19	0.5	0.14	0.58
(1,749)	1:A:96:ILE:HA	1:A:96:ILE:HG13	19	0.22	0.04	0.23
(1,1096)	1:A:67:VAL:HG21	1:A:81:SER:HA	18	2.15	0.36	2.22


¹Number of violated models, ²Standard deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,931)	1:A:41:VAL:HG11	1:A:68:TYR:HE1	18	4.22
(1,931)	1:A:41:VAL:HG12	1:A:68:TYR:HE1	18	4.22
(1,931)	1:A:41:VAL:HG13	1:A:68:TYR:HE1	18	4.22
(1,931)	1:A:41:VAL:HG11	1:A:68:TYR:HE1	10	4.05
(1,931)	1:A:41:VAL:HG12	1:A:68:TYR:HE1	10	4.05
(1,931)	1:A:41:VAL:HG13	1:A:68:TYR:HE1	10	4.05
(1,931)	1:A:41:VAL:HG11	1:A:68:TYR:HE1	6	3.97
(1,931)	1:A:41:VAL:HG12	1:A:68:TYR:HE1	6	3.97
(1,931)	1:A:41:VAL:HG13	1:A:68:TYR:HE1	6	3.97
(1,1514)	1:A:39:TYR:H	1:A:68:TYR:HE1	8	2.96
(1,845)	1:A:42:TYR:HB3	1:A:62:LEU:HD21	15	2.85



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,845)	1:A:42:TYR:HB3	1:A:62:LEU:HD22	15	2.85
(1,845)	1:A:42:TYR:HB3	1:A:62:LEU:HD23	15	2.85
(1,277)	1:A:68:TYR:HE1	1:A:70:GLY:H	8	2.84
(1,845)	1:A:42:TYR:HB3	1:A:62:LEU:HD21	20	2.77
(1,845)	1:A:42:TYR:HB3	1:A:62:LEU:HD22	20	2.77
(1,845)	1:A:42:TYR:HB3	1:A:62:LEU:HD23	20	2.77
(1,707)	1:A:35:LYS:HD3	1:A:104:GLU:HB3	15	2.77
(1,931)	1:A:41:VAL:HG11	1:A:68:TYR:HE1	12	2.76
(1,931)	1:A:41:VAL:HG12	1:A:68:TYR:HE1	12	2.76
(1,931)	1:A:41:VAL:HG13	1:A:68:TYR:HE1	12	2.76
(1,931)	1:A:41:VAL:HG11	1:A:68:TYR:HE1	16	2.75



10 Dihedral-angle violation analysis (i)

No dihedral-angle restraints found

