



# Full wwPDB NMR Structure Validation Report ⓘ

Jun 6, 2023 – 04:54 AM EDT

PDB ID : 2MMU  
BMRB ID : 19867  
Title : Structure of CrgA, a Cell Division Structural and Regulatory Protein from Mycobacterium tuberculosis, in Lipid Bilayers  
Authors : Das, N.; Dai, J.; Hung, I.; Rajagopalan, M.; Zhou, H.; Cross, T.A.  
Deposited on : 2014-03-18

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
BMRB Restraints Analysis : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.33


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLID-STATE NMR*

The overall completeness of chemical shifts assignment is 5%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	101	 36% 12% . 50%

## 2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 846 atoms, of which 431 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Cell division protein CrgA.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	50	846	291	431	61	59	4	0


There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	94	LEU	-	expression tag	UNP P67376
A	95	GLU	-	expression tag	UNP P67376
A	96	HIS	-	expression tag	UNP P67376
A	97	HIS	-	expression tag	UNP P67376
A	98	HIS	-	expression tag	UNP P67376
A	99	HIS	-	expression tag	UNP P67376
A	100	HIS	-	expression tag	UNP P67376
A	101	HIS	-	expression tag	UNP P67376

## 4 Residue-property plots [i](#)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

### • Molecule 1: Cell division protein CrgA

Chain A:  36% 12% 50%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, molecular dynamics*.

Of the 1000 calculated structures, 1 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.34
NAMD	refinement	2.9
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	40
Number of shifts mapped to atoms	40
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	5%

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.71	4/429 ( 0.9%)	1.92	10/583 ( 1.7%)
All	All	1.71	4/429 ( 0.9%)	1.92	10/583 ( 1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0	1
All	All	0	1

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	79	PHE	CG-CD1	6.28	1.48	1.38
1	A	82	MET	CA-CB	5.87	1.66	1.53
1	A	35	SER	CA-CB	5.15	1.60	1.52
1	A	37	PHE	CG-CD2	5.08	1.46	1.38

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	33	PHE	CB-CG-CD1	-8.32	114.97	120.80
1	A	81	PHE	CB-CG-CD2	7.90	126.33	120.80
1	A	75	TYR	CB-CG-CD1	7.58	125.55	121.00
1	A	89	THR	CA-CB-CG2	-6.97	102.65	112.40
1	A	33	PHE	CB-CG-CD2	6.85	125.59	120.80
1	A	81	PHE	CB-CG-CD1	-6.79	116.05	120.80
1	A	89	THR	CA-CB-OG1	6.32	122.26	109.00
1	A	91	ARG	NE-CZ-NH1	6.25	123.42	120.30
1	A	49	MET	CG-SD-CE	-5.99	90.61	100.20
1	A	75	TYR	CG-CD2-CE2	5.79	125.93	121.30

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	75	TYR	Sidechain

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	415	431	429	2
All	All	415	431	429	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:73:TRP:CE2	1:A:77:ILE:HD11	0.46	2.46
1:A:93:HIS:CE1	1:A:94:LEU:HD13	0.41	2.51

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	46/101 (46%)	42 (91%)	3 (7%)	1 (2%)	10	49
All	All	46/101 (46%)	42 (91%)	3 (7%)	1 (2%)	10	49

All 1 Ramachandran outliers are listed below.

Mol	Chain	Res	Type
1	A	94	LEU



### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	43/86 (50%)	43 (100%)	0 (0%)	100	100
All	All	43/86 (50%)	43 (100%)	0 (0%)	100	100

There are no protein residues with a non-rotameric sidechain to report.

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation i

The completeness of assignment taking into account all chemical shift lists is 5% for the well-defined parts and 5% for the entire structure.

### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping i

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	40
Number of shifts mapped to atoms	40
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	40

#### 7.1.2 Chemical shift referencing i

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	None (insufficient data)
$^{13}\text{C}_\beta$	0	—	None (insufficient data)
$^{13}\text{C}'$	0	—	None (insufficient data)
$^{15}\text{N}$	40	0.00 $\pm$ 0.00	None needed (< 0.5 ppm)

#### 7.1.3 Completeness of resonance assignments i

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 5%, i.e. 40 atoms were assigned a chemical shift out of a possible 758. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	40/252 (16%)	0/103 (0%)	0/100 (0%)	40/49 (82%)
Sidechain	0/392 (0%)	0/267 (0%)	0/120 (0%)	0/5 (0%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	0/114 (0%)	0/57 (0%)	0/52 (0%)	0/5 (0%)
Overall	40/758 (5%)	0/427 (0%)	0/272 (0%)	40/59 (68%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 5%, i.e. 40 atoms were assigned a chemical shift out of a possible 758. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	40/252 (16%)	0/103 (0%)	0/100 (0%)	40/49 (82%)
Sidechain	0/392 (0%)	0/267 (0%)	0/120 (0%)	0/5 (0%)
Aromatic	0/114 (0%)	0/57 (0%)	0/52 (0%)	0/5 (0%)
Overall	40/758 (5%)	0/427 (0%)	0/272 (0%)	40/59 (68%)

Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	35	SER	N	227.50	99.14 – 133.45	32.4
1	A	39	GLY	N	220.40	91.59 – 127.52	30.9
1	A	80	ALA	N	228.50	106.13 – 140.55	30.6
1	A	49	MET	N	222.50	102.99 – 137.21	29.9
1	A	87	LEU	N	230.40	102.77 – 140.89	28.5
1	A	76	ALA	N	220.90	106.13 – 140.55	28.3
1	A	42	LEU	N	226.00	102.77 – 140.89	27.3
1	A	45	LEU	N	224.60	102.77 – 140.89	27.0
1	A	85	GLY	N	203.30	91.59 – 127.52	26.1
1	A	73	TRP	N	224.60	101.51 – 141.60	25.7
1	A	51	PHE	N	225.10	99.93 – 140.82	25.6
1	A	88	LEU	N	219.10	102.77 – 140.89	25.5
1	A	90	MET	N	207.40	102.99 – 137.21	25.5
1	A	46	ILE	N	225.30	100.55 – 142.30	24.9
1	A	77	ILE	N	224.80	100.55 – 142.30	24.8
1	A	48	LEU	N	215.80	102.77 – 140.89	24.6

*Continued on next page...*


Continued from previous page...

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	41	MET	N	202.90	102.99 – 137.21	24.2
1	A	79	PHE	N	219.10	99.93 – 140.82	24.1
1	A	38	ILE	N	222.00	100.55 – 142.30	24.1
1	A	74	ASN	N	208.90	99.66 – 138.23	23.3
1	A	83	ILE	N	218.10	100.55 – 142.30	23.2
1	A	84	THR	N	223.00	91.89 – 138.78	23.0
1	A	31	VAL	N	221.30	99.23 – 142.92	22.9
1	A	78	ALA	N	200.80	106.13 – 140.55	22.5
1	A	36	LEU	N	206.90	102.77 – 140.89	22.3
1	A	32	TRP	N	210.30	101.51 – 141.60	22.1
1	A	81	PHE	N	210.00	99.93 – 140.82	21.9
1	A	37	PHE	N	209.40	99.93 – 140.82	21.8
1	A	82	MET	N	193.70	102.99 – 137.21	21.5
1	A	86	LEU	N	203.80	102.77 – 140.89	21.5
1	A	89	THR	N	215.80	91.89 – 138.78	21.4
1	A	43	ILE	N	208.60	100.55 – 142.30	20.9
1	A	34	VAL	N	211.50	99.23 – 142.92	20.7
1	A	33	PHE	N	203.50	99.93 – 140.82	20.3
1	A	40	LEU	N	198.30	102.77 – 140.89	20.1
1	A	75	TYR	N	189.70	100.12 – 140.79	17.0
1	A	47	TRP	N	188.60	101.51 – 141.60	16.7
1	A	91	ARG	N	177.30	102.91 – 138.82	15.7
1	A	53	LEU	N	174.30	102.77 – 140.89	13.8
1	A	50	VAL	N	171.30	99.23 – 142.92	11.5

### 7.1.5 Random Coil Index (RCI) plots

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



## 8 NMR restraints analysis

No restraints data found

## 9 Distance violation analysis

No distance restraints data found

## 10 Dihedral-angle violation analysis

No dihedral-angle restraints found