

wwPDB NMR Structure Validation Summary Report (i)

Jun 3, 2023 – 05:17 PM EDT

PDB ID	:	2MMZ
BMRB ID	:	19872
Title	:	Solution structure of the apo form of human glutaredoxin 5
Authors	:	Banci, L.; Brancaccio, D.; Ciofi-Baffoni, S.; Del Conte, R.; Gadepalli, R.;
		Mikolajczyk, M.; Neri, S.; Piccioli, M.; Winkelmann, J.
Deposited on	:	2014-03-25

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 80%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	${f NMR} ext{ archive} \ (\# ext{Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	А	121	74%	14%	••	10%	



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 12 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core	Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1 A:42-A:150 (109) 0.61 12						

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	$1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, \\18, 19$
2	13, 20



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1840 atoms, of which 915 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Glutaredoxin-related protein 5, mitochondrial.

Mol	Chain	Residues	Atoms				Trace		
1	٨	191	Total	С	Н	Ν	0	S	0
	A	121	1840	586	915	157	178	4	

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q86SX6
A	2	SER	-	expression tag	UNP Q86SX6
А	3	PHE	-	expression tag	UNP Q86SX6
А	4	THR	-	expression tag	UNP Q86SX6
А	5	MET	-	expression tag	UNP Q86SX6




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Glutaredoxin-related protein 5, mitochondrial



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 12. Colouring as in section 4.1 above.

• Molecule 1: Glutared oxin-related protein 5, mitochondrial





5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 80 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1274
Number of shifts mapped to atoms	1274
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	80%



6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol C	Chain	B	ond lengths	Bond angles		
	Unam	RMSZ	#Z > 5	RMSZ	#Z > 5	
1	А	$0.65 {\pm} 0.01$	$0{\pm}0/874~(~0.0{\pm}~0.0\%)$	$1.19{\pm}0.03$	$8{\pm}2/1185~(~0.6{\pm}~0.1\%)$	
All	All	0.65	0/17480~(~0.0%)	1.19	151/23700 ($0.6%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	А	$0.0{\pm}0.0$	$2.6{\pm}1.2$
All	All	0	53

There are no bond-length outliers.

5 of 16 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Z Observed $(^{o})$		Moo	dels
	Ullalli	nes	туре	Atoms		Observed()	$\mathrm{Ideal}(^{o})$	Worst	Total
1	А	83	ARG	NE-CZ-NH1	9.64	125.12	120.30	9	18
1	А	97	ARG	NE-CZ-NH1	9.02	124.81	120.30	17	14
1	А	78	ARG	NE-CZ-NH1	7.96	124.28	120.30	9	16
1	А	85	TYR	CB-CG-CD1	-7.91	116.26	121.00	17	18
1	А	118	PHE	CB-CG-CD2	-7.67	115.43	120.80	3	11

There are no chirality outliers.

5 of 10 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	А	118	PHE	Peptide	15
1	А	119	VAL	Peptide	13
1	А	85	TYR	Sidechain	10



Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	А	109	ILE	Peptide	4
1	А	103	TYR	Sidechain	3

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	858	855	855	0±1
All	All	17160	17100	17100	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:91:LEU:H	1:A:91:LEU:HD23	0.70	1.45	9	1
1:A:91:LEU:H	1:A:91:LEU:CD2	0.62	2.08	9	1
1:A:89:ASN:ND2	1:A:91:LEU:HD21	0.57	2.15	9	1
1:A:91:LEU:HD23	1:A:91:LEU:N	0.54	2.17	9	1
1:A:69:PHE:CD2	1:A:122:CYS:HB3	0.43	2.49	16	1

All unique clashes are listed below, sorted by their clash magnitude.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	108/121~(89%)	$100\pm2~(93\pm2\%)$	$6\pm2~(6\pm2\%)$	$1\pm1 (1\pm1\%)$	18	66
All	All	2160/2420~(89%)	2010~(93%)	126~(6%)	24 (1%)	18	66

5 of 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of



Mol	Chain	\mathbf{Res}	Type	Models (Total)
1	А	119	VAL	14
1	А	69	PHE	4
1	А	108	THR	2
1	А	66	GLN	2
1	А	67	CYS	1

occurrence in the ensemble.

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	94/99~(95%)	81 ± 2 (86±3%)	$13\pm2~(14\pm3\%)$	7 47
All	All	1880/1980~(95%)	1618 (86%)	262 (14%)	7 47

5 of 37 unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	53	LYS	20
1	А	54	VAL	20
1	А	58	LEU	20
1	А	126	LEU	20
1	А	106	TRP	19

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.



6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 80% for the well-defined parts and 79% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1274
Number of shifts mapped to atoms	1274
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction}\pm{\rm precision},ppm$	Suggested action
$^{13}C_{\alpha}$	120	0.41 ± 0.14	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	106	0.77 ± 0.13	Should be applied
$^{13}C'$	117	0.19 ± 0.23	None needed (< 0.5 ppm)
¹⁵ N	113	0.68 ± 0.35	None needed (imprecise)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 80%, i.e. 1201 atoms were assigned a chemical shift out of a possible 1500. 0 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	427/544~(78%)	109/222 (49%)	215/218~(99%)	103/104~(99%)
Sidechain	740/854~(87%)	502/555~(90%)	238/270~(88%)	0/29~(0%)



	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Aromatic	34/102~(33%)	29/49~(59%)	0/46~(0%)	5/7~(71%)
Overall	1201/1500~(80%)	640/826~(77%)	453/534 ($85%$)	108/140~(77%)

Continued from previous page...


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1995
Intra-residue (i-j =0)	536
Sequential (i-j =1)	534
Medium range ($ i-j >1$ and $ i-j <5$)	391
Long range $(i-j \ge 5)$	496
Inter-chain	0
Hydrogen bond restraints	38
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	16.5
Number of long range restraints per residue ¹	4.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	22.2	0.2
0.2-0.5 (Medium)	3.8	0.5
>0.5 (Large)	4.3	1.4



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destroints type	Count	$\%^1$	Vic	lated	3	Consis	tentl	y Violated ⁴
Restraints type	Count	701	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	536	26.9	32	6.0	1.6	2	0.4	0.1
Backbone-Backbone	5	0.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	426	21.4	29	6.8	1.5	2	0.5	0.1
Sidechain-Sidechain	105	5.3	3	2.9	0.2	0	0.0	0.0
Sequential (i-j =1)	534	26.8	25	4.7	1.3	2	0.4	0.1
Backbone-Backbone	149	7.5	2	1.3	0.1	0	0.0	0.0
Backbone-Sidechain	317	15.9	16	5.0	0.8	0	0.0	0.0
Sidechain-Sidechain	68	3.4	7	10.3	0.4	2	2.9	0.1
Medium range ($ i-j > 1 \& i-j < 5$)	391	19.6	18	4.6	0.9	1	0.3	0.1
Backbone-Backbone	107	5.4	2	1.9	0.1	0	0.0	0.0
Backbone-Sidechain	175	8.8	9	5.1	0.5	1	0.6	0.1
Sidechain-Sidechain	109	5.5	7	6.4	0.4	0	0.0	0.0
Long range $(i-j \ge 5)$	496	24.9	31	6.2	1.6	3	0.6	0.2
Backbone-Backbone	58	2.9	2	3.4	0.1	0	0.0	0.0
Backbone-Sidechain	202	10.1	11	5.4	0.6	0	0.0	0.0
Sidechain-Sidechain	236	11.8	18	7.6	0.9	3	1.3	0.2
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	38	1.9	1	2.6	0.1	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1995	100.0	107	5.4	5.4	8	0.4	0.4
Backbone-Backbone	357	17.9	7	2.0	0.4	0	0.0	0.0
Backbone-Sidechain	1120	56.1	65	5.8	3.3	3	0.3	0.2
Sidechain-Sidechain	518	26.0	35	6.8	1.8	5	1.0	0.3

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	5	Maan (Å)	Mar (Å)	SD^6 (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (Å)	Max (Å)	$SD^{*}(A)$	Median (Å)
1	8	7	3	10	0	28	0.24	1.12	0.23	0.14
2	9	5	4	8	0	26	0.26	0.93	0.23	0.14
3	8	9	4	8	0	29	0.24	1.4	0.27	0.13
4	9	6	3	8	0	26	0.21	0.85	0.16	0.14
5	8	10	7	12	0	37	0.34	1.19	0.33	0.14
6	11	9	4	8	0	32	0.2	0.77	0.17	0.13
7	8	4	5	9	0	26	0.25	0.84	0.23	0.14
8	10	7	6	7	0	30	0.2	0.84	0.15	0.15
9	9	7	5	13	0	34	0.24	0.91	0.21	0.15
10	13	4	3	8	0	28	0.25	0.92	0.23	0.15
11	12	8	3	5	0	28	0.23	0.75	0.19	0.14




Madal ID	Number of violations					5	Mean (Å)	Max (Å)		Madian (Å)
Model ID	IR^{1}	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	SD^{6} (Å)	Median (Å)
12	12	8	6	9	0	35	0.26	1.32	0.26	0.14
13	8	7	3	9	0	27	0.21	0.7	0.18	0.13
14	12	11	3	7	0	33	0.23	0.77	0.2	0.14
15	6	10	5	8	0	29	0.26	0.9	0.2	0.16
16	8	6	3	7	0	24	0.28	0.88	0.23	0.15
17	11	9	3	7	0	30	0.24	0.89	0.23	0.14
18	7	10	4	7	0	28	0.21	0.74	0.15	0.14
19	12	9	3	9	0	33	0.22	0.78	0.16	0.14
20	11	7	10	14	0	42	0.35	1.4	0.35	0.15

Continued from previous page...

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation





The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1851(IR:504, SQ:509, MR:373, LR:465, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio		restra	aints	Fractio	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC ⁵	Total	Count^6	%
9	10	6	10	0	35	1	5.0
6	2	7	5	0	20	2	10.0
3	3	2	3	0	11	3	15.0
3	2	0	2	0	7	4	20.0
0	0	0	4	0	4	5	25.0
2	2	0	0	0	4	6	30.0
2	0	0	1	0	3	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
1	0	0	1	0	2	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	1	0	0	0	1	16	80.0
2	1	0	0	0	3	17	85.0
1	1	1	2	0	5	18	90.0
1	1	1	0	0	3	19	95.0
2	2	1	3	0	8	20	100.0

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations




Intra-residue

MediumRange LongRange

Sequential

InterChain




9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,1016)	1:A:55:VAL:HG21	1:A:88:TYR:HD1	20	0.78	0.07	0.76
(1,1016)	1:A:55:VAL:HG21	1:A:88:TYR:HD2	20	0.78	0.07	0.76
(1,1016)	1:A:55:VAL:HG22	1:A:88:TYR:HD1	20	0.78	0.07	0.76
(1,1016)	1:A:55:VAL:HG22	1:A:88:TYR:HD2	20	0.78	0.07	0.76
(1,1016)	1:A:55:VAL:HG23	1:A:88:TYR:HD1	20	0.78	0.07	0.76
(1,1016)	1:A:55:VAL:HG23	1:A:88:TYR:HD2	20	0.78	0.07	0.76
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD11	20	0.73	0.27	0.75
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD12	20	0.73	0.27	0.75
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD13	20	0.73	0.27	0.75
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD11	20	0.73	0.27	0.75
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD12	20	0.73	0.27	0.75
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD13	20	0.73	0.27	0.75
(1,949)	1:A:58:LEU:HD11	1:A:85:TYR:HD1	20	0.52	0.12	0.54
(1,949)	1:A:58:LEU:HD11	1:A:85:TYR:HD2	20	0.52	0.12	0.54
(1,949)	1:A:58:LEU:HD12	1:A:85:TYR:HD1	20	0.52	0.12	0.54
(1,949)	1:A:58:LEU:HD12	1:A:85:TYR:HD2	20	0.52	0.12	0.54



Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,949)	1:A:58:LEU:HD13	1:A:85:TYR:HD1	20	0.52	0.12	0.54
(1,949)	1:A:58:LEU:HD13	1:A:85:TYR:HD2	20	0.52	0.12	0.54
(1,879)	1:A:43:GLU:HG2	1:A:44:GLN:HB2	20	0.52	0.16	0.54
(1,879)	1:A:43:GLU:HG2	1:A:44:GLN:HB3	20	0.52	0.16	0.54
(1,879)	1:A:43:GLU:HG3	1:A:44:GLN:HB2	20	0.52	0.16	0.54
(1,879)	1:A:43:GLU:HG3	1:A:44:GLN:HB3	20	0.52	0.16	0.54
(1,731)	1:A:85:TYR:HB3	1:A:87:ALA:HA	20	0.19	0.02	0.18
(1,702)	1:A:56:VAL:HG21	1:A:57:PHE:HB2	20	0.14	0.01	0.15
(1,702)	1:A:56:VAL:HG22	1:A:57:PHE:HB2	20	0.14	0.01	0.15
(1,702)	1:A:56:VAL:HG23	1:A:57:PHE:HB2	20	0.14	0.01	0.15
(1,1440)	1:A:100:ILE:HA	1:A:100:ILE:HG12	20	0.14	0.01	0.14
(1,843)	1:A:73:VAL:H	1:A:73:VAL:HB	20	0.13	0.01	0.13
(1,1290)	1:A:48:LEU:HD11	1:A:50:LYS:H	19	0.29	0.07	0.29
(1,1290)	1:A:48:LEU:HD12	1:A:50:LYS:H	19	0.29	0.07	0.29
(1,1290)	1:A:48:LEU:HD13	1:A:50:LYS:H	19	0.29	0.07	0.29
(1,1290)	1:A:48:LEU:HD21	1:A:50:LYS:H	19	0.29	0.07	0.29
(1,1290)	1:A:48:LEU:HD22	1:A:50:LYS:H	19	0.29	0.07	0.29
(1,1290)	1:A:48:LEU:HD23	1:A:50:LYS:H	19	0.29	0.07	0.29
(1,1208)	1:A:95:GLU:H	1:A:95:GLU:HB3	19	0.14	0.01	0.14

Continued from previous page...


¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD11	3	1.4
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD12	3	1.4
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD13	3	1.4
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD11	3	1.4
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD12	3	1.4
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD13	3	1.4
(1,1091)	1:A:109:ILE:HG21	1:A:111:GLN:HG2	20	1.4
(1,1091)	1:A:109:ILE:HG21	1:A:111:GLN:HG3	20	1.4
(1,1091)	1:A:109:ILE:HG22	1:A:111:GLN:HG2	20	1.4
(1,1091)	1:A:109:ILE:HG22	1:A:111:GLN:HG3	20	1.4
(1,1091)	1:A:109:ILE:HG23	1:A:111:GLN:HG2	20	1.4
(1,1091)	1:A:109:ILE:HG23	1:A:111:GLN:HG3	20	1.4
(1,1091)	1:A:109:ILE:HG21	1:A:111:GLN:HG2	12	1.32
(1,1091)	1:A:109:ILE:HG21	1:A:111:GLN:HG3	12	1.32
(1,1091)	1:A:109:ILE:HG22	1:A:111:GLN:HG2	12	1.32
(1,1091)	1:A:109:ILE:HG22	1:A:111:GLN:HG3	12	1.32
(1,1091)	1:A:109:ILE:HG23	1:A:111:GLN:HG2	12	1.32



Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1091)	1:A:109:ILE:HG23	1:A:111:GLN:HG3	12	1.32
(1,1213)	1:A:89:ASN:HD22	1:A:91:LEU:H	20	1.24
(1,1213)	1:A:89:ASN:HD22	1:A:91:LEU:H	5	1.19
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD11	1	1.12
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD12	1	1.12
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD13	1	1.12
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD11	1	1.12
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD12	1	1.12
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD13	1	1.12
(1,333)	1:A:89:ASN:HD22	1:A:91:LEU:HG	20	1.06
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD11	5	1.03
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD12	5	1.03
(1,1831)	1:A:57:PHE:HD1	1:A:96:LEU:HD13	5	1.03
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD11	5	1.03
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD12	5	1.03
(1,1831)	1:A:57:PHE:HD2	1:A:96:LEU:HD13	5	1.03
(1,1406)	1:A:60:GLY:HA2	1:A:89:ASN:HD22	5	1.02
(1,1858)	1:A:144:HIS:HE1	1:A:148:LEU:HD11	20	1.0

Continued from previous page...



10 Dihedral-angle violation analysis (i)

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value

