



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2023 – 08:49 AM EDT

PDB ID : 2R2N  
Title : The crystal structure of human kynurenine aminotransferase II in complex with kynurenine  
Authors : Han, Q.; Robinson, H.; Li, J.  
Deposited on : 2007-08-27  
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	425	
1	B	425	
1	C	425	
1	D	425	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	429	-	-	X	-
4	GOL	C	429	-	-	X	-
4	GOL	D	427	-	-	X	-
4	GOL	D	429	-	-	X	-

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 14831 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kynurenine/alpha-aminoadipate aminotransferase mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	425	Total 3332	C 2139	N 559	O 616	S 18	0	0	0
1	B	425	Total 3333	C 2139	N 559	O 617	S 18	0	0	0
1	C	425	Total 3332	C 2139	N 559	O 616	S 18	0	0	0
1	D	425	Total 3333	C 2139	N 559	O 617	S 18	0	0	0

- Molecule 2 is 4'-DEOXY-4'-AMINOPYRIDOXAL-5'-PHOSPHATE (three-letter code: PMP) (formula: C<sub>8</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 16	C 8	N 2	O 5	P 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			16	8	2	5	1		
2	C	1	Total	C	N	O	P	0	0
			16	8	2	5	1		
2	D	1	Total	C	N	O	P	0	0
			16	8	2	5	1		

- Molecule 3 is (2S)-2-amino-4-(2-aminophenyl)-4-oxobutanoic acid (three-letter code: KYN) (formula: C<sub>10</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	10	2	3		
3	A	1	Total	C	N	O	0	0
			15	10	2	3		
3	C	1	Total	C	N	O	0	0
			15	10	2	3		
3	C	1	Total	C	N	O	0	0
			15	10	2	3		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	314	Total O 314 314	0	0
5	B	325	Total O 325 325	0	0
5	C	352	Total O 352 352	0	0
5	D	350	Total O 350 350	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial




- Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial





- Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial





• Molecule 1: Kynurenine/alpha-aminoadipate aminotransferase mitochondrial





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.84Å 109.36Å 119.35Å 90.00° 94.68° 90.00°	Depositor
Resolution (Å)	26.68 – 1.95 26.67 – 1.95	Depositor EDS
% Data completeness (in resolution range)	93.1 (26.68-1.95) 93.1 (26.67-1.95)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.93 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.194 , 0.223 0.194 , 0.224	Depositor DCC
$R_{free}$ test set	6185 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.1	Xtrriage
Anisotropy	0.020	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	14831	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3905e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PMP, KYN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.76	0/3414	0.85	5/4634 (0.1%)
1	B	0.77	0/3415	0.88	6/4634 (0.1%)
1	C	0.84	1/3414 (0.0%)	0.91	8/4634 (0.2%)
1	D	0.87	0/3415	0.94	9/4634 (0.2%)
All	All	0.81	1/13658 (0.0%)	0.89	28/18536 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	C	0	3
1	D	0	2
All	All	0	10

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	112	CYS	CB-SG	-5.07	1.73	1.81

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	283	ARG	NE-CZ-NH1	13.52	127.06	120.30
1	B	257	ARG	NE-CZ-NH2	-12.38	114.11	120.30
1	A	257	ARG	NE-CZ-NH2	-12.16	114.22	120.30
1	B	257	ARG	NE-CZ-NH1	11.55	126.08	120.30
1	C	321	ARG	NE-CZ-NH2	-11.26	114.67	120.30

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	180	PRO	Peptide
1	A	181	GLU	Peptide
1	A	187	GLN	Peptide
1	A	341	LEU	Peptide
1	A	342	THR	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3332	0	3353	131	0
1	B	3333	0	3353	105	0
1	C	3332	0	3353	109	0
1	D	3333	0	3353	94	0
2	A	16	0	11	3	0
2	B	16	0	11	2	0
2	C	16	0	11	2	0
2	D	16	0	11	4	0
3	A	30	0	23	4	0
3	C	30	0	22	5	0
4	A	6	0	7	6	0
4	C	6	0	8	6	0
4	D	24	0	32	12	0
5	A	314	0	0	20	0
5	B	325	0	0	17	0
5	C	352	0	0	19	0
5	D	350	0	0	12	0
All	All	14831	0	13548	433	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 433 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:SER:HB2	1:B:18:PRO:HA	1.23	1.17
1:C:370:ILE:HD11	1:C:398:LEU:CD2	1.76	1.14
1:C:370:ILE:HD11	1:C:398:LEU:HD21	1.15	1.09
1:D:381:MET:HE2	1:D:398:LEU:HD13	1.34	1.05
1:D:78:ALA:HB3	4:D:427:GOL:H12	1.34	1.04

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423/425 (100%)	402 (95%)	12 (3%)	9 (2%)	<b>7</b> <b>1</b>
1	B	423/425 (100%)	409 (97%)	13 (3%)	1 (0%)	47 38
1	C	423/425 (100%)	405 (96%)	15 (4%)	3 (1%)	<b>22</b> <b>11</b>
1	D	423/425 (100%)	408 (96%)	12 (3%)	3 (1%)	<b>22</b> <b>11</b>
All	All	1692/1700 (100%)	1624 (96%)	52 (3%)	16 (1%)	<b>17</b> <b>8</b>

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	PRO
1	A	19	ILE
1	A	182	ASP
1	A	184	LYS
1	B	17	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/370 (100%)	330 (89%)	40 (11%)	6	1
1	B	370/370 (100%)	337 (91%)	33 (9%)	9	2
1	C	370/370 (100%)	337 (91%)	33 (9%)	9	2
1	D	370/370 (100%)	339 (92%)	31 (8%)	11	3
All	All	1480/1480 (100%)	1343 (91%)	137 (9%)	9	2

5 of 137 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	177	ARG
1	D	221	ARG
1	D	313	GLU
1	B	99	THR
1	B	87	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	118	GLN
1	C	318	HIS
1	D	419	GLN
1	C	187	GLN
1	C	294	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	D	429	-	5,5,5	0.66	0	5,5,5	0.70	0
3	KYN	A	427	-	13,15,15	1.85	2 (15%)	17,20,20	1.50	4 (23%)
4	GOL	D	427	-	5,5,5	0.55	0	5,5,5	0.98	0
2	PMP	D	426	-	16,16,16	1.28	2 (12%)	21,23,23	1.39	4 (19%)
3	KYN	A	428	-	13,15,15	1.70	1 (7%)	17,20,20	1.02	1 (5%)
4	GOL	C	429	-	5,5,5	0.79	0	5,5,5	0.76	0
4	GOL	A	429	-	5,5,5	1.16	1 (20%)	5,5,5	1.10	1 (20%)
3	KYN	C	428	-	13,15,15	1.48	2 (15%)	17,20,20	1.41	4 (23%)
4	GOL	D	430	-	5,5,5	0.47	0	5,5,5	0.59	0
2	PMP	B	426	-	16,16,16	1.29	1 (6%)	21,23,23	1.40	3 (14%)
2	PMP	C	426	-	16,16,16	1.25	2 (12%)	21,23,23	1.39	3 (14%)
2	PMP	A	426	-	16,16,16	1.28	2 (12%)	21,23,23	1.56	4 (19%)
3	KYN	C	427	-	13,15,15	1.47	2 (15%)	17,20,20	1.25	2 (11%)
4	GOL	D	428	-	5,5,5	0.47	0	5,5,5	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	429	-	-	4/4/4/4	-
3	KYN	A	427	-	-	7/12/12/12	0/1/1/1
4	GOL	D	427	-	-	2/4/4/4	-
2	PMP	D	426	-	-	5/8/8/8	0/1/1/1
3	KYN	A	428	-	-	7/12/12/12	0/1/1/1
4	GOL	C	429	-	-	0/4/4/4	-
4	GOL	A	429	-	-	0/4/4/4	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KYN	C	428	-	-	6/12/12/12	0/1/1/1
4	GOL	D	430	-	-	3/4/4/4	-
2	PMP	B	426	-	-	5/8/8/8	0/1/1/1
2	PMP	C	426	-	-	3/8/8/8	0/1/1/1
2	PMP	A	426	-	-	7/8/8/8	0/1/1/1
3	KYN	C	427	-	-	4/12/12/12	0/1/1/1
4	GOL	D	428	-	-	4/4/4/4	-

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	428	KYN	OXT-C	-5.51	1.12	1.30
3	A	427	KYN	OXT-C	4.90	1.46	1.30
3	C	428	KYN	OXT-C	-4.41	1.16	1.30
3	C	427	KYN	OXT-C	-4.26	1.16	1.30
3	A	427	KYN	CD2-CG	-3.95	1.36	1.41

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	426	PMP	O2P-P-O4P	3.37	115.71	106.73
2	B	426	PMP	C3-C4-C5	3.16	121.75	118.72
3	C	427	KYN	CA-CB-C1	-3.16	109.68	113.70
3	A	427	KYN	CD2-CG-N1	-3.11	118.56	122.67
3	C	428	KYN	CD1-CG-CD2	3.00	121.01	118.10

There are no chirality outliers.

5 of 57 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	426	PMP	C5A-O4P-P-O1P
2	A	426	PMP	C5A-O4P-P-O2P
2	A	426	PMP	C5A-O4P-P-O3P
2	B	426	PMP	C3-C4-C4A-N4A
2	B	426	PMP	C5A-O4P-P-O2P

There are no ring outliers.

13 monomers are involved in 42 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	429	GOL	4	0
3	A	427	KYN	4	0
4	D	427	GOL	4	0
2	D	426	PMP	4	0
4	C	429	GOL	6	0
4	A	429	GOL	6	0
3	C	428	KYN	2	0
4	D	430	GOL	1	0
2	B	426	PMP	2	0
2	C	426	PMP	2	0
2	A	426	PMP	3	0
3	C	427	KYN	3	0
4	D	428	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	425/425 (100%)	0.18	26 (6%) 21 29	18, 29, 52, 78	0
1	B	425/425 (100%)	0.16	22 (5%) 27 37	18, 29, 46, 78	0
1	C	425/425 (100%)	0.05	17 (4%) 38 48	14, 23, 43, 85	0
1	D	425/425 (100%)	-0.07	17 (4%) 38 48	14, 22, 39, 79	0
All	All	1700/1700 (100%)	0.08	82 (4%) 30 40	14, 26, 46, 85	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	17	SER	9.6
1	C	20	ARG	8.4
1	C	21	THR	8.0
1	C	19	ILE	7.5
1	A	17	SER	6.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	KYN	C	427	15/15	0.65	0.28	61,63,69,69	0
3	KYN	C	428	15/15	0.70	0.31	62,65,72,72	0
3	KYN	A	428	15/15	0.75	0.26	62,65,69,69	0
4	GOL	C	429	6/6	0.79	0.24	33,34,36,38	0
3	KYN	A	427	15/15	0.81	0.27	66,69,74,75	0
4	GOL	D	429	6/6	0.83	0.27	35,39,40,41	0
4	GOL	D	430	6/6	0.83	0.41	39,43,44,45	0
4	GOL	D	427	6/6	0.88	0.26	21,32,36,42	0
4	GOL	A	429	6/6	0.89	0.28	26,33,34,34	0
4	GOL	D	428	6/6	0.89	0.21	25,37,42,45	0
2	PMP	B	426	16/16	0.94	0.13	29,36,40,44	0
2	PMP	A	426	16/16	0.95	0.11	28,33,36,39	0
2	PMP	C	426	16/16	0.96	0.11	20,28,34,38	0
2	PMP	D	426	16/16	0.96	0.10	18,32,36,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.