



wwPDB X-ray Structure Validation Summary Report

Oct 11, 2021 – 05:49 AM EDT

PDB ID : 2RJP
Title : Crystal structure of ADAMTS4 with inhibitor bound
Authors : Mosyak, L.; Stahl, M.; Somers, W.
Deposited on : 2007-10-15
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	316	
1	B	316	
1	C	316	
1	D	316	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 8872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADAMTS-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	291	2188	1368	390	406	24	0	0	0
1	B	284	2128	1329	379	396	24	0	0	0
1	C	290	2183	1365	389	405	24	0	0	0
1	D	278	2093	1307	374	388	24	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	362	GLN	GLU	engineered mutation	UNP O75173
A	521	ASP	-	expression tag	UNP O75173
A	522	TYR	-	expression tag	UNP O75173
A	523	LYS	-	expression tag	UNP O75173
A	524	ASP	-	expression tag	UNP O75173
A	525	ASP	-	expression tag	UNP O75173
A	526	ASP	-	expression tag	UNP O75173
A	527	ASP	-	expression tag	UNP O75173
A	528	LYS	-	expression tag	UNP O75173
B	362	GLN	GLU	engineered mutation	UNP O75173
B	521	ASP	-	expression tag	UNP O75173
B	522	TYR	-	expression tag	UNP O75173
B	523	LYS	-	expression tag	UNP O75173
B	524	ASP	-	expression tag	UNP O75173
B	525	ASP	-	expression tag	UNP O75173
B	526	ASP	-	expression tag	UNP O75173
B	527	ASP	-	expression tag	UNP O75173
B	528	LYS	-	expression tag	UNP O75173
C	362	GLN	GLU	engineered mutation	UNP O75173
C	521	ASP	-	expression tag	UNP O75173
C	522	TYR	-	expression tag	UNP O75173

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	523	LYS	-	expression tag	UNP O75173
C	524	ASP	-	expression tag	UNP O75173
C	525	ASP	-	expression tag	UNP O75173
C	526	ASP	-	expression tag	UNP O75173
C	527	ASP	-	expression tag	UNP O75173
C	528	LYS	-	expression tag	UNP O75173
D	362	GLN	GLU	engineered mutation	UNP O75173
D	521	ASP	-	expression tag	UNP O75173
D	522	TYR	-	expression tag	UNP O75173
D	523	LYS	-	expression tag	UNP O75173
D	524	ASP	-	expression tag	UNP O75173
D	525	ASP	-	expression tag	UNP O75173
D	526	ASP	-	expression tag	UNP O75173
D	527	ASP	-	expression tag	UNP O75173
D	528	LYS	-	expression tag	UNP O75173


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Ca 3 3	0	0
3	B	3	Total Ca 3 3	0	0
3	C	3	Total Ca 3 3	0	0
3	D	3	Total Ca 3 3	0	0

- Molecule 4 is N-({4'-[(4-isobutyrylphenoxy)methyl]biphenyl-4-yl}sulfonyl)-D-valine (three-letter code: 886) (formula: C₂₈H₃₁NO₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	Total	C	N	O	S	0	0
			36	28	1	6	1		
4	B	1	Total	C	N	O	S	0	0
			36	28	1	6	1		
4	C	1	Total	C	N	O	S	0	0
			36	28	1	6	1		
4	D	1	Total	C	N	O	S	0	0
			36	28	1	6	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	33	Total	O	0	0
			33	33		
5	B	45	Total	O	0	0
			45	45		
5	C	29	Total	O	0	0
			29	29		
5	D	13	Total	O	0	0
			13	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: ADAMTS-4

Chain A:  83% 9% 8%





- Molecule 1: ADAMTS-4

Chain B:  80% 10% 10%




- Molecule 1: ADAMTS-4

Chain C:  82% 10% 8%



- Molecule 1: ADAMTS-4

Chain D:  79% 9% 12%



M396	L416	P428	P431	K441	P461	GLN	LEU	PRO	PRO	P466	P494	P497	C506	LEU	HIS	MET	ASP	GLN	LEU	GLN	ASP	PHE	ASN	PRO	GLN	ALA	ASP	TYR	LYS	ASP	ASP	ASP	ASP	LYS
------	------	------	------	------	------	-----	-----	-----	-----	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.57Å 82.62Å 99.33Å 90.00° 90.63° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80	Depositor
% Data completeness (in resolution range)	100.0 (50.00-2.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.206 , 0.267	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	8872	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 886, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2249	0.56	0/3068
1	B	0.41	0/2184	0.53	0/2973
1	C	0.42	0/2244	0.54	0/3061
1	D	0.60	4/2148 (0.2%)	0.59	2/2923 (0.1%)
All	All	0.48	4/8825 (0.0%)	0.55	2/12025 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	441	LYS	CD-CE	16.46	1.92	1.51
1	D	280	GLN	CD-OE1	8.17	1.42	1.24
1	D	441	LYS	CE-NZ	6.34	1.65	1.49
1	D	441	LYS	CG-CD	5.10	1.69	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	441	LYS	CD-CE-NZ	-7.90	93.52	111.70
1	D	441	LYS	CG-CD-CE	-7.57	89.18	111.90

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	464	PRO	Peptide
1	C	463	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2188	0	2090	16	0
1	B	2128	0	2035	16	0
1	C	2183	0	2087	18	0
1	D	2093	0	1996	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0
4	A	36	0	30	4	0
4	B	36	0	30	3	0
4	C	36	0	30	3	0
4	D	36	0	30	2	0
5	A	33	0	0	0	0
5	B	45	0	0	1	0
5	C	29	0	0	0	0
5	D	13	0	0	0	0
All	All	8872	0	8328	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:441:LYS:CD	1:D:441:LYS:CE	1.92	1.47
1:D:441:LYS:CE	1:D:441:LYS:CG	2.54	0.85
1:C:296:GLN:HG3	1:C:297:ARG:N	1.99	0.77
1:C:464:PRO:O	1:C:466:PRO:HD3	1.95	0.67
1:A:463:LEU:O	1:A:465:PRO:HD2	1.95	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/316 (91%)	276 (96%)	8 (3%)	3 (1%)	15	44
1	B	276/316 (87%)	265 (96%)	10 (4%)	1 (0%)	34	66
1	C	286/316 (90%)	270 (94%)	14 (5%)	2 (1%)	22	53
1	D	270/316 (85%)	259 (96%)	7 (3%)	4 (2%)	10	33
All	All	1119/1264 (88%)	1070 (96%)	39 (4%)	10 (1%)	17	46

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	465	PRO
1	D	395	MET
1	B	228	ASP
1	C	228	ASP
1	A	228	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/261 (92%)	233 (98%)	6 (2%)	47	80
1	B	232/261 (89%)	229 (99%)	3 (1%)	69	91
1	C	239/261 (92%)	235 (98%)	4 (2%)	60	87
1	D	228/261 (87%)	224 (98%)	4 (2%)	59	86
All	All	938/1044 (90%)	921 (98%)	17 (2%)	59	86

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	264	LEU
1	D	345	CYS
1	B	306	ASP
1	B	481	MET
1	C	277	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	234	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 16 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	886	C	510	2	34,38,38	1.21	2 (5%)	49,54,54	1.66	6 (12%)
4	886	B	510	2	34,38,38	1.27	2 (5%)	49,54,54	1.51	7 (14%)
4	886	A	510	2	34,38,38	1.36	3 (8%)	49,54,54	1.23	6 (12%)
4	886	D	510	2	34,38,38	1.38	2 (5%)	49,54,54	1.12	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	886	C	510	2	-	11/32/36/36	0/3/3/3
4	886	B	510	2	-	10/32/36/36	0/3/3/3
4	886	A	510	2	-	9/32/36/36	0/3/3/3
4	886	D	510	2	-	9/32/36/36	0/3/3/3

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	510	886	S1-N1	4.30	1.69	1.61
4	D	510	886	C4-N1	3.95	1.53	1.47
4	A	510	886	S1-N1	3.80	1.68	1.61
4	B	510	886	S1-N1	3.76	1.68	1.61
4	C	510	886	S1-N1	3.34	1.67	1.61

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	510	886	C3-C2-C4	6.89	117.88	111.24
4	D	510	886	C3-C2-C4	5.28	116.32	111.24
4	B	510	886	C3-C2-C4	4.97	116.03	111.24
4	A	510	886	O6-C25-C22	-4.20	115.23	120.66
4	C	510	886	O6-C25-C22	-4.04	115.44	120.66

There are no chirality outliers.

5 of 39 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	510	886	C1-C2-C4-N1
4	B	510	886	C3-C2-C4-N1
4	C	510	886	C1-C2-C4-C5
4	C	510	886	C1-C2-C4-N1
4	C	510	886	C3-C2-C4-C5


There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	510	886	3	0
4	B	510	886	3	0
4	A	510	886	4	0
4	D	510	886	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.