



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2024 – 12:36 pm BST

PDB ID : 2V28
Title : Apo structure of the cold active phenylalanine hydroxylase from *Colwellia psychrerythraea* 34H
Authors : Leiros, H.-K.S.; Pey, A.L.; Innselset, M.; Moe, E.; Leiros, I.; Steen, I.H.; Martinez, A.
Deposited on : 2007-06-04
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 2% 81% 15% ...
1	B	267	 4% 82% 13% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHENYLALANINE-4-HYDROXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	Total 2236	C 1442	N 361	O 422	S 11	32	12	0
1	B	261	Total 2166	C 1401	N 348	O 407	S 10	27	6	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	365	Total 365	O 365	0	0
3	B	264	Total 264	O 264	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PHENYLALANINE-4-HYDROXYLASE



- Molecule 1: PHENYLALANINE-4-HYDROXYLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.43Å 86.29Å 87.67Å 90.00° 96.64° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 19.83 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.5 (20.00-1.95) 97.4 (19.83-1.95)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.94Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.198 , 0.265 0.198 , 0.264	Depositor DCC
R_{free} test set	2173 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtrriage
Anisotropy	0.095	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 74.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5041	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	2.29	13/2290 (0.6%)	1.16	14/3102 (0.5%)
1	B	1.50	11/2220 (0.5%)	1.26	23/3009 (0.8%)
All	All	1.94	24/4510 (0.5%)	1.21	37/6111 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	LYS	CD-CE	78.90	3.48	1.51
1	A	152	LYS	CE-NZ	-42.75	0.42	1.49
1	B	60	ARG	CD-NE	37.18	2.09	1.46
1	B	84	ALA	C-O	28.77	1.78	1.23
1	A	11	VAL	CB-CG2	-27.95	0.94	1.52
1	B	84	ALA	CA-C	24.03	2.15	1.52
1	A	116	GLN	CD-OE1	22.58	1.73	1.24
1	B	85	LEU	CB-CG	19.39	2.08	1.52
1	B	40	LYS	CE-NZ	-18.93	1.01	1.49
1	A	229	LYS	CE-NZ	18.90	1.96	1.49
1	A	40	LYS	CB-CG	18.42	2.02	1.52
1	A	116	GLN	CG-CD	16.43	1.88	1.51
1	A	11	VAL	CB-CG1	15.52	1.85	1.52
1	B	152	LYS	CD-CE	14.62	1.87	1.51
1	B	84	ALA	C-N	13.38	1.64	1.34
1	B	60	ARG	NE-CZ	12.75	1.49	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	LYS	CE-NZ	11.40	1.77	1.49
1	A	6	LYS	CG-CD	11.16	1.90	1.52
1	B	152	LYS	CG-CD	9.36	1.84	1.52
1	B	65	ASP	CG-OD1	9.05	1.46	1.25
1	B	116	GLN	CG-CD	7.35	1.68	1.51
1	A	85	LEU	CA-CB	6.34	1.68	1.53
1	A	13	ASP	C-N	-5.25	1.22	1.34
1	A	175	LYS	CD-CE	-5.06	1.38	1.51

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	LYS	CD-CE-NZ	-37.38	25.74	111.70
1	B	60	ARG	NE-CZ-NH2	-24.03	108.29	120.30
1	B	60	ARG	NE-CZ-NH1	21.52	131.06	120.30
1	B	40	LYS	CD-CE-NZ	-19.82	66.13	111.70
1	B	84	ALA	CB-CA-C	-17.35	84.08	110.10
1	A	11	VAL	CA-CB-CG2	15.61	134.31	110.90
1	B	60	ARG	CG-CD-NE	-13.62	83.20	111.80
1	B	152	LYS	CG-CD-CE	13.35	151.94	111.90
1	B	116	GLN	CG-CD-NE2	13.26	148.53	116.70
1	B	85	LEU	CA-CB-CG	-13.22	84.89	115.30
1	A	13	ASP	C-N-CA	-12.99	89.23	121.70
1	B	152	LYS	CD-CE-NZ	-12.03	84.03	111.70
1	A	116	GLN	OE1-CD-NE2	-11.20	96.14	121.90
1	B	116	GLN	OE1-CD-NE2	-10.79	97.07	121.90
1	B	85	LEU	CB-CG-CD2	10.49	128.84	111.00
1	B	84	ALA	N-CA-CB	-10.18	95.85	110.10
1	A	229	LYS	CD-CE-NZ	-10.11	88.44	111.70
1	A	152	LYS	CD-CE-NZ	-10.10	88.46	111.70
1	A	116	GLN	CG-CD-OE1	9.81	141.22	121.60
1	B	84	ALA	C-N-CA	-9.29	98.47	121.70
1	A	11	VAL	CG1-CB-CG2	8.95	125.22	110.90
1	A	85	LEU	CA-CB-CG	8.58	135.04	115.30
1	B	84	ALA	CA-C-O	-8.38	102.50	120.10
1	B	40	LYS	CA-CB-CG	8.37	131.81	113.40
1	B	84	ALA	O-C-N	8.32	136.01	122.70
1	B	84	ALA	N-CA-C	-7.79	89.98	111.00
1	A	11	VAL	CA-CB-CG1	-6.96	100.46	110.90
1	A	116	GLN	CB-CG-CD	-6.95	93.53	111.60
1	B	65	ASP	CB-CG-OD1	-6.79	112.19	118.30
1	B	116	GLN	CB-CG-CD	-6.78	93.96	111.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	152	LYS	CB-CG-CD	-6.32	95.17	111.60
1	B	110	GLU	CA-CB-CG	-5.74	100.76	113.40
1	A	40	LYS	CA-CB-CG	-5.71	100.84	113.40
1	A	14	GLU	CB-CA-C	5.59	121.57	110.40
1	B	40	LYS	CB-CG-CD	5.58	126.11	111.60
1	B	117	GLU	CA-CB-CG	-5.45	101.42	113.40
1	A	40	LYS	CB-CG-CD	5.30	125.37	111.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	65	ASP	Sidechain
1	B	84	ALA	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2236	0	2192	34	0
1	B	2166	0	2125	18	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	365	0	0	8	0
3	B	264	0	0	2	0
All	All	5041	0	4317	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:LYS:NZ	1:A:6:LYS:CE	1.77	1.45
1:B:201[A]:ASP:OD2	1:B:260:LYS:NZ	2.06	0.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:ASP:HB2	1:A:17:PHE:H	1.35	0.88
1:A:49:GLU:HG2	3:A:2095:HOH:O	1.78	0.84
1:A:33[A]:THR:HG23	3:A:2119:HOH:O	1.80	0.81
1:B:138:ASN:HB3	1:B:237:ARG:HD2	1.76	0.67
1:A:30:GLU:HG3	3:A:2061:HOH:O	1.96	0.66
1:A:47:TYR:CE1	1:A:185[A]:LEU:HD13	2.34	0.63
1:A:13:ASP:HB3	1:A:15:HIS:H	1.65	0.62
1:A:155:ARG:HD3	3:A:2227:HOH:O	1.98	0.61
1:B:127:HIS:CE1	1:B:182:GLY:HA3	2.36	0.61
1:A:38[B]:CYS:O	1:A:42:LYS:HD2	2.02	0.60
1:A:13:ASP:HB3	1:A:15:HIS:N	2.17	0.59
1:A:118:PRO:HB3	1:A:122:HIS:CD2	2.37	0.59
1:A:13:ASP:O	1:A:14:GLU:C	2.26	0.58
1:A:179:ILE:HB	1:A:185[B]:LEU:HD21	1.87	0.56
1:B:133:ASN:HD22	1:B:136:PHE:H	1.54	0.55
1:A:69:LYS:HE3	3:A:2139:HOH:O	2.06	0.55
1:A:13:ASP:HB2	1:A:17:PHE:N	2.15	0.55
1:B:223:ILE:HD11	1:B:260:LYS:HE3	1.90	0.52
1:B:35:GLN:HE22	1:B:188:PRO:HD3	1.73	0.52
1:A:133:ASN:HD22	1:A:136:PHE:H	1.56	0.52
1:A:138:ASN:HB3	1:A:237[A]:ARG:HD2	1.94	0.50
1:A:216:ARG:HG3	1:A:266:ALA:HB3	1.92	0.50
1:A:69:LYS:HE2	3:A:2135:HOH:O	2.11	0.50
1:A:118:PRO:HG2	3:A:2204:HOH:O	2.12	0.49
1:A:12:PRO:HB2	1:A:16:GLY:HA2	1.95	0.49
1:A:242:ASP:O	1:A:246[A]:GLU:HG3	2.13	0.48
1:B:7:TYR:N	3:B:2001:HOH:O	2.48	0.47
1:A:112:MET:HE3	1:A:112:MET:HB2	1.64	0.46
1:B:72:LYS:HE3	1:B:78:GLU:HB3	1.97	0.46
1:B:34:ARG:HB3	3:B:2035:HOH:O	2.16	0.46
1:B:35:GLN:HG2	1:B:125:PHE:CE2	2.51	0.45
1:A:11:VAL:HA	1:A:12:PRO:HD3	1.86	0.44
1:B:93:ARG:O	1:B:97:GLU:HG3	2.18	0.44
1:A:179:ILE:HG22	1:A:185[B]:LEU:HG	2.00	0.44
1:B:30:GLU:CD	1:B:109:ARG:HH22	2.22	0.44
1:A:92:PHE:CD2	1:A:147:GLY:HA3	2.53	0.43
1:A:127:HIS:HE1	1:A:167:GLU:OE1	2.01	0.43
1:A:237[B]:ARG:NH2	3:A:2323:HOH:O	2.51	0.43
1:A:91:PHE:CZ	1:A:95:LEU:HD11	2.55	0.41
1:B:36:ILE:HD13	1:B:36:ILE:HA	1.85	0.41
1:B:162:TYR:CE2	1:B:167:GLU:HG2	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:GLN:HE22	1:A:188:PRO:HD3	1.85	0.41
1:B:206:ASP:HB3	1:B:209:ASP:HB2	2.03	0.41
1:B:200[B]:VAL:HG12	1:B:201[B]:ASP:N	2.35	0.41
1:B:29:GLN:O	1:B:33:THR:HG23	2.21	0.41
1:A:42:LYS:HD3	1:A:195:MET:HG3	2.03	0.41
1:B:100:PHE:HA	1:B:101:PRO:HD3	1.95	0.41
1:A:142:ALA:O	1:A:146:MET:HG3	2.21	0.40
1:A:14:GLU:C	1:A:16:GLY:H	2.24	0.40
1:A:179:ILE:CG2	1:A:185[B]:LEU:HG	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/267 (103%)	267 (97%)	7 (3%)	0	100	100
1	B	265/267 (99%)	258 (97%)	6 (2%)	1 (0%)	34	22
All	All	539/534 (101%)	525 (97%)	13 (2%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	85	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/235 (104%)	238 (97%)	7 (3%)	42	31
1	B	237/235 (101%)	232 (98%)	5 (2%)	53	46
All	All	482/470 (103%)	470 (98%)	12 (2%)	46	38

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LYS
1	A	13	ASP
1	A	85	LEU
1	A	112	MET
1	A	133	ASN
1	A	229	LYS
1	A	267	SER
1	B	36	ILE
1	B	40	LYS
1	B	41	ASP
1	B	152	LYS
1	B	235	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	133	ASN
1	B	35	GLN
1	B	133	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	1268	-	4,4,4	0.18	0	6,6,6	0.30	0
2	SO4	B	1268	-	4,4,4	0.15	0	6,6,6	0.11	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	84:ALA	C	85:LEU	N	1.64

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/267 (98%)	-0.16	5 (1%) 66 74	7, 14, 32, 50	22 (8%)
1	B	261/267 (97%)	0.36	11 (4%) 36 45	12, 26, 47, 67	18 (6%)
All	All	525/534 (98%)	0.10	16 (3%) 50 59	7, 20, 45, 67	40 (7%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	8	VAL	10.1
1	A	4	GLY	6.4
1	B	7	TYR	5.9
1	A	5	THR	4.6
1	B	15	HIS	4.4
1	A	15	HIS	4.2
1	B	267	SER	4.2
1	A	14	GLU	4.1
1	B	83	PRO	3.9
1	B	85	LEU	3.8
1	B	86	ILE	3.6
1	B	88	PHE	3.4
1	B	22	THR	3.3
1	B	9	SER	3.2
1	B	84	ALA	3.2
1	A	85	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	B	1268	5/5	0.86	0.19	70,70,71,71	0
2	SO4	A	1268	5/5	0.92	0.19	43,43,44,45	5

6.5 Other polymers [i](#)

There are no such residues in this entry.