



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 9, 2023 – 09:22 am GMT

PDB ID : 2VUS
Title : Crystal structure of unliganded NmrA-AreA zinc finger complex
Authors : Kotaka, M.; Johnson, C.; Lamb, H.K.; Hawkins, A.R.; Ren, J.; Stammers, D.K.
Deposited on : 2008-05-30
Resolution : 2.60 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	
1	B	352	
1	C	352	
1	D	352	
1	E	352	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	352	<p>%</p> <p>59% 31% 10%</p>
1	G	352	<p>59% 30% 10%</p>
1	H	352	<p>59% 30% 10%</p>
2	I	43	<p>7%</p> <p>63% 35%</p>
2	J	43	<p>2%</p> <p>56% 40%</p>
2	K	43	<p>5%</p> <p>51% 40% 5% 5%</p>
2	L	43	<p>56% 42%</p>
2	M	43	<p>9%</p> <p>67% 30%</p>
2	N	43	<p>5%</p> <p>63% 35%</p>
2	O	43	<p>7%</p> <p>63% 30% 5%</p>
2	P	43	<p>9%</p> <p>63% 33%</p>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 24506 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITROGEN METABOLITE REPRESSION REGULATOR NMRA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	318	2530	1637	428	457	8	0	0	0
1	B	318	2530	1637	428	457	8	0	0	0
1	C	318	2530	1637	428	457	8	0	0	0
1	D	318	2530	1637	428	457	8	0	0	0
1	E	318	2530	1637	428	457	8	0	0	0
1	F	318	2530	1637	428	457	8	0	0	0
1	G	318	2530	1637	428	457	8	0	0	0
1	H	318	2530	1637	428	457	8	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	238	ARG	LEU	conflict	UNP O59919
B	238	ARG	LEU	conflict	UNP O59919
C	238	ARG	LEU	conflict	UNP O59919
D	238	ARG	LEU	conflict	UNP O59919
E	238	ARG	LEU	conflict	UNP O59919
F	238	ARG	LEU	conflict	UNP O59919
G	238	ARG	LEU	conflict	UNP O59919
H	238	ARG	LEU	conflict	UNP O59919

- Molecule 2 is a protein called NITROGEN REGULATORY PROTEIN AREA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	J	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	K	41	Total	C	N	O	S	0	0	0
			318	200	59	55	4			
2	L	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	M	43	Total	C	N	O	S	0	0	0
			333	211	61	57	4			
2	N	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	O	41	Total	C	N	O	S	0	0	0
			318	200	59	55	4			
2	P	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O S	0	0
			5	4 1		
3	B	1	Total	O S	0	0
			5	4 1		
3	C	1	Total	O S	0	0
			5	4 1		
3	D	1	Total	O S	0	0
			5	4 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		
4	F	1	Total	Cl	0	0
			1	1		
4	G	2	Total	Cl	0	0
			2	2		
4	H	1	Total	Cl	0	0
			1	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	I	1	Total	Zn	0	0
			1	1		
5	J	1	Total	Zn	0	0
			1	1		
5	K	1	Total	Zn	0	0
			1	1		
5	L	1	Total	Zn	0	0
			1	1		
5	M	1	Total	Zn	0	0
			1	1		
5	N	1	Total	Zn	0	0
			1	1		
5	O	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	P	1	Total 1	Zn 1	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	148	Total 148	O 148	0	0
6	B	181	Total 181	O 181	0	0
6	C	192	Total 192	O 192	0	0
6	D	184	Total 184	O 184	0	0
6	E	207	Total 207	O 207	0	0
6	F	170	Total 170	O 170	0	0
6	G	212	Total 212	O 212	0	0
6	H	177	Total 177	O 177	0	0
6	I	26	Total 26	O 26	0	0
6	J	17	Total 17	O 17	0	0
6	K	20	Total 20	O 20	0	0
6	L	16	Total 16	O 16	0	0
6	M	17	Total 17	O 17	0	0
6	N	17	Total 17	O 17	0	0
6	O	17	Total 17	O 17	0	0
6	P	11	Total 11	O 11	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA

Chain A: 




- Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA

Chain B: 




- Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA


Chain C: 




- Molecule 2: NITROGEN REGULATORY PROTEIN AREA




- Molecule 2: NITROGEN REGULATORY PROTEIN AREA




- Molecule 2: NITROGEN REGULATORY PROTEIN AREA




- Molecule 2: NITROGEN REGULATORY PROTEIN AREA



- Molecule 2: NITROGEN REGULATORY PROTEIN AREA



- Molecule 2: NITROGEN REGULATORY PROTEIN AREA



- Molecule 2: NITROGEN REGULATORY PROTEIN AREA



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	228.79Å 228.79Å 222.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.97 – 2.60 29.97 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.7 (29.97-2.60) 98.8 (29.97-2.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.61Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.242 , 0.306 0.243 , 0.306	Depositor DCC
R_{free} test set	6691 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	25.8	Xtrriage
Anisotropy	0.356	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 18.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.387 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24506	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/2606	0.64	0/3557
1	B	0.38	0/2606	0.67	0/3557
1	C	0.39	0/2606	0.64	0/3557
1	D	0.39	0/2606	0.65	0/3557
1	E	0.39	0/2606	0.66	0/3557
1	F	0.38	0/2606	0.63	0/3557
1	G	0.40	0/2606	0.66	1/3557 (0.0%)
1	H	0.38	0/2606	0.64	0/3557
2	I	0.39	0/334	0.61	0/456
2	J	0.38	0/334	0.65	0/456
2	K	0.38	0/326	0.66	0/445
2	L	0.35	0/334	0.64	0/456
2	M	0.37	0/342	0.58	0/467
2	N	0.39	0/334	0.63	0/456
2	O	0.41	0/326	0.60	0/445
2	P	0.38	0/334	0.64	0/456
All	All	0.39	0/23512	0.65	1/32093 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	167	MET	N-CA-C	-5.03	97.42	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2530	0	2476	127	0
1	B	2530	0	2476	106	0
1	C	2530	0	2476	103	0
1	D	2530	0	2476	123	0
1	E	2530	0	2476	89	0
1	F	2530	0	2476	103	0
1	G	2530	0	2476	92	0
1	H	2530	0	2476	88	0
2	I	326	0	326	11	0
2	J	326	0	326	18	0
2	K	318	0	315	25	0
2	L	326	0	326	16	0
2	M	333	0	334	11	0
2	N	326	0	326	15	0
2	O	318	0	315	13	0
2	P	326	0	326	12	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
3	F	5	0	0	0	0
3	G	5	0	0	0	0
3	H	5	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	F	1	0	0	0	0
4	G	2	0	0	0	0
4	H	1	0	0	0	0
5	I	1	0	0	0	0
5	J	1	0	0	0	0
5	K	1	0	0	0	0
5	L	1	0	0	0	0
5	M	1	0	0	0	0
5	N	1	0	0	0	0
5	O	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	P	1	0	0	0	0
6	A	148	0	0	11	0
6	B	181	0	0	17	0
6	C	192	0	0	13	0
6	D	184	0	0	18	0
6	E	207	0	0	10	0
6	F	170	0	0	9	0
6	G	212	0	0	18	0
6	H	177	0	0	8	0
6	I	26	0	0	0	0
6	J	17	0	0	3	0
6	K	20	0	0	0	0
6	L	16	0	0	3	0
6	M	17	0	0	0	0
6	N	17	0	0	3	0
6	O	17	0	0	0	0
6	P	11	0	0	0	0
All	All	24506	0	22402	924	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 924 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:687:ASN:HD21	2:J:691:GLN:HB2	1.14	1.06
1:G:352:LEU:OXT	6:G:2212:HOH:O	1.72	1.06
2:N:687:ASN:HD22	2:N:691:GLN:HB2	1.17	1.05
1:A:261:GLN:HE21	1:A:265:ILE:HD11	1.21	1.04
2:O:687:ASN:HD21	2:O:691:GLN:HB2	1.19	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/352 (89%)	291 (93%)	20 (6%)	3 (1%)	15	32
1	B	314/352 (89%)	287 (91%)	24 (8%)	3 (1%)	15	32
1	C	314/352 (89%)	290 (92%)	22 (7%)	2 (1%)	25	47
1	D	314/352 (89%)	292 (93%)	19 (6%)	3 (1%)	15	32
1	E	314/352 (89%)	285 (91%)	27 (9%)	2 (1%)	25	47
1	F	314/352 (89%)	280 (89%)	33 (10%)	1 (0%)	41	64
1	G	314/352 (89%)	294 (94%)	18 (6%)	2 (1%)	25	47
1	H	314/352 (89%)	288 (92%)	21 (7%)	5 (2%)	9	19
2	I	40/43 (93%)	37 (92%)	2 (5%)	1 (2%)	5	9
2	J	40/43 (93%)	36 (90%)	3 (8%)	1 (2%)	5	9
2	K	39/43 (91%)	35 (90%)	3 (8%)	1 (3%)	5	9
2	L	40/43 (93%)	31 (78%)	8 (20%)	1 (2%)	5	9
2	M	41/43 (95%)	38 (93%)	3 (7%)	0	100	100
2	N	40/43 (93%)	37 (92%)	3 (8%)	0	100	100
2	O	39/43 (91%)	37 (95%)	1 (3%)	1 (3%)	5	9
2	P	40/43 (93%)	38 (95%)	1 (2%)	1 (2%)	5	9
All	All	2831/3160 (90%)	2596 (92%)	208 (7%)	27 (1%)	15	32

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	350	TRP
1	G	116	HIS
1	A	253	ASN
1	B	319	ASP
1	F	350	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/284 (94%)	259 (97%)	7 (3%)	46	72
1	B	266/284 (94%)	261 (98%)	5 (2%)	57	79
1	C	266/284 (94%)	263 (99%)	3 (1%)	73	88
1	D	266/284 (94%)	257 (97%)	9 (3%)	37	63
1	E	266/284 (94%)	260 (98%)	6 (2%)	50	75
1	F	266/284 (94%)	261 (98%)	5 (2%)	57	79
1	G	266/284 (94%)	262 (98%)	4 (2%)	65	83
1	H	266/284 (94%)	259 (97%)	7 (3%)	46	72
2	I	38/39 (97%)	38 (100%)	0	100	100
2	J	38/39 (97%)	37 (97%)	1 (3%)	46	72
2	K	37/39 (95%)	36 (97%)	1 (3%)	44	71
2	L	38/39 (97%)	38 (100%)	0	100	100
2	M	39/39 (100%)	38 (97%)	1 (3%)	46	72
2	N	38/39 (97%)	38 (100%)	0	100	100
2	O	37/39 (95%)	34 (92%)	3 (8%)	11	23
2	P	38/39 (97%)	38 (100%)	0	100	100
All	All	2431/2584 (94%)	2379 (98%)	52 (2%)	53	77

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	212	ASN
1	G	28	VAL
2	O	671	THR
1	E	351	MET
1	F	155	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 99 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	84	GLN
1	G	84	GLN
1	F	136	ASN
1	F	244	GLN
1	G	166	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 15 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	E	1353	-	4,4,4	0.29	0	6,6,6	0.12	0
3	SO4	C	1353	-	4,4,4	0.34	0	6,6,6	0.08	0
3	SO4	H	1353	-	4,4,4	0.26	0	6,6,6	0.21	0
3	SO4	B	1353	-	4,4,4	0.30	0	6,6,6	0.17	0
3	SO4	D	1353	-	4,4,4	0.32	0	6,6,6	0.21	0
3	SO4	G	1353	-	4,4,4	0.31	0	6,6,6	0.18	0
3	SO4	F	1353	-	4,4,4	0.26	0	6,6,6	0.22	0
3	SO4	A	1353	-	4,4,4	0.39	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/352 (90%)	-0.54	0 100 100	8, 28, 61, 91	0
1	B	318/352 (90%)	-0.53	0 100 100	9, 26, 53, 73	0
1	C	318/352 (90%)	-0.68	1 (0%) 94 93	10, 22, 45, 69	0
1	D	318/352 (90%)	-0.63	0 100 100	10, 26, 57, 72	0
1	E	318/352 (90%)	-0.57	0 100 100	8, 24, 52, 75	0
1	F	318/352 (90%)	-0.51	2 (0%) 89 88	9, 25, 55, 70	0
1	G	318/352 (90%)	-0.67	0 100 100	5, 23, 50, 75	0
1	H	318/352 (90%)	-0.63	0 100 100	7, 24, 59, 82	0
2	I	42/43 (97%)	-0.03	3 (7%) 16 11	8, 33, 92, 111	0
2	J	42/43 (97%)	-0.14	1 (2%) 59 53	14, 37, 89, 101	0
2	K	41/43 (95%)	-0.21	2 (4%) 29 23	12, 37, 100, 104	0
2	L	42/43 (97%)	-0.12	0 100 100	16, 41, 96, 119	0
2	M	43/43 (100%)	0.17	4 (9%) 8 6	13, 35, 101, 116	0
2	N	42/43 (97%)	-0.11	2 (4%) 30 24	9, 31, 90, 110	0
2	O	41/43 (95%)	-0.01	3 (7%) 15 11	20, 34, 89, 106	0
2	P	42/43 (97%)	0.03	4 (9%) 8 5	14, 39, 99, 107	0
All	All	2879/3160 (91%)	-0.53	22 (0%) 86 84	5, 26, 59, 119	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	688	PRO	6.9
2	O	688	PRO	5.2
2	I	688	PRO	5.1
2	M	670	PRO	4.3
2	P	686	ARG	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	B	1353	5/5	0.92	0.17	19,46,53,63	0
4	CL	A	1354	1/1	0.92	0.23	58,58,58,58	0
4	CL	F	1354	1/1	0.93	0.12	71,71,71,71	0
3	SO4	A	1353	5/5	0.94	0.13	30,31,52,71	0
3	SO4	F	1353	5/5	0.94	0.15	20,31,46,51	0
5	ZN	M	1713	1/1	0.95	0.11	43,43,43,43	0
4	CL	D	1354	1/1	0.96	0.11	56,56,56,56	0
3	SO4	C	1353	5/5	0.96	0.12	25,29,58,59	0
4	CL	G	1354	1/1	0.96	0.10	37,37,37,37	0
3	SO4	E	1353	5/5	0.96	0.12	35,37,54,58	0
5	ZN	O	1712	1/1	0.96	0.19	57,57,57,57	0
3	SO4	G	1353	5/5	0.97	0.10	24,34,46,62	0
3	SO4	H	1353	5/5	0.97	0.12	38,51,57,63	0
3	SO4	D	1353	5/5	0.98	0.10	18,23,47,55	0
4	CL	H	1354	1/1	0.98	0.09	44,44,44,44	0
5	ZN	I	1713	1/1	0.99	0.11	38,38,38,38	0
5	ZN	J	1713	1/1	0.99	0.10	40,40,40,40	0
5	ZN	K	1712	1/1	0.99	0.12	30,30,30,30	0
5	ZN	L	1713	1/1	0.99	0.10	36,36,36,36	0
4	CL	G	1355	1/1	0.99	0.05	41,41,41,41	0
5	ZN	N	1713	1/1	0.99	0.11	34,34,34,34	0
4	CL	C	1354	1/1	0.99	0.12	49,49,49,49	0
5	ZN	P	1713	1/1	0.99	0.09	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.