



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2023 – 07:55 PM EDT

PDB ID : 2Z6K  
Title : Crystal structure of full-length human RPA14/32 heterodimer  
Authors : Deng, X.; Habel, J.E.; Kabaleswaran, V.; Borgstahl, G.E.  
Deposited on : 2007-08-03  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	
1	B	270	
2	C	142	
2	D	142	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3818 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Replication protein A 32 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	127	998	635	173	184	6	0	0	0
1	B	126	990	631	171	182	6	0	0	0

- Molecule 2 is a protein called Replication protein A 14 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	117	915	590	148	170	7	0	0	0
2	D	117	915	590	148	170	7	0	0	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	-20	MET	-	expression tag	UNP P35244
C	-19	GLY	-	expression tag	UNP P35244
C	-18	HIS	-	expression tag	UNP P35244
C	-17	HIS	-	expression tag	UNP P35244
C	-16	HIS	-	expression tag	UNP P35244
C	-15	HIS	-	expression tag	UNP P35244
C	-14	HIS	-	expression tag	UNP P35244
C	-13	HIS	-	expression tag	UNP P35244
C	-12	HIS	-	expression tag	UNP P35244
C	-11	HIS	-	expression tag	UNP P35244
C	-10	HIS	-	expression tag	UNP P35244
C	-9	HIS	-	expression tag	UNP P35244
C	-8	SER	-	expression tag	UNP P35244
C	-7	SER	-	expression tag	UNP P35244
C	-6	GLY	-	expression tag	UNP P35244
C	-5	HIS	-	expression tag	UNP P35244

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	ILE	-	expression tag	UNP P35244
C	-3	GLU	-	expression tag	UNP P35244
C	-2	GLY	-	expression tag	UNP P35244
C	-1	ARG	-	expression tag	UNP P35244
C	0	HIS	-	expression tag	UNP P35244
D	-20	MET	-	expression tag	UNP P35244
D	-19	GLY	-	expression tag	UNP P35244
D	-18	HIS	-	expression tag	UNP P35244
D	-17	HIS	-	expression tag	UNP P35244
D	-16	HIS	-	expression tag	UNP P35244
D	-15	HIS	-	expression tag	UNP P35244
D	-14	HIS	-	expression tag	UNP P35244
D	-13	HIS	-	expression tag	UNP P35244
D	-12	HIS	-	expression tag	UNP P35244
D	-11	HIS	-	expression tag	UNP P35244
D	-10	HIS	-	expression tag	UNP P35244
D	-9	HIS	-	expression tag	UNP P35244
D	-8	SER	-	expression tag	UNP P35244
D	-7	SER	-	expression tag	UNP P35244
D	-6	GLY	-	expression tag	UNP P35244
D	-5	HIS	-	expression tag	UNP P35244
D	-4	ILE	-	expression tag	UNP P35244
D	-3	GLU	-	expression tag	UNP P35244
D	-2	GLY	-	expression tag	UNP P35244
D	-1	ARG	-	expression tag	UNP P35244
D	0	HIS	-	expression tag	UNP P35244





- Molecule 2: Replication protein A 14 kDa subunit

Chain D: 53% 27% 18%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.46Å 97.46Å 125.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00 31.76 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-3.00) 99.9 (31.76-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.76 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.227 , 0.269 0.232 , 0.267	Depositor DCC
$R_{free}$ test set	1471 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.4	Xtrriage
Anisotropy	0.115	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 56.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.045 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3818	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/1015	0.72	0/1379
1	B	0.52	0/1007	0.64	0/1368
2	C	0.57	0/934	0.67	0/1262
2	D	0.48	0/934	0.68	0/1262
All	All	0.55	0/3890	0.68	0/5271

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	4	1
1	B	1	0
2	D	0	1
All	All	5	2

There are no bond length outliers.

There are no bond angle outliers.

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	88	THR	CB
1	A	97	MET	CA
1	A	156	THR	CB
1	A	175	GLN	CA
1	B	97	MET	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	97	MET	Peptide

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Group
2	D	59	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	998	0	1022	35	0
1	B	990	0	1016	37	0
2	C	915	0	921	24	0
2	D	915	0	921	29	0
All	All	3818	0	3880	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:4:MET:HE1	2:D:82:THR:HG21	1.26	1.12
1:A:88:THR:HG23	1:A:89:ASN:N	1.75	1.01
1:A:61:ASP:OD1	1:A:62:GLU:OE1	1.77	1.01
1:A:88:THR:HG23	1:A:89:ASN:H	1.26	0.99
1:B:86:ALA:HB3	1:B:89:ASN:OD1	1.63	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	123/270 (46%)	109 (89%)	12 (10%)	2 (2%)	9	40
1	B	122/270 (45%)	103 (84%)	16 (13%)	3 (2%)	5	28
2	C	115/142 (81%)	109 (95%)	5 (4%)	1 (1%)	17	55
2	D	115/142 (81%)	105 (91%)	7 (6%)	3 (3%)	5	27
All	All	475/824 (58%)	426 (90%)	40 (8%)	9 (2%)	8	36

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	68	ASN
1	B	87	PRO
2	C	3	ASP
1	A	123	GLU
1	A	152	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/228 (50%)	107 (94%)	7 (6%)	18	53
1	B	113/228 (50%)	109 (96%)	4 (4%)	36	71
2	C	102/124 (82%)	100 (98%)	2 (2%)	55	83
2	D	102/124 (82%)	96 (94%)	6 (6%)	19	54
All	All	431/704 (61%)	412 (96%)	19 (4%)	28	65

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	59	LEU
2	D	84	TYR
2	D	90	ASP
2	D	76	LYS
1	B	97	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	158	HIS
1	B	173	ASN
2	D	86	GLN
2	C	35	HIS
1	A	173	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	127/270 (47%)	-0.23	3 (2%) 59 30	60, 79, 99, 110	1 (0%)
1	B	126/270 (46%)	0.25	11 (8%) 10 3	75, 108, 131, 138	1 (0%)
2	C	117/142 (82%)	-0.55	0 100 100	65, 75, 94, 101	0
2	D	117/142 (82%)	-0.23	0 100 100	72, 90, 119, 123	0
All	All	487/824 (59%)	-0.18	14 (2%) 51 23	60, 84, 122, 138	2 (0%)

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	117	ASN	4.7
1	B	42	ARG	4.3
1	B	87	PRO	3.5
1	B	85	LYS	3.5
1	B	88	THR	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.