



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 06:29 pm BST

PDB ID : 3B4P
Title : Crystal structure of phenazine biosynthesis protein PhzA/B from Burkholderia cepacia R18194, complex with 2-(cyclohexylamino)benzoic acid
Authors : Ahuja, E.G.; Janning, P.; Mentel, M.; Graebisch, A.; Breinbauer, R.; Blankenfeldt, W.
Deposited on : 2007-10-24
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	185	 71% 14% • 15%
1	B	185	 69% 14% • 15%

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 2954 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenazine biosynthesis protein A/B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	157	1346	846	253	242	5	0	5	0
1	B	158	1320	832	246	237	5	0	1	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q396C9
A	-18	GLY	-	expression tag	UNP Q396C9
A	-17	SER	-	expression tag	UNP Q396C9
A	-16	SER	-	expression tag	UNP Q396C9
A	-15	HIS	-	expression tag	UNP Q396C9
A	-14	HIS	-	expression tag	UNP Q396C9
A	-13	HIS	-	expression tag	UNP Q396C9
A	-12	HIS	-	expression tag	UNP Q396C9
A	-11	HIS	-	expression tag	UNP Q396C9
A	-10	HIS	-	expression tag	UNP Q396C9
A	-9	SER	-	expression tag	UNP Q396C9
A	-8	SER	-	expression tag	UNP Q396C9
A	-7	GLY	-	expression tag	UNP Q396C9
A	-6	LEU	-	expression tag	UNP Q396C9
A	-5	VAL	-	expression tag	UNP Q396C9
A	-4	PRO	-	expression tag	UNP Q396C9
A	-3	ARG	-	expression tag	UNP Q396C9
A	-2	GLY	-	expression tag	UNP Q396C9
A	-1	SER	-	expression tag	UNP Q396C9
A	0	HIS	-	expression tag	UNP Q396C9
B	-19	MET	-	expression tag	UNP Q396C9
B	-18	GLY	-	expression tag	UNP Q396C9
B	-17	SER	-	expression tag	UNP Q396C9
B	-16	SER	-	expression tag	UNP Q396C9
B	-15	HIS	-	expression tag	UNP Q396C9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP Q396C9
B	-13	HIS	-	expression tag	UNP Q396C9
B	-12	HIS	-	expression tag	UNP Q396C9
B	-11	HIS	-	expression tag	UNP Q396C9
B	-10	HIS	-	expression tag	UNP Q396C9
B	-9	SER	-	expression tag	UNP Q396C9
B	-8	SER	-	expression tag	UNP Q396C9
B	-7	GLY	-	expression tag	UNP Q396C9
B	-6	LEU	-	expression tag	UNP Q396C9
B	-5	VAL	-	expression tag	UNP Q396C9
B	-4	PRO	-	expression tag	UNP Q396C9
B	-3	ARG	-	expression tag	UNP Q396C9
B	-2	GLY	-	expression tag	UNP Q396C9
B	-1	SER	-	expression tag	UNP Q396C9
B	0	HIS	-	expression tag	UNP Q396C9

- Molecule 2 is 2-(cyclohexylamino)benzoic acid (three-letter code: 3B4) (formula: C₁₃H₁₇NO₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
2	A	1	16	13	1	2	0	0
2	B	1	16	13	1	2	0	0

- Molecule 3 is AZIDE ION (three-letter code: AZI) (formula: N₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total N 3 3	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	111	Total 111	O 111	0	0
5	B	138	Total 138	O 138	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phenazine biosynthesis protein A/B

Chain A: 



- Molecule 1: Phenazine biosynthesis protein A/B

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	64.55Å 64.55Å 160.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.65 – 1.70 19.65 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.65-1.70) 100.0 (19.65-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.41 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.170 , 0.214 0.245 , 0.285	Depositor DCC
R_{free} test set	2191 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.043 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2954	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AZI, 3B4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.27	6/1388 (0.4%)	1.15	6/1875 (0.3%)
1	B	1.27	4/1361 (0.3%)	1.18	9/1839 (0.5%)
All	All	1.27	10/2749 (0.4%)	1.16	15/3714 (0.4%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	137	GLU	CD-OE2	6.78	1.33	1.25
1	B	68	GLU	CG-CD	5.96	1.60	1.51
1	B	84	TRP	CZ3-CH2	5.87	1.49	1.40
1	B	107	GLU	CB-CG	-5.71	1.41	1.52
1	A	140	GLU	CD-OE2	-5.65	1.19	1.25
1	B	72	GLU	CG-CD	5.62	1.60	1.51
1	A	107	GLU	CB-CG	-5.59	1.41	1.52
1	A	101	TRP	CE3-CZ3	5.29	1.47	1.38
1	A	120	TYR	CD1-CE1	5.05	1.47	1.39
1	A	92	PHE	CE1-CZ	-5.05	1.27	1.37

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	34	ARG	NE-CZ-NH2	-11.18	114.71	120.30
1	B	65	ARG	NE-CZ-NH1	11.03	125.81	120.30
1	B	149	ARG	NE-CZ-NH2	-7.37	116.61	120.30
1	B	65	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	B	34	ARG	NE-CZ-NH1	-6.03	117.28	120.30
1	A	88[A]	ASP	CB-CG-OD1	5.93	123.64	118.30
1	A	88[B]	ASP	CB-CG-OD1	5.93	123.64	118.30
1	B	24	ARG	NE-CZ-NH2	-5.57	117.52	120.30
1	A	14	ARG	NE-CZ-NH1	5.54	123.07	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	121	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	B	96	ASP	CB-CG-OD1	5.38	123.14	118.30
1	B	139	ARG	NE-CZ-NH1	-5.34	117.63	120.30
1	A	34	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	B	159	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	A	38	ARG	NE-CZ-NH2	-5.21	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1346	0	1269	16	0
1	B	1320	0	1252	23	0
2	A	16	0	16	0	0
2	B	16	0	16	2	0
3	A	3	0	0	0	0
4	B	4	0	3	1	0
5	A	111	0	0	9	0
5	B	138	0	0	10	0
All	All	2954	0	2556	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:ASP:OD2	1:B:136:LYS:NZ	1.87	1.07
1:A:149:ARG:NH1	5:A:608:HOH:O	1.85	1.06
1:A:117:ARG:HD3	5:A:707:HOH:O	1.79	0.82
1:B:41:ARG:NH1	5:B:714:HOH:O	2.12	0.82
1:B:48:ASP:CG	1:B:136:LYS:NZ	2.40	0.75
1:B:48:ASP:CG	1:B:136:LYS:HZ1	1.90	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:34:ARG:NH1	5:B:713:HOH:O	2.14	0.70
1:B:60:GLN:NE2	5:B:712:HOH:O	2.10	0.59
1:B:25:LYS:HD2	5:B:653:HOH:O	2.02	0.58
1:B:29:GLN:OE1	5:B:650:HOH:O	2.17	0.57
1:B:136:LYS:HE3	5:B:682:HOH:O	2.05	0.56
1:B:107:GLU:OE1	1:B:121:ARG:HD3	2.06	0.55
1:B:81:PHE:CE1	2:B:500:3B4:HAJA	2.41	0.54
1:B:84:TRP:CD1	4:B:600:ACT:H3	2.42	0.54
1:A:25:LYS:CD	5:A:708:HOH:O	2.56	0.53
1:B:122:ASN:HD22	1:B:123:HIS:H	1.56	0.52
1:A:122:ASN:HD22	1:A:123[B]:HIS:H	1.57	0.51
1:A:163:LEU:HD12	1:A:164:PRO:HD2	1.93	0.50
1:A:149:ARG:NE	5:A:665:HOH:O	2.43	0.50
1:A:25:LYS:CE	5:A:704:HOH:O	2.60	0.49
1:A:125:LEU:HD12	1:B:125:LEU:HD12	1.95	0.48
1:A:28:GLU:OE1	5:A:708:HOH:O	2.19	0.48
1:A:143:ASN:N	5:A:705:HOH:O	2.48	0.46
1:A:149:ARG:NH2	5:A:665:HOH:O	2.49	0.46
1:B:90[B]:GLN:HG2	1:B:92:PHE:CZ	2.52	0.44
1:B:132:ASN:HA	5:B:735:HOH:O	2.17	0.44
1:A:122:ASN:HD22	1:A:123[A]:HIS:H	1.63	0.44
1:A:50:VAL:HG22	1:A:65[B]:ARG:HG2	1.98	0.44
1:A:25:LYS:HD3	1:A:25:LYS:HA	1.76	0.43
1:B:107:GLU:HG3	1:B:121:ARG:HG2	2.01	0.43
1:A:93:GLU:O	1:B:94:THR:HA	2.19	0.43
1:B:25:LYS:HG2	5:B:669:HOH:O	2.18	0.42
1:A:25:LYS:HE2	5:A:704:HOH:O	2.19	0.42
1:B:105:ARG:NH1	5:B:662:HOH:O	2.52	0.42
1:B:52:GLY:H	1:B:138:GLN:NE2	2.18	0.41
1:B:48:ASP:CG	1:B:136:LYS:HZ2	2.14	0.41
1:B:90[A]:GLN:NE2	5:B:690:HOH:O	2.54	0.41
2:B:500:3B4:NAL	2:B:500:3B4:OAA	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/185 (86%)	158 (99%)	2 (1%)	0	100	100
1	B	157/185 (85%)	155 (99%)	2 (1%)	0	100	100
All	All	317/370 (86%)	313 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/161 (88%)	137 (97%)	4 (3%)	43	25
1	B	138/161 (86%)	134 (97%)	4 (3%)	42	23
All	All	279/322 (87%)	271 (97%)	8 (3%)	42	23

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	ASN
1	A	138	GLN
1	A	142	MET
1	A	165	SER
1	B	16	GLN
1	B	122	ASN
1	B	138	GLN
1	B	142	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	122	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	138	GLN
1	B	16	GLN
1	B	32	HIS
1	B	122	ASN
1	B	138	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	3B4	B	500	-	15,17,17	1.95	5 (33%)	18,22,22	2.38	7 (38%)
3	AZI	A	600	-	0,2,2	0.00	-	0,1,1	0.00	-
4	ACT	B	600	-	1,3,3	1.21	0	0,3,3	0.00	-
2	3B4	A	500	-	15,17,17	1.89	4 (26%)	18,22,22	1.58	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3B4	B	500	-	-	3/4/16/16	0/2/2/2
2	3B4	A	500	-	-	3/4/16/16	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	3B4	CAO-CAM	4.33	1.51	1.47
2	B	500	3B4	CAO-CAN	-3.59	1.35	1.40
2	A	500	3B4	CAJ-CAP	-3.58	1.43	1.52
2	B	500	3B4	CAP-NAL	3.46	1.54	1.46
2	A	500	3B4	CAK-CAP	3.18	1.59	1.52
2	B	500	3B4	CAH-CAJ	3.09	1.61	1.53
2	B	500	3B4	CAO-CAM	2.82	1.50	1.47
2	A	500	3B4	CAP-NAL	2.62	1.52	1.46
2	B	500	3B4	CAG-CAH	2.06	1.59	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	3B4	CAJ-CAP-NAL	5.50	119.17	110.60
2	A	500	3B4	CAK-CAP-NAL	-3.71	104.82	110.60
2	A	500	3B4	CAE-CAN-NAL	-3.70	116.38	122.25
2	B	500	3B4	CAK-CAP-NAL	-3.46	105.21	110.60
2	B	500	3B4	CAF-CAO-CAN	3.34	121.28	117.92
2	B	500	3B4	CAK-CAP-CAJ	-3.16	105.33	110.82
2	A	500	3B4	CAN-NAL-CAP	-3.06	119.43	124.69
2	B	500	3B4	CAH-CAJ-CAP	-3.04	105.38	111.11
2	B	500	3B4	CAE-CAN-NAL	-2.30	118.60	122.25
2	B	500	3B4	CAG-CAI-CAK	-2.04	107.26	111.42

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	500	3B4	CAJ-CAP-NAL-CAN
2	B	500	3B4	CAE-CAN-NAL-CAP
2	B	500	3B4	CAO-CAN-NAL-CAP
2	A	500	3B4	CAE-CAN-NAL-CAP
2	A	500	3B4	CAO-CAN-NAL-CAP
2	A	500	3B4	CAJ-CAP-NAL-CAN

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	500	3B4	2	0
4	B	600	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.