



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2023 – 11:23 PM EDT

PDB ID : 3B6H  
Title : Crystal structure of human prostacyclin synthase in complex with inhibitor minoxidil  
Authors : Li, Y.-C.; Chiang, C.-W.; Yeh, H.-C.; Hsu, P.-Y.; Whitby, F.G.; Wang, L.-H.; Chan, N.-L.  
Deposited on : 2007-10-29  
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4693 (1.64-1.60)
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)
RSRZ outliers	127900	4609 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center; margin-top: 5px;">7%      86%      7%      6%</p>
1	B	498	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center; margin-top: 5px;">9%      85%      9%      6%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BOG	B	702	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8552 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostacyclin synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	469	3812	2452	671	674	15	0	1	0
1	B	470	3819	2456	672	676	15	0	1	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	11	MET	-	expression tag	UNP Q16647
A	12	ALA	-	expression tag	UNP Q16647
A	13	LYS	-	expression tag	UNP Q16647
A	14	LYS	-	expression tag	UNP Q16647
A	15	THR	-	expression tag	UNP Q16647
A	16	SER	-	expression tag	UNP Q16647
A	17	SER	-	expression tag	UNP Q16647
A	501	HIS	-	expression tag	UNP Q16647
A	502	HIS	-	expression tag	UNP Q16647
A	503	HIS	-	expression tag	UNP Q16647
A	504	HIS	-	expression tag	UNP Q16647
A	505	HIS	-	expression tag	UNP Q16647
A	506	HIS	-	expression tag	UNP Q16647
A	507	HIS	-	expression tag	UNP Q16647
A	508	HIS	-	expression tag	UNP Q16647
B	11	MET	-	expression tag	UNP Q16647
B	12	ALA	-	expression tag	UNP Q16647
B	13	LYS	-	expression tag	UNP Q16647
B	14	LYS	-	expression tag	UNP Q16647
B	15	THR	-	expression tag	UNP Q16647
B	16	SER	-	expression tag	UNP Q16647
B	17	SER	-	expression tag	UNP Q16647
B	501	HIS	-	expression tag	UNP Q16647
B	502	HIS	-	expression tag	UNP Q16647
B	503	HIS	-	expression tag	UNP Q16647

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	504	HIS	-	expression tag	UNP Q16647
B	505	HIS	-	expression tag	UNP Q16647
B	506	HIS	-	expression tag	UNP Q16647
B	507	HIS	-	expression tag	UNP Q16647
B	508	HIS	-	expression tag	UNP Q16647

- Molecule 2 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	A	1	Total	C	O	0	0
				20	14		
2	B	1	Total	C	O	0	0
				20	14		

- Molecule 3 is 6-PIPERIDIN-1-YL PYRIMIDINE-2,4-DIAMINE 3-OXIDE (three-letter code: MXD) (formula: C<sub>9</sub>H<sub>15</sub>N<sub>5</sub>O).



*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
4	B	1	43	34	1	4	4	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	401	Total	O	0	0
			401	401		
5	B	364	Total	O	0	0
			364	364		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Prostacyclin synthase



- Molecule 1: Prostacyclin synthase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.76Å 106.07Å 73.90Å 90.00° 91.81° 90.00°	Depositor
Resolution (Å)	29.09 – 1.62 29.08 – 1.62	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.09-1.62) 99.6 (29.08-1.62)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.41 (at 1.62Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.203 , 0.228 0.202 , 0.227	Depositor DCC
$R_{free}$ test set	6770 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.1	Xtrriage
Anisotropy	0.018	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 57.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.018 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8552	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MXD, BOG, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3914	0.53	0/5306
1	B	0.37	0/3921	0.53	0/5316
All	All	0.37	0/7835	0.53	0/10622

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3812	0	3792	32	0
1	B	3819	0	3799	31	0
2	A	20	0	28	5	0
2	B	20	0	28	4	0
3	A	15	0	15	2	0
3	B	15	0	15	1	0
4	A	43	0	30	0	0
4	B	43	0	30	6	0
5	A	401	0	0	5	0
5	B	364	0	0	4	0
All	All	8552	0	7737	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:ARG:HH11	1:A:236:ARG:HG3	1.07	1.11
1:A:236:ARG:HG3	1:A:236:ARG:NH1	1.89	0.86
1:B:288:MET:HG2	4:B:600:HEM:HBB1	1.58	0.85
1:B:131:ARG:HH11	1:B:134:GLN:HE22	1.24	0.83
1:B:288:MET:HG2	4:B:600:HEM:CBB	2.09	0.82
4:B:600:HEM:HBB2	4:B:600:HEM:HHC	1.65	0.77
1:B:382:ARG:HH22	2:B:702:BOG:H2	1.51	0.76
1:A:236:ARG:HH11	1:A:236:ARG:CG	1.91	0.76
1:A:245:ARG:HG2	1:A:245:ARG:HH11	1.53	0.73
1:A:211:ASP:O	1:A:480:ARG:NH2	2.23	0.71
1:B:219:ARG:HH22	1:B:479:SER:HB2	1.57	0.69
1:B:141:TYR:CE2	1:B:329:THR:HB	2.29	0.68
1:B:382:ARG:NH2	2:B:702:BOG:H2	2.09	0.66
1:B:99:TYR:CE1	2:B:702:BOG:H3'2	2.30	0.66
1:A:245:ARG:HH11	1:A:245:ARG:CG	2.08	0.65
1:A:422:LYS:HE2	5:A:954:HOH:O	1.96	0.65
1:B:219:ARG:HG3	5:B:928:HOH:O	1.98	0.63
1:A:121:LYS:HE2	5:A:933:HOH:O	1.98	0.62
1:A:501:HIS:HB2	5:B:964:HOH:O	2.00	0.62
1:B:288:MET:CG	4:B:600:HEM:CBB	2.81	0.57
1:B:141:TYR:HE2	1:B:329:THR:HB	1.69	0.57
1:A:382:ARG:HH12	2:A:701:BOG:H4	1.72	0.55
1:B:131:ARG:HH11	1:B:134:GLN:NE2	2.00	0.55
1:B:211:ASP:O	1:B:480:ARG:NH2	2.40	0.55
1:B:131:ARG:NH1	1:B:134:GLN:HE22	2.00	0.52
1:B:216:LYS:HG3	1:B:221:SER:OG	2.09	0.52
1:B:222:LEU:HD22	1:B:226:ASP:HB3	1.92	0.52
1:A:382:ARG:HH22	2:A:701:BOG:H2	1.76	0.50
1:A:382:ARG:NH2	2:A:701:BOG:H2	2.26	0.50
1:B:206:THR:CG2	1:B:233:VAL:HG23	2.42	0.49
1:A:356:PHE:HD2	2:A:701:BOG:H4'1	1.77	0.49
1:A:181:TYR:HA	1:A:256:LEU:HD22	1.94	0.49
1:A:145:HIS:NE2	1:A:149:LEU:HD12	2.28	0.48
1:B:88:TRP:CE2	1:B:437:GLY:HA2	2.49	0.48
3:B:551:MXD:HAL2	4:B:600:HEM:CGA	2.42	0.48
1:B:137:THR:HG21	1:B:332:GLN:HE22	1.79	0.48
1:B:209:GLN:OE1	1:B:229:HIS:HE1	1.96	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:THR:HG21	1:A:188:ARG:HH22	1.79	0.47
1:A:88:TRP:CE2	1:A:437:GLY:HA2	2.50	0.47
1:A:188:ARG:HG3	1:A:193:GLN:HG3	1.98	0.46
1:B:257:GLU:O	1:B:261:LEU:HD22	2.17	0.45
1:A:333:LYS:HA	1:A:336:ASP:OD1	2.17	0.44
1:A:499:ARG:NH2	5:A:964:HOH:O	2.50	0.44
1:B:91:ARG:HG3	1:B:96:PHE:CE2	2.54	0.43
1:A:121:LYS:NZ	1:A:442:LEU:HD21	2.34	0.42
1:A:111:GLN:NE2	5:A:1069:HOH:O	2.52	0.42
1:B:371:ASP:OD1	1:B:373:ARG:HD3	2.19	0.42
1:A:88:TRP:CZ2	1:A:437:GLY:HA2	2.54	0.42
1:A:103:LEU:HD23	3:A:551:MXD:HAH2	2.02	0.42
1:B:245:ARG:HD2	1:B:248:ARG:HH21	1.83	0.42
1:A:102:PHE:O	1:A:106:ARG:HB3	2.20	0.41
1:B:206:THR:HG23	1:B:233:VAL:HG23	2.00	0.41
1:A:125:LYS:HD3	1:A:442:LEU:HD22	2.02	0.41
3:A:551:MXD:HAF	3:A:551:MXD:HAK2	1.81	0.41
1:B:196:ASP:HA	5:B:934:HOH:O	2.20	0.41
1:B:313:LEU:HD22	1:B:456:PHE:CZ	2.55	0.41
1:B:475:GLU:O	1:B:489:GLU:HG2	2.21	0.41
4:B:600:HEM:HBB2	4:B:600:HEM:CHC	2.39	0.41
1:A:236:ARG:NH1	1:A:236:ARG:CG	2.60	0.40
1:A:486:MET:HE3	5:A:834:HOH:O	2.21	0.40
1:B:124:MET:HE2	5:B:799:HOH:O	2.21	0.40
1:A:141:TYR:CE1	1:A:330:LEU:HD23	2.56	0.40
1:A:245:ARG:CG	1:A:245:ARG:NH1	2.74	0.40
1:A:77:LEU:HB3	1:A:385:LEU:HD22	2.02	0.40
1:B:356:PHE:HD2	2:B:702:BOG:H4'2	1.87	0.40
1:A:99:TYR:CD1	2:A:701:BOG:H3'1	2.57	0.40
1:B:193:GLN:HE21	1:B:193:GLN:HB3	1.70	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/498 (94%)	460 (99%)	6 (1%)	0	100	100
1	B	467/498 (94%)	462 (99%)	5 (1%)	0	100	100
All	All	933/996 (94%)	922 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/433 (94%)	399 (98%)	8 (2%)	55	29
1	B	408/433 (94%)	395 (97%)	13 (3%)	39	14
All	All	815/866 (94%)	794 (97%)	21 (3%)	46	19

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	MET
1	A	148	LEU
1	A	169	TYR
1	A	213	LEU
1	A	236	ARG
1	A	317	LEU
1	A	373	ARG
1	A	499	ARG
1	B	45	ASP
1	B	77	LEU
1	B	169	TYR
1	B	186	LEU
1	B	213	LEU
1	B	240	LEU
1	B	245	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	261	LEU
1	B	304	GLU
1	B	306	LEU
1	B	329	THR
1	B	373	ARG
1	B	459	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	GLN
1	A	190	HIS
1	A	449	ASN
1	A	453	GLN
1	A	468	ASN
1	A	487	GLN
1	A	490	HIS
1	B	97	HIS
1	B	130	HIS
1	B	134	GLN
1	B	195	GLN
1	B	229	HIS
1	B	332	GLN
1	B	449	ASN
1	B	468	ASN
1	B	490	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BOG	B	702	-	20,20,20	0.66	1 (5%)	25,25,25	0.66	0
2	BOG	A	701	-	20,20,20	0.63	1 (5%)	25,25,25	0.63	0
4	HEM	A	600	3,1	41,50,50	1.86	6 (14%)	45,82,82	1.71	8 (17%)
3	MXD	A	551	4	14,16,16	3.14	1 (7%)	16,22,22	2.07	2 (12%)
4	HEM	B	600	3,1	41,50,50	1.94	7 (17%)	45,82,82	1.77	8 (17%)
3	MXD	B	551	4	14,16,16	3.00	1 (7%)	16,22,22	2.15	4 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BOG	B	702	-	-	6/11/31/31	0/1/1/1
2	BOG	A	701	-	-	6/11/31/31	0/1/1/1
4	HEM	A	600	3,1	-	4/12/54/54	-
3	MXD	A	551	4	-	0/4/12/12	0/2/2/2
4	HEM	B	600	3,1	-	5/12/54/54	-
3	MXD	B	551	4	-	0/4/12/12	0/2/2/2

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	551	MXD	OAN-NAC	-11.47	1.24	1.38
3	B	551	MXD	OAN-NAC	-10.79	1.25	1.38
4	B	600	HEM	C3D-C2D	7.67	1.53	1.36
4	A	600	HEM	C3D-C2D	7.48	1.52	1.36
4	B	600	HEM	C3C-C2C	-4.68	1.33	1.40
4	A	600	HEM	C3C-C2C	-3.88	1.35	1.40

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	600	HEM	C3C-CAC	3.79	1.55	1.47
4	A	600	HEM	C3C-CAC	3.68	1.55	1.47
4	B	600	HEM	CAB-C3B	2.89	1.55	1.47
4	A	600	HEM	CAB-C3B	2.87	1.55	1.47
2	B	702	BOG	O1-C1	2.42	1.44	1.40
4	A	600	HEM	FE-ND	2.35	2.08	1.96
2	A	701	BOG	O1-C1	2.27	1.44	1.40
4	A	600	HEM	CAA-C2A	2.26	1.55	1.52
4	B	600	HEM	CAA-C2A	2.22	1.55	1.52
4	B	600	HEM	FE-ND	2.09	2.07	1.96
4	B	600	HEM	FE-NB	2.06	2.07	1.96

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	600	HEM	C4D-ND-C1D	5.74	111.00	105.07
4	A	600	HEM	C4D-ND-C1D	5.70	110.96	105.07
3	A	551	MXD	NAM-CAA-NAC	5.53	120.25	116.46
3	B	551	MXD	NAM-CAA-NAC	5.01	119.89	116.46
4	B	600	HEM	CBA-CAA-C2A	-4.85	104.34	112.62
3	A	551	MXD	CAK-NAI-CAG	4.83	122.18	111.52
3	B	551	MXD	CAK-NAI-CAG	4.72	121.94	111.52
4	A	600	HEM	CBA-CAA-C2A	-3.71	106.29	112.62
3	B	551	MXD	NAB-CAD-NAI	3.40	120.36	116.55
4	A	600	HEM	C4C-CHD-C1D	3.07	126.61	122.56
4	A	600	HEM	C1B-NB-C4B	3.03	108.20	105.07
4	B	600	HEM	C1B-NB-C4B	2.96	108.13	105.07
4	B	600	HEM	C3B-C2B-C1B	2.84	108.59	106.49
4	B	600	HEM	C4B-CHC-C1C	2.83	126.29	122.56
4	B	600	HEM	C4C-CHD-C1D	2.79	126.23	122.56
4	A	600	HEM	C4B-CHC-C1C	2.61	126.01	122.56
4	A	600	HEM	CMA-C3A-C4A	-2.47	124.66	128.46
4	B	600	HEM	CBD-CAD-C3D	-2.30	106.24	112.63
4	A	600	HEM	C3B-C2B-C1B	2.29	108.18	106.49
3	B	551	MXD	CAF-CAD-NAI	-2.27	119.59	122.29
4	B	600	HEM	CHC-C4B-C3B	2.17	127.88	124.57
4	A	600	HEM	CMD-C2D-C1D	2.10	128.24	125.04

There are no chirality outliers.

All (21) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	B	702	BOG	O5-C1-O1-C1'
2	B	702	BOG	C2'-C1'-O1-C1
2	A	701	BOG	C2-C1-O1-C1'
2	B	702	BOG	C2-C1-O1-C1'
2	B	702	BOG	C4'-C5'-C6'-C7'
2	B	702	BOG	C3'-C4'-C5'-C6'
2	A	701	BOG	C5'-C6'-C7'-C8'
2	A	701	BOG	O1-C1'-C2'-C3'
2	A	701	BOG	C4'-C5'-C6'-C7'
4	B	600	HEM	C4B-C3B-CAB-CBB
2	B	702	BOG	C1'-C2'-C3'-C4'
2	A	701	BOG	C4-C5-C6-O6
4	A	600	HEM	CAD-CBD-CGD-O1D
4	B	600	HEM	CAA-CBA-CGA-O1A
4	A	600	HEM	CAA-CBA-CGA-O1A
4	B	600	HEM	CAA-CBA-CGA-O2A
4	A	600	HEM	CAD-CBD-CGD-O2D
4	A	600	HEM	CAA-CBA-CGA-O2A
4	B	600	HEM	CAD-CBD-CGD-O2D
4	B	600	HEM	CAD-CBD-CGD-O1D
2	A	701	BOG	O5-C1-O1-C1'


There are no ring outliers.


5 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	702	BOG	4	0
2	A	701	BOG	5	0
3	A	551	MXD	2	0
4	B	600	HEM	6	0
3	B	551	MXD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	469/498 (94%)	0.40	33 (7%) 16 14	9, 19, 31, 40	0
1	B	470/498 (94%)	0.42	44 (9%) 8 7	10, 20, 33, 40	0
All	All	939/996 (94%)	0.41	77 (8%) 11 9	9, 20, 32, 40	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	TRP	7.2
1	B	114	HIS	7.0
1	A	115	TYR	5.5
1	A	114	HIS	5.3
1	A	331	PRO	4.9
1	B	115	TYR	4.7
1	B	154	GLU	4.6
1	B	329	THR	4.5
1	B	40	LEU	4.3
1	A	186	LEU	4.2
1	B	318	TRP	4.1
1	B	156	GLY	4.0
1	B	37	ILE	4.0
1	B	38	PRO	3.9
1	B	501	HIS	3.8
1	A	501	HIS	3.7
1	B	319	GLN	3.7
1	B	187	PRO	3.6
1	A	330	LEU	3.5
1	A	130	HIS	3.5
1	B	261	LEU	3.4
1	A	113	PRO	3.4
1	A	332	GLN	3.4
1	B	317	LEU	3.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	261	LEU	3.2
1	A	334	VAL	3.2
1	A	248	ARG	3.1
1	B	248	ARG	3.1
1	A	36	SER	2.9
1	B	333	LYS	2.9
1	A	40	LEU	2.9
1	B	44	LEU	2.8
1	A	38	PRO	2.8
1	B	186	LEU	2.7
1	B	456	PHE	2.7
1	A	187	PRO	2.7
1	B	414	GLY	2.7
1	A	39	TRP	2.7
1	B	117	PRO	2.7
1	B	158	GLY	2.6
1	A	91	ARG	2.6
1	B	219	ARG	2.6
1	B	331	PRO	2.6
1	B	155	ALA	2.6
1	B	330	LEU	2.6
1	A	133	LEU	2.5
1	A	333	LYS	2.5
1	B	84	ASP	2.5
1	A	111	GLN	2.4
1	B	85	ALA	2.4
1	B	316	ILE	2.4
1	A	119	ASP	2.4
1	A	372	GLY	2.4
1	B	119	ASP	2.3
1	B	413	ASP	2.3
1	A	190	HIS	2.3
1	A	173	LEU	2.2
1	A	185	ALA	2.2
1	B	91	ARG	2.2
1	B	334	VAL	2.2
1	A	335	LEU	2.2
1	B	113	PRO	2.2
1	B	160	HIS	2.2
1	B	36	SER	2.2
1	B	116	SER	2.2
1	A	85	ALA	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	127	THR	2.2
1	A	470	ASP	2.2
1	B	412	PRO	2.1
1	A	126	LEU	2.1
1	B	265	GLU	2.1
1	B	424	GLY	2.1
1	B	336	ASP	2.1
1	A	219	ARG	2.1
1	B	118	SER	2.1
1	B	92	THR	2.0
1	B	315	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BOG	B	702	20/20	0.13	0.44	61,66,67,67	0
2	BOG	A	701	20/20	0.29	0.37	37,49,50,50	0
3	MXD	B	551	15/15	0.90	0.13	16,19,25,26	0
3	MXD	A	551	15/15	0.94	0.11	15,18,24,25	0
4	HEM	B	600	43/43	0.97	0.11	12,14,24,24	0
4	HEM	A	600	43/43	0.98	0.11	11,13,19,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around BOG B 702:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)


**Electron density around BOG A 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









## 6.5 Other polymers [i](#)

There are no such residues in this entry.