



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 24, 2021 – 01:24 PM EST

PDB ID : 3C4R  
Title : Crystal structure of an uncharacterized protein encoded by cryptic prophage  
Authors : Sugadev, R.; Burley, S.K.; Swaminathan, S.; Ozyurt, S.; Luz, J.; New York  
SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2008-01-30  
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	 4% 55% 25% 18%
1	B	151	 4% 50% 28% 18%
1	C	151	 0% 58% 22% 16%
1	D	151	 3% 47% 35% 16%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	124	1001	638	177	181	5	0	0	0
1	B	124	1001	638	177	181	5	0	0	0
1	C	127	1026	652	182	186	6	0	0	0
1	D	127	1026	652	182	186	6	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	expression tag	UNP Q8FILO
A	0	SER	-	expression tag	UNP Q8FILO
A	1	LEU	-	expression tag	UNP Q8FILO
A	142	GLU	-	expression tag	UNP Q8FILO
A	143	GLY	-	expression tag	UNP Q8FILO
A	144	HIS	-	expression tag	UNP Q8FILO
A	145	HIS	-	expression tag	UNP Q8FILO
A	146	HIS	-	expression tag	UNP Q8FILO
A	147	HIS	-	expression tag	UNP Q8FILO
A	148	HIS	-	expression tag	UNP Q8FILO
A	149	HIS	-	expression tag	UNP Q8FILO
B	-1	MSE	-	expression tag	UNP Q8FILO
B	0	SER	-	expression tag	UNP Q8FILO
B	1	LEU	-	expression tag	UNP Q8FILO
B	142	GLU	-	expression tag	UNP Q8FILO
B	143	GLY	-	expression tag	UNP Q8FILO
B	144	HIS	-	expression tag	UNP Q8FILO
B	145	HIS	-	expression tag	UNP Q8FILO
B	146	HIS	-	expression tag	UNP Q8FILO
B	147	HIS	-	expression tag	UNP Q8FILO
B	148	HIS	-	expression tag	UNP Q8FILO

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	149	HIS	-	expression tag	UNP Q8FIL0
C	-1	MSE	-	expression tag	UNP Q8FIL0
C	0	SER	-	expression tag	UNP Q8FIL0
C	1	LEU	-	expression tag	UNP Q8FIL0
C	142	GLU	-	expression tag	UNP Q8FIL0
C	143	GLY	-	expression tag	UNP Q8FIL0
C	144	HIS	-	expression tag	UNP Q8FIL0
C	145	HIS	-	expression tag	UNP Q8FIL0
C	146	HIS	-	expression tag	UNP Q8FIL0
C	147	HIS	-	expression tag	UNP Q8FIL0
C	148	HIS	-	expression tag	UNP Q8FIL0
C	149	HIS	-	expression tag	UNP Q8FIL0
D	-1	MSE	-	expression tag	UNP Q8FIL0
D	0	SER	-	expression tag	UNP Q8FIL0
D	1	LEU	-	expression tag	UNP Q8FIL0
D	142	GLU	-	expression tag	UNP Q8FIL0
D	143	GLY	-	expression tag	UNP Q8FIL0
D	144	HIS	-	expression tag	UNP Q8FIL0
D	145	HIS	-	expression tag	UNP Q8FIL0
D	146	HIS	-	expression tag	UNP Q8FIL0
D	147	HIS	-	expression tag	UNP Q8FIL0
D	148	HIS	-	expression tag	UNP Q8FIL0
D	149	HIS	-	expression tag	UNP Q8FIL0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	22	Total O 22 22	0	0
2	B	13	Total O 13 13	0	0
2	C	15	Total O 15 15	0	0
2	D	15	Total O 15 15	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	36.83Å 106.57Å 88.25Å 90.00° 96.42° 90.00°	Depositor
Resolution (Å)	31.09 – 2.30 45.54 – 2.31	Depositor EDS
% Data completeness (in resolution range)	79.9 (31.09-2.30) 80.0 (45.54-2.31)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.08 (at 2.32Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.242 , 0.282 0.241 , 0.280	Depositor DCC
$R_{free}$ test set	1394 reflections (5.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.8	Xtrriage
Anisotropy	0.694	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 44.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4119	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/1021	0.61	0/1374
1	B	0.37	0/1021	0.56	0/1374
1	C	0.39	0/1045	0.64	1/1404 (0.1%)
1	D	0.41	0/1045	0.61	0/1404
All	All	0.39	0/4132	0.61	1/5556 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	123	ALA	N-CA-C	-5.27	96.77	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1001	0	991	29	0
1	B	1001	0	991	44	0
1	C	1026	0	1014	27	0
1	D	1026	0	1014	58	0
2	A	22	0	0	1	0
2	B	13	0	0	0	0
2	C	15	0	0	1	0
2	D	15	0	0	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4119	0	4010	147	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 147 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:8:ILE:HA	1:B:92:MSE:HE2	1.40	1.03
1:B:11:ALA:HB3	1:B:92:MSE:HE3	1.40	0.99
1:D:84:MSE:HE3	1:D:89:VAL:HG22	1.48	0.94
1:B:10:MSE:HG3	1:B:11:ALA:N	1.93	0.84
1:D:44:ASP:HB2	2:D:150:HOH:O	1.80	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	122/151 (81%)	115 (94%)	4 (3%)	3 (2%)	5	4
1	B	122/151 (81%)	118 (97%)	2 (2%)	2 (2%)	9	9
1	C	125/151 (83%)	115 (92%)	8 (6%)	2 (2%)	9	9
1	D	125/151 (83%)	119 (95%)	6 (5%)	0	100	100
All	All	494/604 (82%)	467 (94%)	20 (4%)	7 (1%)	11	11

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	18	PRO
1	B	48	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	50	ALA
1	C	18	PRO
1	A	48	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/118 (88%)	98 (94%)	6 (6%)	20	27
1	B	104/118 (88%)	96 (92%)	8 (8%)	13	16
1	C	107/118 (91%)	101 (94%)	6 (6%)	21	29
1	D	107/118 (91%)	100 (94%)	7 (6%)	17	23
All	All	422/472 (89%)	395 (94%)	27 (6%)	17	23

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	117	PHE
1	C	30	ARG
1	D	96	SER
1	B	122	ILE
1	A	110	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	13	ASN
1	D	56	GLN
1	B	95	HIS
1	B	7	HIS
1	B	54	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	119/151 (78%)	0.47	2 (1%) 70 76	31, 44, 70, 78	0
1	B	119/151 (78%)	0.59	6 (5%) 28 35	32, 55, 84, 96	0
1	C	121/151 (80%)	0.42	2 (1%) 70 76	28, 46, 62, 78	0
1	D	121/151 (80%)	0.65	4 (3%) 46 53	31, 48, 80, 97	0
All	All	480/604 (79%)	0.53	14 (2%) 51 58	28, 48, 76, 97	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	125	PHE	8.1
1	D	127	GLN	3.9
1	B	51	LEU	3.4
1	B	43	TYR	3.0
1	C	2	LYS	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.