



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 08:17 pm BST

PDB ID : 3COO
Title : The crystal structure of Reelin-N domain of F-spondin
Authors : Tan, K.; Lawler, J.; Wang, J.H.
Deposited on : 2008-03-28
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	180	60% 16% • 21%
1	B	180	63% 10% •• 23%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2440 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spondin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	143	1161	733	202	221	5	0	3	0
1	B	138	1126	715	197	209	5	0	4	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	ARG	-	EXPRESSION TAG	UNP Q9HCB6
A	24	SER	-	EXPRESSION TAG	UNP Q9HCB6
A	25	PRO	-	EXPRESSION TAG	UNP Q9HCB6
A	26	TRP	-	EXPRESSION TAG	UNP Q9HCB6
A	27	PRO	-	EXPRESSION TAG	UNP Q9HCB6
A	28	GLY	-	EXPRESSION TAG	UNP Q9HCB6
A	195	THR	-	EXPRESSION TAG	UNP Q9HCB6
A	196	GLY	-	EXPRESSION TAG	UNP Q9HCB6
A	197	HIS	-	EXPRESSION TAG	UNP Q9HCB6
A	198	HIS	-	EXPRESSION TAG	UNP Q9HCB6
A	199	HIS	-	EXPRESSION TAG	UNP Q9HCB6
A	200	HIS	-	EXPRESSION TAG	UNP Q9HCB6
A	201	HIS	-	EXPRESSION TAG	UNP Q9HCB6
A	202	HIS	-	EXPRESSION TAG	UNP Q9HCB6
B	23	ARG	-	EXPRESSION TAG	UNP Q9HCB6
B	24	SER	-	EXPRESSION TAG	UNP Q9HCB6
B	25	PRO	-	EXPRESSION TAG	UNP Q9HCB6
B	26	TRP	-	EXPRESSION TAG	UNP Q9HCB6
B	27	PRO	-	EXPRESSION TAG	UNP Q9HCB6
B	28	GLY	-	EXPRESSION TAG	UNP Q9HCB6
B	195	THR	-	EXPRESSION TAG	UNP Q9HCB6
B	196	GLY	-	EXPRESSION TAG	UNP Q9HCB6
B	197	HIS	-	EXPRESSION TAG	UNP Q9HCB6
B	198	HIS	-	EXPRESSION TAG	UNP Q9HCB6
B	199	HIS	-	EXPRESSION TAG	UNP Q9HCB6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	200	HIS	-	EXPRESSION TAG	UNP Q9HCB6
B	201	HIS	-	EXPRESSION TAG	UNP Q9HCB6
B	202	HIS	-	EXPRESSION TAG	UNP Q9HCB6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	79	Total O 79 79	0	0
2	B	74	Total O 74 74	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	58.60Å 58.60Å 219.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.06 – 2.00 40.06 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.0 (40.06-2.00) 99.0 (40.06-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.93 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.206 , 0.241 0.204 , 0.241	Depositor DCC
R_{free} test set	1346 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	36.8	Xtrriage
Anisotropy	0.066	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2440	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/1190	0.93	5/1606 (0.3%)
1	B	0.85	0/1161	0.84	2/1567 (0.1%)
All	All	0.84	0/2351	0.89	7/3173 (0.2%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	59	THR	N-CA-CB	-5.63	99.59	110.30
1	A	138	ARG	NE-CZ-NH2	5.54	123.07	120.30
1	B	95	LEU	CA-CB-CG	5.51	127.97	115.30
1	A	99[A]	ARG	NE-CZ-NH1	-5.29	117.66	120.30
1	A	99[B]	ARG	NE-CZ-NH1	-5.29	117.66	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1161	0	1145	21	0
1	B	1126	0	1125	16	0
2	A	79	0	0	1	0
2	B	74	0	0	3	0
All	All	2440	0	2270	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 37 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:GLN:HG2	2:A:257:HOH:O	1.40	1.16
1:A:121:GLU:HA	1:A:135:SER:OG	1.78	0.83
1:A:63:LEU:O	1:A:179:LYS:HE3	1.81	0.81
1:A:79:ARG:HD3	1:A:144:GLN:HE22	1.56	0.70
1:B:59:THR:HG21	2:B:227:HOH:O	1.91	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/180 (79%)	139 (98%)	3 (2%)	0	100	100
1	B	138/180 (77%)	133 (96%)	5 (4%)	0	100	100
All	All	280/360 (78%)	272 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/157 (81%)	118 (93%)	9 (7%)	14	10
1	B	123/157 (78%)	111 (90%)	12 (10%)	8	4
All	All	250/314 (80%)	229 (92%)	21 (8%)	12	7

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	43	TYR
1	B	47[B]	ILE
1	B	138	ARG
1	A	174	GLU
1	B	141	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	184	GLN
1	B	165	GLN
1	B	115	GLN
1	A	144	GLN
1	B	123	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.