

Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2021 – 09:12 PM EDT

PDB ID 3CUL

> Title : Aminoacyl-tRNA synthetase ribozyme

Authors Xiao, H.; Murakami, H.; Suga, H.; Ferre-D'Amare, A.R.

2008-04-16 Deposited on

2.80 Å(reported) Resolution

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as541be (2020) Mogul

Xtriage (Phenix) 1.13 EDS 2.23.2

20191225.v01 (using entries in the PDB archive December 25th 2019) Percentile statistics

> Refmac 5.8.0158

CCP4 7.0.044 (Gargrove) Engh & Huber (2001)

Ideal geometry (proteins) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)
RNA backbone	3102	1227 (3.10-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	С	92	42%	48%	10%			
2	D	92	52%	40%	5% •			
3	A	98	63%	27%	• 6%			
3	В	98	55%	36%	• 7%			



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	С	508	-	-	-	X



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called RNA (92-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	С	92	Total	C 071	N	O	P	0	0	0
			1965	871	348	652	94			

• Molecule 2 is a RNA chain called RNA (92-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	D	92	Total 1964	C 875	N 350	O 647	P 92	0	0	0

• Molecule 3 is a protein called U1 small nuclear ribonucleoprotein A.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	A	92	Total 706	_	N 122	_	Se 3	0	0	0
3	В	91	Total 688	C 444	N 117	O 124	Se 3	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	326	HIS	TYR	engineered mutation	UNP P09012
A	331	ARG	GLN	engineered mutation	UNP P09012
В	426	HIS	TYR	engineered mutation	UNP P09012
В	431	ARG	GLN	engineered mutation	UNP P09012

• Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	8	Total Mg 8 8	0	0
4	D	2	Total Mg 2 2	0	0



• Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	С	1	Total K 1 1	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	С	5	Total O 5 5	0	0
6	D	2	Total O 2 2	0	0
6	A	1	Total O 1 1	0	0
6	В	2	Total O 2 2	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA (92-MER)





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	192.15Å 48.73Å 90.52Å	Donositor
a, b, c, α , β , γ	90.00° 93.50° 90.00°	Depositor
Resolution (Å)	29.57 - 2.80	Depositor
rtesolution (A)	29.57 - 2.75	EDS
% Data completeness	99.1 (29.57-2.80)	Depositor
(in resolution range)	99.0 (29.57-2.75)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$< I/\sigma(I) > 1$	2.49 (at 2.76Å)	Xtriage
Refinement program	CNS 1.1	Depositor
P. P.	0.227 , 0.283	Depositor
R, R_{free}	0.225 , 0.279	DCC
R_{free} test set	2181 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å ²)	82.2	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.26 , 48.6	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5344	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GTP, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
MIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	С	0.55	$1/2159 \ (0.0\%)$	0.80	3/3364 (0.1%)	
2	D	0.44	0/2194	0.77	5/3419 (0.1%)	
3	A	0.51	0/715	0.72	0/959	
3	В	0.42	0/698	0.67	0/940	
All	All	0.49	$1/5766 \ (0.0\%)$	0.76	8/8682 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	1
2	D	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
1	С	2	G	O3'-P	-9.53	1.49	1.61

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
1	С	34	С	N1-C1'-C2'	8.76	125.39	114.00
2	D	134	С	C2'-C3'-O3'	8.18	127.49	109.50
1	С	53	U	C2'-C3'-O3'	8.10	127.32	109.50
2	D	153	U	C4'-C3'-O3'	5.60	124.20	113.00
2	D	153	U	C4'-C3'-C2'	5.43	108.03	102.60
1	С	53	U	C4'-C3'-C2'	5.29	107.89	102.60



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	D	134	С	C4'-C3'-C2'	5.15	107.75	102.60
2	D	175	A	N9-C1'-C2'	5.00	120.51	114.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	С	30	A	Sidechain
2	D	108	G	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1965	0	989	48	0
2	D	1964	0	993	33	0
3	A	706	0	687	26	0
3	В	688	0	661	42	0
4	С	8	0	0	0	0
4	D	2	0	0	0	0
5	С	1	0	0	0	0
6	A	1	0	0	0	0
6	В	2	0	0	1	0
6	С	5	0	0	0	0
6	D	2	0	0	0	0
All	All	5344	0	3330	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (140) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)	
1:C:32:U:H3	3:A:311:ASN:HD22	1.10	0.98	
3:B:444:LEU:HG	3:B:445:LYS:HE3	1.66	0.78	



 $Continued\ from\ previous\ page...$

Continuea from prev		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$\text{overlap } (\mathring{\mathbf{A}})$	
1:C:32:U:H3	3:A:311:ASN:ND2	1.79	0.78	
1:C:86:G:H2'	1:C:87:C:H5'	1.66	0.78	
3:B:443:SER:O	3:B:447:ARG:HG3	1.87	0.75	
3:A:357:VAL:O	3:A:361:THR:HG22	1.86	0.74	
1:C:56:G:H2'	1:C:57:U:C6	2.23	0.72	
3:B:465:ARG:HH11	3:B:465:ARG:HG3	1.53	0.72	
2:D:127:C:H2'	2:D:128:C:C6	2.25	0.72	
3:B:405:HIS:CD2	3:B:457:VAL:HG22	2.31	0.66	
2:D:127:C:H2'	2:D:128:C:H6	1.62	0.64	
2:D:122:C:H2'	2:D:123:A:C8	2.31	0.64	
1:C:76:U:H2'	1:C:77:C:C6	2.33	0.64	
1:C:57:U:O5'	1:C:57:U:H6	1.82	0.63	
1:C:25:G:O2'	1:C:26:C:H5'	1.99	0.63	
1:C:22:C:H2'	1:C:23:A:C8	2.34	0.62	
1:C:40:G:H1'	3:A:344:LEU:HD22	1.83	0.61	
3:B:416:ILE:HG12	3:B:473:TYR:CD2	2.36	0.61	
3:B:485:ASP:OD2	3:B:489:ILE:HG22	2.00	0.61	
1:C:40:G:OP2	3:A:343:SER:HA	2.00	0.60	
1:C:64:A:H2'	1:C:65:C:C6	2.36	0.60	
3:B:488:ILE:C	3:B:488:ILE:HD12	2.21	0.59	
3:B:405:HIS:NE2	3:B:457:VAL:HG22	2.17	0.59	
3:B:445:LYS:HA	3:B:445:LYS:HE2	1.85	0.59	
1:C:82:U:H2'	1:C:83:A:H8	1.66	0.59	
3:B:484:THR:HG22	3:B:485:ASP:N	2.17	0.59	
3:B:440:VAL:HG22	3:B:450:ALA:HB2	1.85	0.59	
3:B:416:ILE:HD12	3:B:416:ILE:N	2.18	0.58	
3:B:410:ASN:H	3:B:410:ASN:HD22	1.50	0.58	
3:B:447:ARG:O	3:B:449:GLN:HG2	2.04	0.57	
1:C:35:A:H2	3:A:339:LEU:HD12	1.69	0.57	
2:D:129:C:H2'	2:D:130:A:O4'	2.04	0.57	
3:A:364:LEU:C	3:A:364:LEU:HD12	2.25	0.57	
3:A:368:GLN:HE22	3:A:379:ILE:H	1.53	0.57	
2:D:122:C:H5'	2:D:154:A:N6	2.20	0.56	
1:C:86:G:H2'	1:C:87:C:C5'	2.34	0.55	
1:C:1:GTP:H1'	1:C:52:U:O4	2.05	0.55	
2:D:162:G:H2'	2:D:163:U:C6	2.42	0.55	
1:C:40:G:C1'	3:A:344:LEU:HD22	2.37	0.55	
2:D:157:U:H2'	2:D:158:G:C8	2.42	0.54	
2:D:144:U:H2'	2:D:145:A:C8	2.42	0.54	
3:A:306:THR:OG1	3:A:353:ILE:HG23	2.08	0.54	
1:C:86:G:C2'	1:C:87:C:H5'	2.36	0.54	



 $Continued\ from\ previous\ page...$

A + 1	A4 0	Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$\text{overlap } (\mathring{\mathbf{A}})$	
3:B:412:LEU:HD12	3:B:449:GLN:HA	1.89	0.54	
1:C:89:G:H2'	1:C:90:C:C6	2.43	0.54	
3:B:416:ILE:HG12	3:B:473:TYR:CE2	2.43	0.54	
3:A:303:PRO:HA	3:A:381:TYR:CE1	2.42	0.54	
1:C:31:U:O2	1:C:33:G:N2	2.41	0.53	
3:B:465:ARG:HG3	3:B:465:ARG:NH1	2.22	0.53	
1:C:11:A:H2'	1:C:12:G:O4'	2.09	0.53	
2:D:141:G:H2'	2:D:142:G:H8	1.74	0.53	
3:B:438:ILE:HG23	3:B:452:VAL:HG22	1.91	0.53	
3:B:404:ASN:HD21	3:B:485:ASP:HA	1.73	0.53	
3:B:404:ASN:OD1	3:B:405:HIS:N	2.42	0.53	
3:A:332:PHE:HZ	3:A:367:MSE:HE3	1.74	0.52	
3:B:484:THR:CG2	3:B:485:ASP:N	2.72	0.52	
1:C:12:G:H2'	1:C:13:C:C6	2.45	0.52	
3:B:440:VAL:HG22	3:B:450:ALA:CB	2.39	0.52	
1:C:57:U:C6	1:C:57:U:O5'	2.63	0.51	
3:A:321:LEU:HD23	3:A:340:VAL:HG11	1.91	0.51	
1:C:82:U:H2'	1:C:83:A:C8	2.44	0.51	
2:D:158:G:H8	2:D:158:G:O5'	1.93	0.51	
2:D:187:C:H2'	2:D:188:A:C8	2.46	0.51	
2:D:144:U:H2'	2:D:145:A:H8	1.76	0.51	
2:D:167:A:H2'	2:D:168:G:H8	1.74	0.51	
2:D:186:G:H2'	2:D:187:C:C6	2.46	0.51	
2:D:167:A:H2'	2:D:168:G:C8	2.45	0.51	
1:C:68:G:H2'	1:C:69:G:O4'	2.12	0.50	
1:C:2:G:C6	1:C:3:A:C5	2.99	0.50	
3:B:444:LEU:CG	3:B:445:LYS:HE3	2.37	0.50	
3:B:441:SER:HB3	3:B:446:MSE:CB	2.42	0.50	
1:C:2:G:O2'	1:C:3:A:H5'	2.12	0.49	
3:B:444:LEU:CD2	3:B:445:LYS:HE3	2.42	0.49	
1:C:2:G:C6	1:C:3:A:C6	3.01	0.49	
2:D:151:G:H4'	2:D:152:U:OP1	2.12	0.49	
3:A:341:SER:HB3	3:A:346:MSE:HB3	1.93	0.48	
1:C:14:C:O2'	1:C:15:A:H5'	2.12	0.48	
1:C:25:G:C2'	1:C:26:C:H5'	2.43	0.48	
3:A:358:SER:HA	3:A:361:THR:CG2	2.43	0.48	
3:B:413:ASN:HB3	3:B:416:ILE:HD13	1.96	0.47	
1:C:16:U:O2'	1:C:17:U:H5'	2.14	0.47	
3:A:303:PRO:HA	3:A:381:TYR:CD1	2.50	0.47	
3:B:472:PHE:CD2	3:B:477:MSE:HE3	2.50	0.47	
3:B:432:PHE:O	3:B:459:SER:HB2	2.14	0.47	



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)	
2:D:153:U:H6	2:D:153:U:O5'	1.97	0.47	
3:A:316:ILE:HG13	3:A:373:TYR:CD1	2.50	0.46	
3:B:470:PHE:CD2	3:B:471:PRO:O	2.68	0.46	
2:D:110:A:H2'	2:D:111:A:C8	2.51	0.46	
2:D:134:C:O2'	2:D:135:A:OP2	2.29	0.46	
1:C:87:C:H2'	1:C:88:A:C8	2.51	0.46	
1:C:80:C:H6	1:C:80:C:O5'	1.99	0.45	
1:C:15:A:H2'	1:C:16:U:C6	2.52	0.45	
2:D:163:U:O5'	2:D:163:U:H6	2.00	0.45	
1:C:89:G:H2'	1:C:90:C:H6	1.80	0.45	
3:B:441:SER:HB3	3:B:446:MSE:HB2	1.98	0.45	
2:D:130:A:C2	2:D:140:G:C6	3.05	0.45	
1:C:58:G:H8	1:C:58:G:O5'	2.00	0.44	
2:D:127:C:O2'	2:D:128:C:H5'	2.18	0.44	
2:D:156:G:H1	2:D:190:C:H42	1.64	0.44	
1:C:35:A:H1'	3:A:346:MSE:HE1	1.98	0.44	
2:D:190:C:O2'	2:D:191:C:H5'	2.17	0.44	
1:C:78:C:H2'	1:C:79:U:C6	2.52	0.44	
1:C:36:C:O2	3:A:387:ASP:HB2	2.17	0.44	
3:B:488:ILE:HD12	3:B:489:ILE:HG12	2.00	0.44	
1:C:61:G:H2'	1:C:62:G:O4'	2.18	0.44	
2:D:164:A:H8	2:D:164:A:O5'	2.00	0.44	
1:C:88:A:O2'	1:C:89:G:H5'	2.18	0.43	
3:A:364:LEU:HD12	3:A:364:LEU:O	2.18	0.43	
3:B:412:LEU:HD12	3:B:449:GLN:CA	2.48	0.43	
1:C:56:G:H2'	1:C:57:U:C5	2.54	0.43	
2:D:109:A:H2'	2:D:110:A:O4'	2.19	0.43	
2:D:145:A:H3'	2:D:146:U:C6	2.53	0.43	
1:C:10:A:H2'	1:C:11:A:C8	2.54	0.43	
2:D:146:U:H6	2:D:146:U:O5'	2.02	0.42	
3:A:321:LEU:HD23	3:A:340:VAL:CG1	2.48	0.42	
3:B:425:LEU:HD23	3:B:425:LEU:HA	1.90	0.42	
3:B:436:LEU:O	3:B:437:ASP:HB2	2.19	0.42	
2:D:186:G:H2'	2:D:187:C:H6	1.84	0.42	
3:A:327:ALA:HB1	3:B:471:PRO:CG	2.50	0.42	
1:C:15:A:O2'	1:C:16:U:H5'	2.19	0.42	
1:C:76:U:H2'	1:C:77:C:H6	1.77	0.42	
1:C:84:C:H2'	1:C:85:C:C6	2.54	0.42	
3:B:467:MSE:HA	3:B:467:MSE:HE2	2.01	0.42	
3:B:416:ILE:N	3:B:416:ILE:CD1	2.83	0.42	
2:D:141:G:O2'	2:D:142:G:H5'	2.20	0.41	



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:C:52:U:C2	1:C:54:A:N1	2.88	0.41
3:A:370:PHE:HB3	3:A:377:MSE:HE2	2.02	0.41
3:B:464:LEU:HG	3:B:479:ILE:HG22	2.02	0.41
3:B:488:ILE:HD12	3:B:489:ILE:N	2.35	0.41
3:B:473:TYR:O	3:B:474:ASP:HB2	2.21	0.41
2:D:122:C:H2'	2:D:123:A:H8	1.79	0.41
3:A:373:TYR:O	3:A:374:ASP:HB2	2.21	0.41
3:B:405:HIS:HD2	3:B:457:VAL:HG13	1.85	0.41
3:B:412:LEU:HA	6:B:606:HOH:O	2.20	0.41
1:C:85:C:H6	1:C:85:C:O5'	2.04	0.41
2:D:165:C:O2'	2:D:166:G:H5'	2.21	0.40
1:C:29:C:H2'	1:C:30:A:O4'	2.21	0.40
2:D:142:G:O2'	2:D:143:G:H5'	2.21	0.40
3:A:370:PHE:CD2	3:A:371:PRO:O	2.74	0.40
3:A:389:ILE:HD13	3:A:389:ILE:HA	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
3	A	90/98~(92%)	80 (89%)	8 (9%)	2 (2%)	6	22
3	В	89/98~(91%)	76 (85%)	13 (15%)	0	100	100
All	All	179/196 (91%)	156 (87%)	21 (12%)	2 (1%)	14	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	303	PRO
3	A	373	TYR



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	72/83 (87%)	69 (96%)	3 (4%)	30 63
3	В	69/83 (83%)	67 (97%)	2 (3%)	42 76
All	All	141/166 (85%)	136 (96%)	5 (4%)	36 70

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	309	ILE
3	A	358	SER
3	A	364	LEU
3	В	410	ASN
3	В	457	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	310	ASN
3	A	311	ASN
3	A	326	HIS
3	A	368	GLN
3	В	410	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	С	90/92~(97%)	8 (8%)	4 (4%)
2	D	91/92 (98%)	8 (8%)	5 (5%)
All	All	181/184 (98%)	16 (8%)	9 (4%)

All (16) RNA backbone outliers are listed below:



Mol	Chain	Res	Type
1	С	32	U
1	С	33	G
1	C C C C	40	G
1	С	52	U
1	С	53	U
1	С	54	A
1	С	55	G
1		74	A
2	D	118	U
2	D	133	G
2	D	135	A
2	D	140	G
2	D	147	U
2	D	152	U
2	D	153	U
2	D	155	G

All (9) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	С	32	U
1	С	51	G
1	С	52	U
1	С	53	U
2	D	132	U
2	D	134	С
2	D	151	G
2	D	153	U
2	D	175	A

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res Link		Bo	ond leng	ths	В	ond ang	les
MIOI	туре	Chain	rtes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	GTP	С	1	1,4	26,34,34	1.26	3 (11%)	33,54,54	2.57	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GTP	С	1	1,4	-	5/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\mathring{A})$	Ideal(Å)
1	С	1	GTP	C6-N1	3.54	1.39	1.33
1	С	1	GTP	O5'-C5'	-2.39	1.35	1.44
1	С	1	GTP	C8-N7	-2.13	1.30	1.34

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	С	1	GTP	C5-C6-N1	-8.46	111.86	123.43
1	С	1	GTP	C6-N1-C2	5.80	125.14	115.93
1	С	1	GTP	PB-O3B-PG	-5.54	113.82	132.83
1	С	1	GTP	PA-O3A-PB	-5.16	115.11	132.83
1	С	1	GTP	O5'-C5'-C4'	4.71	125.21	108.99
1	С	1	GTP	C2-N3-C4	-3.15	111.76	115.36
1	С	1	GTP	N3-C2-N1	-2.58	123.78	127.22

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	С	1	GTP	C5'-O5'-PA-O3A
1	С	1	GTP	C5'-O5'-PA-O1A
1	С	1	GTP	C5'-O5'-PA-O2A
1	С	1	GTP	O4'-C4'-C5'-O5'
1	С	1	GTP	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	С	1	GTP	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	С	91/92 (98%)	-0.34	0 100 100	58, 87, 140, 150	0
2	D	92/92 (100%)	-0.14	0 100 100	56, 117, 157, 167	0
3	A	88/98 (89%)	0.05	0 100 100	54, 73, 91, 109	0
3	В	88/98 (89%)	-0.10	0 100 100	66, 85, 103, 110	0
All	All	359/380 (94%)	-0.14	0 100 100	54, 84, 142, 167	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	GTP	С	1	32/32	0.92	0.17	69,79,106,110	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	MG	С	508	1/1	0.36	0.59	110,110,110,110	0
4	MG	С	509	1/1	0.40	0.35	95,95,95,95	0
4	MG	С	505	1/1	0.59	0.14	82,82,82,82	0
4	MG	С	507	1/1	0.72	0.20	68,68,68,68	0
4	MG	D	502	1/1	0.81	0.34	71,71,71,71	0
4	MG	С	503	1/1	0.91	0.22	72,72,72,72	0
4	MG	С	506	1/1	0.92	0.27	89,89,89,89	0
4	MG	D	504	1/1	0.95	0.08	66,66,66,66	0
5	K	С	511	1/1	0.96	0.20	71,71,71,71	0
4	MG	С	510	1/1	0.97	0.24	61,61,61,61	0
4	MG	С	501	1/1	0.97	0.31	63,63,63,63	0

6.5 Other polymers (i)

There are no such residues in this entry.

