

wwPDB X-ray Structure Validation Summary Report (i)

May 29, 2020 – 02:29 am BST

PDB ID	:	3DED
Title	:	C-terminal domain of Probable hemolysin from Chromobacterium violaceum
Authors	:	Chang, C.; Xu, X.; Cui, H.; Savchenko, A.; Edwards, A.; Joachimiak, A.;
		Midwest Center for Structural Genomics (MCSG)
Deposited on	:	2008-06-09
$\operatorname{Resolution}$:	2.14 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} {f Whole archive}\ (\#{f Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R _{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	113	%		000/			
1	Π	110	73% 2%	•	23%			
1	В	113	63%	12% •	24%			
1	С	113	<u>4%</u> 72%	8% •	19%			
-		110	%	070 •	1970			
1	D	113	68%	8%	24%			
1	Е	113	<u>4%</u> 66%	9% •	23%			
1	F	113	71%	5%	24%			



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4716 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	87	Total	С	Ν	0	Se	0	2	0
	A	01	713	450	125	135	3	0	2	0
1	В	86	Total	С	Ν	Ο	Se	0	3	0
	D	80	707	449	122	133	3	0	0	0
1	С	91	Total	С	Ν	Ο	Se	0	0	0
	U	91	725	455	127	140	3	0		0
1	D	86	Total	С	Ν	0	Se	0	0	0
	D	80	687	434	119	131	3	0		U
1	Е	87	Total	С	Ν	0	Se	0	2	0
		01	716	452	128	133	3	0	2	0
1	Б	86	Total	С	Ν	Ο	Se	0	2	0
	1 F	86	707	446	123	134	4		2	U

• Molecule 1 is a protein called Probable hemolysin.

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	315	MSE	-	expression tag	UNP Q7P1I2
А	316	GLY	-	expression tag	UNP Q7P1I2
А	317	SER	-	expression tag	UNP Q7P1I2
А	318	SER	-	expression tag	UNP Q7P1I2
А	319	HIS	-	expression tag	UNP Q7P1I2
А	320	HIS	-	expression tag	UNP Q7P1I2
А	321	HIS	-	expression tag	UNP Q7P1I2
A	322	HIS	-	expression tag	UNP Q7P1I2
А	323	HIS	-	expression tag	UNP Q7P1I2
A	324	HIS	-	expression tag	UNP Q7P1I2
А	325	SER	-	expression tag	UNP Q7P1I2
A	326	SER	-	expression tag	UNP Q7P1I2
А	327	GLY	-	expression tag	UNP Q7P1I2
А	328	ARG	-	expression tag	UNP Q7P1I2
А	329	GLU	-	expression tag	UNP Q7P1I2
А	330	ASN	-	expression tag	UNP Q7P1I2
А	331	LEU	-	expression tag	UNP Q7P1I2



Chain	hain Residue Modelled Actual		Comment	Reference	
A	332	TYR		expression tag	UNP Q7P1I2
A	333	PHE	_	expression tag	UNP Q7P1I2
A	334	GLN	_	expression tag	UNP Q7P1I2
A	335	GLY	_	expression tag	UNP Q7P1I2
A	336	HIS	_	expression tag	UNP Q7P1I2
В	315	MSE	_	expression tag	UNP Q7P1I2
В	316	GLY	-	expression tag	UNP Q7P1I2
В	317	SER	_	expression tag	UNP Q7P1I2
В	318	SER	-	expression tag	UNP Q7P1I2
В	319	HIS	-	expression tag	UNP Q7P1I2
В	320	HIS	-	expression tag	UNP Q7P1I2
В	321	HIS	-	expression tag	UNP Q7P1I2
В	322	HIS	-	expression tag	UNP Q7P1I2
В	323	HIS	-	expression tag	UNP Q7P1I2
В	324	HIS	-	expression tag	UNP Q7P1I2
В	325	SER	-	expression tag	UNP Q7P1I2
В	326	SER	-	expression tag	UNP Q7P1I2
В	327	GLY	-	expression tag	UNP Q7P1I2
В	328	ARG	-	expression tag	UNP Q7P1I2
В	329	GLU	-	expression tag	UNP Q7P1I2
В	330	ASN	-	expression tag	UNP Q7P1I2
В	331	LEU	-	expression tag	UNP Q7P1I2
В	332	TYR	-	expression tag	UNP Q7P1I2
В	333	PHE	-	expression tag	UNP Q7P1I2
В	334	GLN	-	expression tag	UNP Q7P1I2
В	335	GLY	-	expression tag	UNP Q7P1I2
В	336	HIS	-	expression tag	UNP Q7P1I2
С	315	MSE	-	expression tag	UNP Q7P1I2
С	316	GLY	-	expression tag	UNP Q7P1I2
С	317	SER	-	expression tag	UNP Q7P1I2
С	318	SER	-	expression tag	UNP Q7P1I2
С	319	HIS	-	expression tag	UNP Q7P1I2
С	320	HIS	_	expression tag	UNP Q7P1I2
С	321	HIS	-	expression tag	UNP Q7P1I2
С	322	HIS	-	expression tag	UNP Q7P1I2
С	323	HIS	-	expression tag	UNP Q7P1I2
С	324	HIS	-	expression tag	UNP Q7P1I2
С	325	SER	-	expression tag	UNP Q7P1I2
С	326	SER	-	expression tag	UNP Q7P1I2
С	327	GLY	-	expression tag	UNP Q7P1I2
С	328	ARG	-	expression tag	UNP Q7P1I2
	220	OT T		•	TIND OFDATA

Continued on next page...

UNP Q7P1I2



expression tag

GLU

-

С

329

Chain	Residue	Modelled	Actual	Comment	Reference
С	330	ASN	-	expression tag	UNP Q7P1I2
С	331	LEU	-	expression tag	UNP Q7P1I2
С	332	TYR	-	expression tag	UNP Q7P1I2
С	333	PHE	-	expression tag	UNP Q7P1I2
С	334	GLN	-	expression tag	UNP Q7P1I2
С	335	GLY	-	expression tag	UNP Q7P1I2
С	336	HIS	-	expression tag	UNP Q7P1I2
D	315	MSE	-	expression tag	UNP Q7P1I2
D	316	GLY	-	expression tag	UNP Q7P1I2
D	317	SER	-	expression tag	UNP Q7P1I2
D	318	SER	-	expression tag	UNP Q7P1I2
D	319	HIS	-	expression tag	UNP Q7P1I2
D	320	HIS	-	expression tag	UNP Q7P1I2
D	321	HIS	-	expression tag	UNP Q7P1I2
D	322	HIS	-	expression tag	UNP Q7P1I2
D	323	HIS	-	expression tag	UNP Q7P1I2
D	324	HIS	-	expression tag	UNP Q7P1I2
D	325	SER	-	expression tag	UNP Q7P1I2
D	326	SER	-	expression tag	UNP Q7P1I2
D	327	GLY	-	expression tag	UNP Q7P1I2
D	328	ARG	-	expression tag	UNP Q7P1I2
D	329	GLU	-	expression tag	UNP Q7P1I2
D	330	ASN	-	expression tag	UNP Q7P1I2
D	331	LEU	-	expression tag	UNP Q7P1I2
D	332	TYR	-	expression tag	UNP Q7P1I2
D	333	PHE	-	expression tag	UNP Q7P1I2
D	334	GLN	-	expression tag	UNP Q7P1I2
D	335	GLY	-	expression tag	UNP Q7P1I2
D	336	HIS	-	expression tag	UNP Q7P1I2
E	315	MSE	-	expression tag	UNP Q7P1I2
E	316	GLY	-	expression tag	UNP Q7P1I2
E	317	SER	-	expression tag	UNP Q7P1I2
E	318	SER	-	expression tag	UNP Q7P1I2
E	319	HIS	-	expression tag	UNP Q7P1I2
E	320	HIS	-	expression tag	UNP Q7P1I2
E	321	HIS	-	expression tag	UNP Q7P1I2
E	322	HIS	-	expression tag	UNP Q7P1I2
E	323	HIS	-	expression tag	UNP Q7P1I2
E	324	HIS	-	expression tag	UNP Q7P1I2
E	325	SER	-	expression tag	UNP Q7P1I2
E	326	SER	-	expression tag	UNP Q7P1I2
E	327	GLY	-	expression tag	UNP Q7P1I2

Continued from previous page...
Chain Besidue Modelled Actual Comment Beference



Chain	Residue	Modelled	Actual	Comment	Reference
Е	328	ARG	-	expression tag	UNP Q7P1I2
Е	329	GLU	_	expression tag	UNP Q7P1I2
Е	330	ASN	-	expression tag	UNP Q7P1I2
Е	331	LEU	-	expression tag	UNP Q7P1I2
Е	332	TYR	-	expression tag	UNP Q7P1I2
Е	333	PHE	-	expression tag	UNP Q7P1I2
Е	334	GLN	-	expression tag	UNP Q7P1I2
Е	335	GLY	-	expression tag	UNP Q7P1I2
Е	336	HIS	-	expression tag	UNP Q7P1I2
F	315	MSE	-	expression tag	UNP Q7P1I2
F	316	GLY	-	expression tag	UNP Q7P1I2
F	317	SER	-	expression tag	UNP Q7P1I2
F	318	SER	-	expression tag	UNP Q7P1I2
F	319	HIS	-	expression tag	UNP Q7P1I2
F	320	HIS	-	expression tag	UNP Q7P1I2
F	321	HIS	-	expression tag	UNP Q7P1I2
F	322	HIS	-	expression tag	UNP Q7P1I2
F	323	HIS	-	expression tag	UNP Q7P1I2
F	324	HIS	-	expression tag	UNP Q7P1I2
F	325	SER	-	expression tag	UNP Q7P1I2
F	326	SER	-	expression tag	UNP Q7P1I2
F	327	GLY	-	expression tag	UNP Q7P1I2
F	328	ARG	-	expression tag	UNP Q7P1I2
F	329	GLU	-	expression tag	UNP Q7P1I2
F	330	ASN	-	expression tag	UNP Q7P1I2
F	331	LEU	-	expression tag	UNP Q7P1I2
F	332	TYR	-	expression tag	UNP Q7P1I2
F	333	PHE	-	expression tag	UNP Q7P1I2
F	334	GLN	-	expression tag	UNP Q7P1I2
F	335	GLY	-	expression tag	UNP Q7P1I2
F	336	HIS	-	expression tag	UNP Q7P1I2

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Ca 1 1	0	0
2	Е	2	Total Ca 2 2	0	0
2	В	1	Total Ca 1 1	0	0
2	С	2	Total Ca 2 2	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Ca 2 2	0	0
2	F	1	Total Ca 1 1	0	0


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	86	Total O 86 86	0	0
3	В	74	Total O 74 74	0	0
3	С	54	$\begin{array}{cc} \text{Total} & \text{O} \\ 54 & 54 \end{array}$	0	0
3	D	66	Total O 66 66	0	0
3	Е	85	Total O 85 85	0	0
3	F	87	Total O 87 87	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Probable hemolysin



Chain F:	71%	5%	24%
MSE SER SER SER HIS HIS HIS HIS SER SER SER SER	SER ASC ASC ASC TTR ASC ASC CLU CLU CLU CLU CLU CLU CLU CLU CLU CL	H126 HIS	



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	66.14Å 55.31 Å 93.73 Å	Depositor
a, b, c, α , β , γ	90.00° 94.34° 90.00°	Depositor
Resolution (Å)	50.00 - 2.14	Depositor
	30.66 - 2.14	EDS
% Data completeness	97.8 (50.00-2.14)	Depositor
(in resolution range)	97.9(30.66-2.14)	EDS
R _{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.38 (at 2.14 \text{\AA})$	Xtriage
Refinement program	REFMAC $5.2.0019$	Depositor
R, R_{free}	0.186 , 0.235	Depositor
III, IIIfree	0.182 , 0.230	DCC
R_{free} test set	1847 reflections (5.00%)	wwPDB-VP
Wilson B-factor $(Å^2)$	27.8	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , 39.7	EDS
L-test for twinning ²	$ \langle L \rangle = 0.49, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4716	wwPDB-VP
Average B, all atoms $(Å^2)$	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles	
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.72	0/732	0.70	0/983
1	В	0.70	0/728	0.70	0/979
1	С	0.64	0/737	0.71	0/990
1	D	0.64	0/699	0.72	0/940
1	Е	0.77	0/735	0.73	0/986
1	F	0.75	0/722	0.70	0/969
All	All	0.70	0/4353	0.71	0/5847

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	713	0	678	3	0
1	В	707	0	682	11	0
1	С	725	0	677	9	0
1	D	687	0	647	6	0
1	Е	716	0	687	13	0
1	F	707	0	672	4	0
2	А	2	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	1	0	0	0	0
2	С	2	0	0	0	0
2	D	1	0	0	0	0
2	Е	2	0	0	0	0
2	F	1	0	0	0	0
3	А	86	0	0	1	0
3	В	74	0	0	1	0
3	С	54	0	0	1	0
3	D	66	0	0	1	0
3	Е	85	0	0	2	0
3	F	87	0	0	1	0
All	All	4716	0	4043	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:338:GLY:HA3	1:C:339:GLU:HB2	1.26	1.16
1:B:342[B]:GLU:HB2	1:B:357[B]:VAL:HG22	1.32	1.11
1:B:342[B]:GLU:HB2	1:B:357[B]:VAL:CG2	1.91	0.98
1:C:338:GLY:HA3	1:C:339:GLU:CB	1.97	0.87
1:C:338:GLY:CA	1:C:339:GLU:HB2	2.12	0.74

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	87/113~(77%)	84 (97%)	3~(3%)	0	100 100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	87/113~(77%)	83~(95%)	4 (5%)	0	100	100
1	С	89/113~(79%)	81 (91%)	7 (8%)	1 (1%)	14	7
1	D	84/113 (74%)	82 (98%)	2(2%)	0	100	100
1	Ε	87/113~(77%)	84 (97%)	3~(3%)	0	100	100
1	F	86/113~(76%)	83 (96%)	2(2%)	1 (1%)	13	6
All	All	520/678~(77%)	497~(96%)	21 (4%)	2(0%)	34	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	339	GLU
1	F	342	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	77/94~(82%)	77~(100%)	0	100 100
1	В	77/94~(82%)	73~(95%)	4(5%)	23 18
1	С	76/94~(81%)	75~(99%)	1 (1%)	69 73
1	D	73/94~(78%)	73~(100%)	0	100 100
1	Ε	77/94~(82%)	74~(96%)	3~(4%)	32 29
1	F	76/94~(81%)	75~(99%)	1 (1%)	69 73
All	All	456/564~(81%)	447~(98%)	9~(2%)	59 57

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	340	GLU
1	F	394	VAL
1	Е	393[A]	ARG
1	В	424	GLN



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type
1	Ε	363	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ $>$ 2	$OWAB(A^2)$	Q<0.9
1	А	84/113~(74%)	-0.04	1 (1%) 79 83	22, 29, 37, 41	0
1	В	83/113~(73%)	0.19	2 (2%) 59 65	21, 27, 38, 43	0
1	С	88/113~(77%)	0.42	5 (5%) 23 30	26, 33, 49, 55	0
1	D	83/113~(73%)	0.04	1 (1%) 79 83	26, 31, 40, 47	0
1	Ε	84/113~(74%)	0.30	5 (5%) 21 27	22, 28, 39, 49	0
1	F	83/113~(73%)	-0.01	0 100 100	19, 25, 38, 43	0
All	All	505/678~(74%)	0.15	14 (2%) 53 61	19, 29, 41, 55	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	RSRZ
1	С	338	GLY	4.5
1	С	339	GLU	3.7
1	С	337	ASP	3.4
1	С	378	GLY	2.9
1	Е	364	GLU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	CA	F	503	1/1	0.99	0.02	$24,\!24,\!24,\!24$	0
2	CA	D	509	1/1	0.99	0.03	$28,\!28,\!28,\!28$	0
2	CA	С	507	1/1	0.99	0.05	$23,\!23,\!23,\!23$	0
2	CA	Е	501	1/1	0.99	0.03	$24,\!24,\!24,\!24$	0
2	CA	Е	502	1/1	0.99	0.02	21,21,21,21	0
2	CA	С	508	1/1	0.99	0.03	$26,\!26,\!26,\!26$	0
2	CA	А	505	1/1	0.99	0.03	20,20,20,20	0
2	CA	В	506	1/1	1.00	0.02	22,22,22,22	0
2	CA	А	504	1/1	1.00	0.03	$20,\!20,\!20,\!20$	0

6.5 Other polymers (i)

There are no such residues in this entry.

