



wwPDB X-ray Structure Validation Summary Report ⓘ

May 4, 2022 – 08:24 PM EDT

PDB ID : 3E8K
Title : Crystal structure of HK97 Prohead II
Authors : Gertsman, I.; Speir, J.; Johnson, J.E.
Deposited on : 2008-08-20
Resolution : 3.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.28.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1037 (3.80-3.52)
Ramachandran outliers	138981	1004 (3.80-3.52)
Sidechain outliers	138945	1002 (3.80-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	273	30% 52% 11% • 6%
1	B	273	27% 49% 15% 8%
1	C	273	31% 51% 9% • 9%
1	D	273	31% 53% 8% • 7%
1	E	273	25% 56% 12% • 7%
1	F	273	23% 52% 14% • 9%
1	G	273	30% 49% 10% 10%

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 13590 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	256	1978	1238	346	384	10	0	0	0
1	B	250	1934	1209	339	377	9	0	0	0
1	C	248	1918	1198	337	375	8	0	0	0
1	D	254	1961	1228	343	381	9	0	0	0
1	E	255	1970	1233	345	383	9	0	0	0
1	F	248	1918	1198	337	375	8	0	0	0
1	G	247	1911	1193	336	374	8	0	0	0

There are 35 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	159	ALA	-	linker	UNP P49861
A	160	PRO	-	linker	UNP P49861
A	161	GLY	-	linker	UNP P49861
A	162	ASP	-	linker	UNP P49861
A	336	PHE	TRP	engineered mutation	UNP P49861
B	159	ALA	-	linker	UNP P49861
B	160	PRO	-	linker	UNP P49861
B	161	GLY	-	linker	UNP P49861
B	162	ASP	-	linker	UNP P49861
B	336	PHE	TRP	engineered mutation	UNP P49861
C	159	ALA	-	linker	UNP P49861
C	160	PRO	-	linker	UNP P49861
C	161	GLY	-	linker	UNP P49861
C	162	ASP	-	linker	UNP P49861
C	336	PHE	TRP	engineered mutation	UNP P49861

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	159	ALA	-	linker	UNP P49861
D	160	PRO	-	linker	UNP P49861
D	161	GLY	-	linker	UNP P49861
D	162	ASP	-	linker	UNP P49861
D	336	PHE	TRP	engineered mutation	UNP P49861
E	159	ALA	-	linker	UNP P49861
E	160	PRO	-	linker	UNP P49861
E	161	GLY	-	linker	UNP P49861
E	162	ASP	-	linker	UNP P49861
E	336	PHE	TRP	engineered mutation	UNP P49861
F	159	ALA	-	linker	UNP P49861
F	160	PRO	-	linker	UNP P49861
F	161	GLY	-	linker	UNP P49861
F	162	ASP	-	linker	UNP P49861
F	336	PHE	TRP	engineered mutation	UNP P49861
G	159	ALA	-	linker	UNP P49861
G	160	PRO	-	linker	UNP P49861
G	161	GLY	-	linker	UNP P49861
G	162	ASP	-	linker	UNP P49861
G	336	PHE	TRP	engineered mutation	UNP P49861

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major capsid protein

Chain A: 




- Molecule 1: Major capsid protein

Chain B: 




- Molecule 1: Major capsid protein

Chain C: 31% 51% 9% 9%




• Molecule 1: Major capsid protein

Chain D: 31% 53% 8% 7%




• Molecule 1: Major capsid protein

Chain E: 25% 56% 12% 7%







• Molecule 1: Major capsid protein



• Molecule 1: Major capsid protein



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	553.03Å 574.39Å 587.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 3.65 34.99 – 3.65	Depositor EDS
% Data completeness (in resolution range)	64.8 (35.00-3.65) 54.3 (34.99-3.65)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 3.66Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.366 , (Not available) 0.310 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	72.5	Xtrriage
Anisotropy	0.013	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , -38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.23$, $\langle L^2 \rangle = 0.09$	Xtrriage
Estimated twinning fraction	0.276 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.75	EDS
Total number of atoms	13590	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.41	1/2013 (0.0%)	0.68	2/2730 (0.1%)
1	B	0.42	0/1968	0.82	11/2669 (0.4%)
1	C	0.36	0/1952	0.68	2/2647 (0.1%)
1	D	0.37	1/1996 (0.1%)	0.68	5/2708 (0.2%)
1	E	0.40	1/2005 (0.0%)	0.70	5/2720 (0.2%)
1	F	0.42	0/1952	0.83	11/2647 (0.4%)
1	G	0.40	0/1944	0.77	9/2636 (0.3%)
All	All	0.40	3/13830 (0.0%)	0.74	45/18757 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	329	PHE	C-N	-5.66	1.21	1.34
1	A	212	MET	C-N	-5.28	1.21	1.34
1	D	288	LEU	C-N	-5.03	1.22	1.34

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	200	ALA	C-N-CD	-9.66	99.35	120.60
1	G	272	ALA	CB-CA-C	-8.25	97.73	110.10
1	G	228	GLY	N-CA-C	8.21	133.63	113.10
1	D	193	SER	N-CA-CB	-7.66	99.01	110.50
1	G	195	GLN	N-CA-C	-7.24	91.46	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	369	ALA	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1978	0	1946	311	0
1	B	1934	0	1896	343	0
1	C	1918	0	1878	343	0
1	D	1961	0	1929	330	0
1	E	1970	0	1937	365	0
1	F	1918	0	1878	347	0
1	G	1911	0	1870	285	0
All	All	13590	0	13334	2213	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 82.

The worst 5 of 2213 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:127:PRO:HB2	1:E:213:TYR:CD1	1.32	1.62
1:D:125:ILE:HD11	1:D:208:ASN:CG	1.32	1.49
1:D:125:ILE:HD11	1:D:208:ASN:ND2	1.26	1.45
1:C:196:VAL:HG12	1:C:203:LEU:CD1	1.44	1.45
1:B:201:PRO:O	1:B:204:GLN:CB	1.67	1.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/273 (93%)	195 (77%)	49 (19%)	10 (4%)	3	26
1	B	248/273 (91%)	189 (76%)	48 (19%)	11 (4%)	2	24
1	C	246/273 (90%)	188 (76%)	47 (19%)	11 (4%)	2	23
1	D	252/273 (92%)	209 (83%)	34 (14%)	9 (4%)	3	29
1	E	253/273 (93%)	190 (75%)	49 (19%)	14 (6%)	2	20
1	F	246/273 (90%)	192 (78%)	40 (16%)	14 (6%)	1	19
1	G	245/273 (90%)	199 (81%)	36 (15%)	10 (4%)	3	25
All	All	1744/1911 (91%)	1362 (78%)	303 (17%)	79 (4%)	2	23

5 of 79 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	124	ILE
1	A	208	ASN
1	B	200	ALA
1	B	297	PHE
1	C	131	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/223 (95%)	184 (87%)	27 (13%)	4	22
1	B	206/223 (92%)	181 (88%)	25 (12%)	5	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	204/223 (92%)	184 (90%)	20 (10%)	8	33
1	D	209/223 (94%)	191 (91%)	18 (9%)	10	39
1	E	210/223 (94%)	185 (88%)	25 (12%)	5	25
1	F	204/223 (92%)	173 (85%)	31 (15%)	3	17
1	G	203/223 (91%)	185 (91%)	18 (9%)	9	38
All	All	1447/1561 (93%)	1283 (89%)	164 (11%)	6	28

5 of 164 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	129	LEU
1	F	361	LEU
1	F	145	SER
1	F	212	MET
1	G	174	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	283	HIS
1	D	334	GLN
1	G	248	ASN
1	C	284	ASN
1	D	158	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.