



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2023 – 04:14 AM EDT

PDB ID : 3EOY
Title : Structure of Reovirus sigma1 in Complex with Its Receptor Junctional Adhesion Molecule-A
Authors : Kirchner, E.; Guglielmi, K.M.; Dermody, T.S.; Stehle, T.
Deposited on : 2008-09-29
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	
1	B	165	
1	C	165	
1	D	165	
1	E	165	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	165	<p>5% 59% 36% . .</p>
2	G	104	<p>12% 60% 34% 5% .</p>
2	H	104	<p>49% 46% . .</p>
2	I	104	<p>6% 44% 49% 5% .</p>
2	J	104	<p>5% 49% 45% . . .</p>
2	K	104	<p>% 54% 40% . .</p>
2	L	104	<p>2% 51% 43% . . .</p>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12227 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer capsid protein sigma-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	161	1241	794	209	233	5	0	0	0
1	B	160	1234	789	208	232	5	0	0	0
1	C	162	1247	797	210	235	5	0	0	0
1	D	162	1249	798	210	236	5	0	0	0
1	E	160	1234	789	208	232	5	0	0	0
1	F	160	1234	789	208	232	5	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	291	GLY	-	expression tag	UNP P03528
A	292	SER	-	expression tag	UNP P03528
A	408	THR	ALA	SEE REMARK 999	UNP P03528
B	291	GLY	-	expression tag	UNP P03528
B	292	SER	-	expression tag	UNP P03528
B	408	THR	ALA	SEE REMARK 999	UNP P03528
C	291	GLY	-	expression tag	UNP P03528
C	292	SER	-	expression tag	UNP P03528
C	408	THR	ALA	SEE REMARK 999	UNP P03528
D	291	GLY	-	expression tag	UNP P03528
D	292	SER	-	expression tag	UNP P03528
D	408	THR	ALA	SEE REMARK 999	UNP P03528
E	291	GLY	-	expression tag	UNP P03528
E	292	SER	-	expression tag	UNP P03528
E	408	THR	ALA	SEE REMARK 999	UNP P03528
F	291	GLY	-	expression tag	UNP P03528
F	292	SER	-	expression tag	UNP P03528

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	408	THR	ALA	SEE REMARK 999	UNP P03528

- Molecule 2 is a protein called Junctional adhesion molecule A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	102	Total	C	N	O	S	0	0	0
			798	500	132	162	4			
2	H	102	Total	C	N	O	S	0	0	0
			798	500	132	162	4			
2	I	102	Total	C	N	O	S	0	0	0
			798	500	132	162	4			
2	J	102	Total	C	N	O	S	0	0	0
			798	500	132	162	4			
2	K	102	Total	C	N	O	S	0	0	0
			798	500	132	162	4			
2	L	102	Total	C	N	O	S	0	0	0
			798	500	132	162	4			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	26	GLY	-	expression tag	UNP Q9Y624
G	27	SER	-	expression tag	UNP Q9Y624
H	26	GLY	-	expression tag	UNP Q9Y624
H	27	SER	-	expression tag	UNP Q9Y624
I	26	GLY	-	expression tag	UNP Q9Y624
I	27	SER	-	expression tag	UNP Q9Y624
J	26	GLY	-	expression tag	UNP Q9Y624
J	27	SER	-	expression tag	UNP Q9Y624
K	26	GLY	-	expression tag	UNP Q9Y624
K	27	SER	-	expression tag	UNP Q9Y624
L	26	GLY	-	expression tag	UNP Q9Y624
L	27	SER	-	expression tag	UNP Q9Y624


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Outer capsid protein sigma-1




- Molecule 1: Outer capsid protein sigma-1




- Molecule 1: Outer capsid protein sigma-1




- Molecule 1: Outer capsid protein sigma-1







• Molecule 1: Outer capsid protein sigma-1




• Molecule 1: Outer capsid protein sigma-1



• Molecule 2: Junctional adhesion molecule A




• Molecule 2: Junctional adhesion molecule A




• Molecule 2: Junctional adhesion molecule A







• Molecule 2: Junctional adhesion molecule A



• Molecule 2: Junctional adhesion molecule A



• Molecule 2: Junctional adhesion molecule A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.94Å 124.34Å 130.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.40 29.89 – 3.40	Depositor EDS
% Data completeness (in resolution range)	90.2 (30.00-3.40) 90.2 (29.89-3.40)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.95 (at 3.39Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.210 , 0.252 0.210 , 0.252	Depositor DCC
R_{free} test set	2187 reflections (9.96%)	wwPDB-VP
Wilson B-factor (Å ²)	64.1	Xtrriage
Anisotropy	0.304	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 27.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.010 for -h,l,k	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	12227	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.55	1/1277 (0.1%)	0.80	3/1742 (0.2%)
1	B	0.58	1/1269 (0.1%)	0.77	3/1731 (0.2%)
1	C	0.53	0/1283	0.71	1/1751 (0.1%)
1	D	0.63	1/1285 (0.1%)	0.87	5/1752 (0.3%)
1	E	0.50	0/1269	0.67	0/1731
1	F	0.51	0/1269	0.67	0/1731
2	G	0.45	0/814	0.65	0/1104
2	H	0.57	0/814	0.88	3/1104 (0.3%)
2	I	0.48	0/814	0.69	0/1104
2	J	0.65	1/814 (0.1%)	0.84	0/1104
2	K	0.51	0/814	0.69	0/1104
2	L	0.52	0/814	0.73	1/1104 (0.1%)
All	All	0.54	4/12536 (0.0%)	0.75	16/17062 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
2	J	0	2
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	435	SER	CA-CB	-6.08	1.43	1.52
1	B	406	GLY	CA-C	5.18	1.60	1.51
1	D	453	SER	CA-CB	-5.10	1.45	1.52
2	J	35	GLU	CG-CD	-5.10	1.44	1.51

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	435	SER	N-CA-CB	-11.84	92.75	110.50
1	B	407	GLY	N-CA-C	10.11	138.37	113.10
1	A	434	GLY	N-CA-C	-9.69	88.88	113.10
2	H	68	ASP	CB-CG-OD1	-9.20	110.02	118.30
1	D	453	SER	CB-CA-C	-8.83	93.32	110.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	433	GLY	Peptide
1	D	453	SER	Mainchain
2	J	115	GLY	Peptide
2	J	119	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1241	0	1193	53	0
1	B	1234	0	1185	53	0
1	C	1247	0	1197	39	0
1	D	1249	0	1200	84	0
1	E	1234	0	1185	52	0
1	F	1234	0	1185	53	0
2	G	798	0	775	45	0
2	H	798	0	775	44	0
2	I	798	0	775	54	0
2	J	798	0	775	49	0
2	K	798	0	775	43	0
2	L	798	0	775	56	0
All	All	12227	0	11795	569	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 569 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:57:SER:O	2:J:114:GLU:HB2	1.34	1.22
2:G:86:ARG:HG3	2:G:86:ARG:HH11	1.13	1.10
1:D:454:PHE:HD2	1:D:455:THR:N	1.54	1.05
1:D:376:PRO:HA	1:D:378:LEU:H	1.19	1.05
1:D:454:PHE:CD2	1:D:455:THR:N	2.36	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/165 (96%)	140 (88%)	18 (11%)	1 (1%)	25	57
1	B	158/165 (96%)	143 (90%)	14 (9%)	1 (1%)	25	57
1	C	160/165 (97%)	144 (90%)	15 (9%)	1 (1%)	25	57
1	D	160/165 (97%)	139 (87%)	19 (12%)	2 (1%)	12	39
1	E	158/165 (96%)	144 (91%)	13 (8%)	1 (1%)	25	57
1	F	158/165 (96%)	143 (90%)	14 (9%)	1 (1%)	25	57
2	G	100/104 (96%)	89 (89%)	10 (10%)	1 (1%)	15	46
2	H	100/104 (96%)	88 (88%)	12 (12%)	0	100	100
2	I	100/104 (96%)	90 (90%)	10 (10%)	0	100	100
2	J	100/104 (96%)	85 (85%)	13 (13%)	2 (2%)	7	30
2	K	100/104 (96%)	91 (91%)	9 (9%)	0	100	100
2	L	100/104 (96%)	89 (89%)	11 (11%)	0	100	100
All	All	1553/1614 (96%)	1385 (89%)	158 (10%)	10 (1%)	25	57

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	407	GLY
2	J	116	GLY
2	J	68	ASP
1	C	407	GLY
1	A	407	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/138 (98%)	127 (94%)	8 (6%)	19	49
1	B	134/138 (97%)	124 (92%)	10 (8%)	13	41
1	C	136/138 (99%)	128 (94%)	8 (6%)	19	49
1	D	136/138 (99%)	122 (90%)	14 (10%)	7	26
1	E	134/138 (97%)	127 (95%)	7 (5%)	23	53
1	F	134/138 (97%)	127 (95%)	7 (5%)	23	53
2	G	93/94 (99%)	87 (94%)	6 (6%)	17	46
2	H	93/94 (99%)	87 (94%)	6 (6%)	17	46
2	I	93/94 (99%)	85 (91%)	8 (9%)	10	35
2	J	93/94 (99%)	87 (94%)	6 (6%)	17	46
2	K	93/94 (99%)	87 (94%)	6 (6%)	17	46
2	L	93/94 (99%)	85 (91%)	8 (9%)	10	35
All	All	1367/1392 (98%)	1273 (93%)	94 (7%)	15	45

5 of 94 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	G	72	LEU
2	I	72	LEU
2	G	101	ARG
2	H	74	CYS
2	J	65	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	336	GLN
2	K	44	ASN
1	F	295	ASN
2	I	44	ASN
1	E	410	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	161/165 (97%)	-0.08	2 (1%) 79 77	32, 55, 76, 83	0
1	B	160/165 (96%)	-0.10	1 (0%) 89 89	34, 47, 66, 75	0
1	C	162/165 (98%)	-0.15	2 (1%) 79 77	36, 53, 75, 82	0
1	D	162/165 (98%)	0.06	5 (3%) 49 48	39, 59, 91, 95	0
1	E	160/165 (96%)	-0.03	0 100 100	34, 56, 72, 77	0
1	F	160/165 (96%)	0.10	8 (5%) 28 29	41, 60, 81, 88	0
2	G	102/104 (98%)	0.83	13 (12%) 3 4	70, 94, 111, 116	0
2	H	102/104 (98%)	-0.02	0 100 100	45, 61, 79, 85	0
2	I	102/104 (98%)	0.23	6 (5%) 22 23	39, 66, 88, 101	0
2	J	102/104 (98%)	0.30	5 (4%) 29 29	52, 73, 92, 104	0
2	K	102/104 (98%)	0.03	1 (0%) 82 81	48, 66, 79, 89	0
2	L	102/104 (98%)	0.30	2 (1%) 65 64	46, 69, 90, 94	0
All	All	1577/1614 (97%)	0.09	45 (2%) 51 50	32, 60, 92, 116	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	307	ILE	4.2
2	G	33	SER	3.7
1	D	294	PRO	3.5
2	G	56	SER	3.4
1	A	294	PRO	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.