



wwPDB EM Validation Summary Report ⓘ

Apr 11, 2024 – 04:39 PM EDT

PDB ID : 3EPF
EMDB ID : EMD-1563
Title : CryoEM structure of poliovirus receptor bound to poliovirus type 2
Authors : Zhang, P.; Mueller, S.; Morais, M.C.; Bator, C.M.; Bowman, V.D.; Hafenstein, S.; Wimmer, E.; Rossmann, M.G.
Deposited on : 2008-09-29
Resolution : 9.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 9.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	213	 36% 43% 16%
2	1	278	 73% 24%
3	2	262	 83% 15%
4	4	68	 78% 18%
5	3	235	 87% 10%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 8461 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poliovirus receptor.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	R	213	1638	1038	281	310	9	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	105	ASP	ASN	engineered mutation	UNP P15151
R	120	SER	ASN	engineered mutation	UNP P15151
R	188	GLN	ASN	engineered mutation	UNP P15151
R	218	GLN	ASN	engineered mutation	UNP P15151
R	237	SER	ASN	engineered mutation	UNP P15151

- Molecule 2 is a protein called Protein VP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	1	272	2129	1361	363	400	5	0	0

- Molecule 3 is a protein called Protein VP2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	2	262	2042	1298	346	384	14	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
2	11	VAL	ASP	conflict	UNP P06210


- Molecule 4 is a protein called Protein VP4.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	4	68	518	318	91	109	0	0

- Molecule 5 is a protein called Protein VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	3	235	1825	1156	303	349	17	0	0

- Molecule 6 is 1[2-CHLORO-4-METHOXY-PHENYL-OXYMETHYL]-4-[2,6-DICHLORO-PHENYL-OXYMETHYL]-BENZENE (three-letter code: SC4) (formula: C₂₁H₁₇Cl₃O₃).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	Cl	O	
6	1	1	27	21	3	3	0

- Molecule 7 is MYRISTIC ACID (three-letter code: MYR) (formula: C₁₄H₂₈O₂).



Mol	Chain	Residues	Atoms		AltConf
7	4	1	Total	C O	0
			11	10 1	


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		AltConf
8	1	136	Total	O	0
			136	136	
8	2	95	Total	O	0
			95	95	
8	4	25	Total	O	0
			25	25	
8	3	15	Total	O	0
			15	15	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Poliovirus receptor




- Molecule 2: Protein VP1





- Molecule 3: Protein VP2



● Molecule 4: Protein VP4

Chain 4:  78% 18%

● Molecule 5: Protein VP3

Chain 3:  87% 10%

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	Not provided	
Resolution determination method	Not provided	
CTF correction method	Not provided	
Microscope	FEI/PHILIPS CM300FEG/T	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	20	Depositor
Minimum defocus (nm)	1236	Depositor
Maximum defocus (nm)	2745	Depositor
Magnification	47000	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	28.879	Depositor
Minimum map value	-9.088	Depositor
Average map value	1.562	Depositor
Map value standard deviation	5.697	Depositor
Recommended contour level	7.26	Depositor
Map size (\AA)	577.22003, 577.22003, 577.22003	wwPDB
Map dimensions	217, 217, 217	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	2.66, 2.66, 2.66	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SC4, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	R	0.41	0/1678	0.77	4/2289 (0.2%)
2	1	0.43	0/2193	0.69	1/2996 (0.0%)
3	2	0.44	0/2098	0.72	1/2863 (0.0%)
4	4	0.52	0/527	0.68	0/713
5	3	0.43	0/1869	0.67	1/2549 (0.0%)
All	All	0.44	0/8365	0.71	7/11410 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	R	1	6
2	1	0	1
All	All	1	7

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	R	211	SER	O-C-N	-8.56	109.01	122.70
1	R	99	LEU	CA-CB-CG	6.25	129.67	115.30
2	1	288	ASP	N-CA-C	5.95	127.06	111.00
1	R	211	SER	CA-C-N	5.90	130.19	117.20
3	2	203	ASN	CB-CA-C	-5.76	98.89	110.40

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	R	143	ALA	CA

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	R	141	VAL	Peptide
1	R	142	LEU	Peptide
1	R	210	PRO	Peptide
1	R	211	SER	Peptide
1	R	212	SER	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	1638	0	1617	322	0
2	1	2129	0	2068	155	0
3	2	2042	0	1969	36	0
4	4	518	0	495	12	0
5	3	1825	0	1800	54	0
6	1	27	0	17	1	0
7	4	11	0	16	0	0
8	1	136	0	0	8	0
8	2	95	0	0	1	0
8	3	15	0	0	1	0
8	4	25	0	0	1	0
All	All	8461	0	7982	410	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 410 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:130:GLN:CG	2:1:106:SER:HA	1.13	1.55
1:R:130:GLN:CG	2:1:106:SER:CA	1.86	1.54
1:R:115:VAL:HG11	1:R:142:LEU:CD1	1.28	1.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:130:GLN:NE2	2:1:105:PHE:CE1	1.76	1.52
1:R:115:VAL:CG1	1:R:142:LEU:HD11	1.03	1.50

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	R	211/213 (99%)	137 (65%)	50 (24%)	24 (11%)	0	7
2	1	268/278 (96%)	250 (93%)	16 (6%)	2 (1%)	22	63
3	2	260/262 (99%)	245 (94%)	13 (5%)	2 (1%)	19	60
4	4	66/68 (97%)	62 (94%)	4 (6%)	0	100	100
5	3	233/235 (99%)	226 (97%)	7 (3%)	0	100	100
All	All	1038/1056 (98%)	920 (89%)	90 (9%)	28 (3%)	8	31

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	R	55	ASN
1	R	99	LEU
1	R	141	VAL
1	R	188	GLN
1	R	210	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	R	185/185 (100%)	149 (80%)	36 (20%)	1 8
2	1	231/235 (98%)	227 (98%)	4 (2%)	60 78
3	2	221/221 (100%)	208 (94%)	13 (6%)	19 45
4	4	56/56 (100%)	53 (95%)	3 (5%)	22 47
5	3	207/207 (100%)	196 (95%)	11 (5%)	22 47
All	All	900/904 (100%)	833 (93%)	67 (7%)	17 38

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	3	15	THR
5	3	99	MET
5	3	224	LEU
1	R	176	GLN
1	R	163	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
3	2	203	ASN
5	3	181	ASN
5	3	233	GLN
5	3	218	ASN
1	R	155	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	MYR	4	1	4	10,10,15	0.59	0	9,9,15	0.65	0
6	SC4	1	999	-	29,29,29	1.32	4 (13%)	39,39,39	1.38	5 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MYR	4	1	4	-	5/7/8/13	-
6	SC4	1	999	-	-	0/12/12/12	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	1	999	SC4	C5-C6	3.37	1.44	1.38
6	1	999	SC4	C15-C16	2.42	1.45	1.40
6	1	999	SC4	C3-C4	2.12	1.43	1.39
6	1	999	SC4	C18-C19	2.11	1.43	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	1	999	SC4	O1-C3-C4	4.45	121.89	116.40
6	1	999	SC4	C21-O3-C6	2.79	123.56	117.51
6	1	999	SC4	O1-C3-C2	-2.77	117.99	123.97
6	1	999	SC4	C15-C16-CL2	2.42	121.58	118.41
6	1	999	SC4	C15-C20-CL3	2.30	121.42	118.41

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	4	1	MYR	C1-C2-C3-C4
7	4	1	MYR	C6-C7-C8-C9
7	4	1	MYR	C5-C6-C7-C8
7	4	1	MYR	C4-C5-C6-C7
7	4	1	MYR	C7-C8-C9-C10

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	1	999	SC4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-1563. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 108



Y Index: 108




Z Index: 108


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 111



Y Index: 105




Z Index: 111


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 7.26. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 16660 nm³; this corresponds to an approximate mass of 15049 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.111\AA^{-1}

8 Fourier-Shell correlation


This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-1563 and PDB model 3EPF. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlays

9.1.1 Map-model overlay [i](#)




9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 7.26 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (7.26).













9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (7.26) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9810	 0.0880
1	 0.9950	 0.0850
2	 0.9840	 0.0910
3	 0.9970	 0.0920
4	 1.0000	 -0.0090
R	 0.9390	 0.1150

