



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 07:19 PM EDT

PDB ID : 3FJT
Title : Crystal structure of a human Fc fragment engineered for extended serum half-life
Authors : Oganessian, V.; Wu, H.; Dall'Acqua, W.F.
Deposited on : 2008-12-15
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	 9% 79% 18%
1	B	209	 10% 79% 19%
2	C	8	 38% 50% 12%
2	D	8	 50% 25% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	1	X	-	-	-
2	FUC	C	8	X	-	-	X
2	FUC	D	8	X	-	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	209	1675	1067	281	322	5	0	0	0
1	B	209	1675	1067	281	322	5	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	252	TYR	MET	engineered mutation	UNP P01857
A	254	THR	SER	engineered mutation	UNP P01857
A	256	GLU	THR	engineered mutation	UNP P01857
B	252	TYR	MET	engineered mutation	UNP P01857
B	254	THR	SER	engineered mutation	UNP P01857
B	256	GLU	THR	engineered mutation	UNP P01857

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	8	99	56	4	39	0	0	0
2	D	8	99	56	4	39	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	34	Total 34	O 34	0	0
3	B	34	Total 34	O 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ig gamma-1 chain C region



- Molecule 1: Ig gamma-1 chain C region




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.66Å 79.61Å 145.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.00 – 2.50 19.90 – 2.49	Depositor EDS
% Data completeness (in resolution range)	93.5 (12.00-2.50) 92.4 (19.90-2.49)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.50Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.227 , 0.291 0.258 , 0.270	Depositor DCC
R_{free} test set	1006 reflections (5.24%)	wwPDB-VP
Wilson B-factor (Å ²)	57.9	Xtrriage
Anisotropy	0.388	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3616	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/1722	0.70	1/2347 (0.0%)
1	B	0.62	0/1722	0.72	1/2347 (0.0%)
All	All	0.63	0/3444	0.71	2/4694 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	291	PRO	CA-N-CD	-8.24	99.96	111.50
1	A	252	TYR	CB-CG-CD2	-5.36	117.79	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1675	0	1633	25	0
1	B	1675	0	1634	27	0
2	C	99	0	85	2	0
2	D	99	0	85	1	0
3	A	34	0	0	1	0
3	B	34	0	0	1	0
All	All	3616	0	3437	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:291:PRO:O	1:B:292:ARG:CG	1.89	1.19
1:B:291:PRO:O	1:B:292:ARG:HG3	1.39	1.18
1:B:297:ASN:O	1:B:298:SER:HB2	1.73	0.88
1:A:297:ASN:O	1:A:298:SER:HB2	1.75	0.86
1:B:291:PRO:O	1:B:292:ARG:HG2	1.77	0.83
1:B:291:PRO:C	1:B:292:ARG:HG3	2.08	0.72
1:A:301:ARG:NE	2:C:2:NAG:H81	2.05	0.72
1:B:347:GLN:OE1	3:B:43:HOH:O	2.08	0.71
1:A:297:ASN:OD1	1:A:299:THR:OG1	2.11	0.68
1:B:271:PRO:HB2	1:B:292:ARG:NH1	2.09	0.68
1:A:273:VAL:CG1	1:A:323:VAL:HG13	2.24	0.66
1:B:368:LEU:HD13	1:B:407:TYR:CZ	2.31	0.66
1:B:297:ASN:O	1:B:298:SER:CB	2.44	0.63
1:A:263:VAL:HG22	1:A:323:VAL:HG21	1.86	0.58
1:A:263:VAL:CG2	1:A:323:VAL:HG21	2.36	0.56
1:A:297:ASN:O	1:A:298:SER:CB	2.51	0.54
1:B:277:TRP:CE3	1:B:306:LEU:HD22	2.42	0.54
1:B:328:LEU:HD13	1:B:332:ILE:HG13	1.90	0.53
1:B:252:TYR:OH	1:B:428:MET:HG2	2.08	0.53
1:B:328:LEU:HD21	1:B:332:ILE:HD11	1.91	0.53
1:B:271:PRO:HB2	1:B:292:ARG:HH12	1.74	0.51
1:B:346:PRO:HB3	1:B:372:PHE:HB3	1.93	0.50
1:A:301:ARG:HE	2:C:2:NAG:H81	1.75	0.50
1:B:429:HIS:CD2	1:B:431:ALA:H	2.28	0.50
1:A:273:VAL:HG13	1:A:323:VAL:HG13	1.93	0.50
1:B:325:ASN:OD1	1:B:326:LYS:N	2.45	0.49
2:D:2:NAG:H83	2:D:6:MAN:O4	2.13	0.49
1:B:368:LEU:HD13	1:B:407:TYR:CE2	2.48	0.48
1:A:365:LEU:HD13	1:A:441:LEU:HD23	1.95	0.48
1:A:398:LEU:HD11	1:A:402:GLY:HA2	1.96	0.48
1:A:271:PRO:HB2	1:A:292:ARG:HH12	1.78	0.47
1:A:253:ILE:HG13	3:A:45:HOH:O	2.14	0.46
1:A:354:SER:CB	1:B:349:TYR:HB3	2.46	0.46
1:B:266:VAL:HG21	1:B:302:VAL:HG23	1.99	0.45
1:B:358:LEU:HD23	1:B:363:VAL:HG11	1.97	0.45
1:B:240:VAL:HG12	1:B:241:PHE:N	2.32	0.44
1:B:291:PRO:HB2	1:B:292:ARG:H	1.42	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:GLU:OE2	1:A:335:THR:HG21	2.18	0.44
1:B:306:LEU:HG	1:B:307:THR:O	2.18	0.44
1:A:397:VAL:HB	1:A:405:PHE:CE2	2.53	0.44
1:A:290:LYS:HA	1:A:291:PRO:HD3	1.88	0.42
1:A:367:CYS:HB2	1:A:381:TRP:CZ2	2.55	0.42
1:A:377:ILE:HD11	1:A:427:VAL:CG1	2.50	0.42
1:B:318:GLU:HG2	1:B:335:THR:CG2	2.50	0.41
1:B:388:GLU:HB3	1:B:410:LEU:HD11	2.03	0.41
1:A:266:VAL:HG23	1:A:300:TYR:O	2.19	0.41
1:A:328:LEU:HD13	1:A:332:ILE:HG13	2.02	0.41
1:A:251:LEU:O	1:A:435:HIS:HD2	2.04	0.41
1:A:365:LEU:HD21	1:A:417:TRP:CE3	2.56	0.40
1:A:253:ILE:HG13	1:A:253:ILE:H	1.61	0.40
1:B:288:LYS:O	1:B:290:LYS:HD2	2.20	0.40
1:A:406:LEU:HD12	1:A:406:LEU:C	2.41	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	207/209 (99%)	193 (93%)	8 (4%)	6 (3%)	4 6
1	B	207/209 (99%)	194 (94%)	11 (5%)	2 (1%)	15 28
All	All	414/418 (99%)	387 (94%)	19 (5%)	8 (2%)	8 13

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	292	ARG
1	B	291	PRO
1	A	294	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	298	SER
1	A	291	PRO
1	A	297	ASN
1	A	329	PRO
1	B	292	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/194 (100%)	183 (94%)	11 (6%)	20	39
1	B	194/194 (100%)	182 (94%)	12 (6%)	18	35
All	All	388/388 (100%)	365 (94%)	23 (6%)	19	37

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	ILE
1	A	268	HIS
1	A	276	ASN
1	A	299	THR
1	A	356	ASP
1	A	359	THR
1	A	361	ASN
1	A	362	GLN
1	A	386	GLN
1	A	390	ASN
1	A	406	LEU
1	B	253	ILE
1	B	268	HIS
1	B	270	ASP
1	B	299	THR
1	B	307	THR
1	B	309	LEU
1	B	335	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	336	ILE
1	B	359	THR
1	B	362	GLN
1	B	442	SER
1	B	444	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	ASN
1	A	362	GLN
1	A	390	ASN
1	A	418	GLN
1	A	429	HIS
1	A	433	HIS
1	A	435	HIS
1	A	438	GLN
1	B	362	GLN
1	B	390	ASN
1	B	429	HIS
1	B	438	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.60	0	17,19,21	1.35	2 (11%)
2	NAG	C	2	2	14,14,15	0.64	0	17,19,21	1.03	1 (5%)
2	BMA	C	3	2	11,11,12	0.31	0	15,15,17	1.55	3 (20%)
2	MAN	C	4	2	11,11,12	0.64	0	15,15,17	1.04	1 (6%)
2	NAG	C	5	2	14,14,15	0.49	0	17,19,21	1.30	2 (11%)
2	MAN	C	6	2	11,11,12	0.57	0	15,15,17	0.94	0
2	NAG	C	7	2	14,14,15	0.54	0	17,19,21	0.61	0
2	FUC	C	8	2	10,10,11	0.65	0	14,14,16	0.76	0
2	NAG	D	1	1,2	14,14,15	0.39	0	17,19,21	1.62	4 (23%)
2	NAG	D	2	2	14,14,15	0.52	0	17,19,21	0.99	1 (5%)
2	BMA	D	3	2	11,11,12	0.23	0	15,15,17	0.64	0
2	MAN	D	4	2	11,11,12	0.58	0	15,15,17	0.85	0
2	NAG	D	5	2	14,14,15	0.62	0	17,19,21	1.56	2 (11%)
2	MAN	D	6	2	11,11,12	0.45	0	15,15,17	1.19	1 (6%)
2	NAG	D	7	2	14,14,15	0.65	0	17,19,21	1.08	0
2	FUC	D	8	2	10,10,11	0.67	0	14,14,16	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	1/1/5/7	1/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	C	5	2	-	3/6/23/26	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1
2	NAG	C	7	2	-	4/6/23/26	0/1/1/1
2	FUC	C	8	2	1/1/4/5	-	0/1/1/1
2	NAG	D	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	2/2/19/22	0/1/1/1
2	NAG	D	5	2	-	3/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	D	6	2	-	0/2/19/22	0/1/1/1
2	NAG	D	7	2	-	2/6/23/26	0/1/1/1
2	FUC	D	8	2	1/1/4/5	-	0/1/1/1

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	5	NAG	C4-C3-C2	4.14	117.09	111.02
2	D	5	NAG	C3-C4-C5	3.99	117.36	110.24
2	C	3	BMA	C1-O5-C5	3.53	116.98	112.19
2	C	5	NAG	C1-O5-C5	3.49	116.92	112.19
2	D	6	MAN	C1-O5-C5	3.43	116.84	112.19
2	D	1	NAG	O5-C5-C6	3.38	112.50	107.20
2	C	3	BMA	C1-C2-C3	3.31	113.73	109.67
2	C	1	NAG	O5-C1-C2	-3.13	106.34	111.29
2	D	1	NAG	O5-C1-C2	-3.06	106.45	111.29
2	C	1	NAG	O5-C5-C6	2.94	111.82	107.20
2	C	5	NAG	C3-C4-C5	2.49	114.68	110.24
2	D	1	NAG	C1-O5-C5	2.32	115.34	112.19
2	C	3	BMA	O5-C1-C2	2.29	114.31	110.77
2	D	2	NAG	C1-C2-N2	2.27	114.36	110.49
2	C	4	MAN	C1-O5-C5	2.12	115.06	112.19
2	D	1	NAG	C4-C3-C2	-2.10	107.94	111.02
2	C	2	NAG	C1-C2-N2	2.01	113.92	110.49

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	1	NAG	C1
2	C	8	FUC	C1
2	D	8	FUC	C1

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	5	NAG	C8-C7-N2-C2
2	C	5	NAG	O7-C7-N2-C2
2	C	7	NAG	C8-C7-N2-C2
2	C	7	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	1	NAG	O7-C7-N2-C2
2	D	5	NAG	C8-C7-N2-C2
2	D	5	NAG	O7-C7-N2-C2
2	D	7	NAG	C8-C7-N2-C2
2	D	7	NAG	O7-C7-N2-C2
2	D	1	NAG	O5-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	D	4	MAN	O5-C5-C6-O6
2	D	4	MAN	C4-C5-C6-O6
2	C	7	NAG	O5-C5-C6-O6
2	C	7	NAG	C4-C5-C6-O6
2	D	5	NAG	O5-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	C	5	NAG	O5-C5-C6-O6
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	NAG	2	0
2	D	6	MAN	1	0
2	D	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	209/209 (100%)	0.44	19 (9%) 9 9	35, 55, 97, 113	0
1	B	209/209 (100%)	0.43	20 (9%) 8 7	38, 57, 97, 107	0
All	All	418/418 (100%)	0.43	39 (9%) 8 8	35, 57, 98, 113	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	299	THR	5.6
1	B	296	TYR	5.2
1	B	236	GLY	4.9
1	B	263	VAL	4.1
1	A	300	TYR	4.1
1	A	444	SER	4.1
1	B	444	SER	4.0
1	B	327	ALA	4.0
1	B	262	VAL	3.7
1	B	274	LYS	3.6
1	A	389	ASN	3.6
1	A	326	LYS	3.3
1	B	268	HIS	3.2
1	B	323	VAL	3.1
1	B	328	LEU	3.0
1	A	293	GLU	2.9
1	A	328	LEU	2.8
1	B	300	TYR	2.8
1	A	236	GLY	2.8
1	B	269	GLU	2.8
1	B	266	VAL	2.7
1	A	327	ALA	2.7
1	A	330	ALA	2.6
1	A	384	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	386	GLN	2.5
1	B	298	SER	2.5
1	B	329	PRO	2.4
1	A	298	SER	2.4
1	B	271	PRO	2.3
1	A	271	PRO	2.3
1	A	272	GLU	2.3
1	B	303	VAL	2.3
1	A	292	ARG	2.2
1	A	324	SER	2.2
1	A	369	VAL	2.2
1	B	282	VAL	2.2
1	A	262	VAL	2.1
1	B	326	LYS	2.1
1	B	293	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	7	14/15	0.67	0.22	69,75,79,79	0
2	FUC	C	8	10/11	0.73	0.43	121,122,122,122	0
2	NAG	C	5	14/15	0.74	0.35	106,107,107,107	0
2	FUC	D	8	10/11	0.77	0.37	117,118,118,119	0
2	NAG	C	1	14/15	0.78	0.21	112,114,117,120	0
2	NAG	D	5	14/15	0.78	0.32	97,99,100,100	0
2	NAG	D	1	14/15	0.79	0.22	107,108,112,115	0
2	NAG	D	2	14/15	0.81	0.20	100,103,105,106	0
2	NAG	C	7	14/15	0.83	0.17	96,98,99,99	0
2	BMA	D	3	11/12	0.86	0.17	88,94,96,97	0
2	NAG	C	2	14/15	0.86	0.21	105,107,108,109	0
2	MAN	D	4	11/12	0.87	0.27	96,96,97,98	0
2	BMA	C	3	11/12	0.88	0.15	103,103,104,104	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MAN	D	6	11/12	0.88	0.12	81,83,84,85	0
2	MAN	C	6	11/12	0.89	0.18	100,102,103,103	0
2	MAN	C	4	11/12	0.90	0.31	104,105,105,106	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.