



# Full wwPDB X-ray Structure Validation Report i

Feb 11, 2024 – 10:17 PM EST

PDB ID : 3GFZ  
Title : Klebsiella pneumoniae BlrP1 pH 6 manganese/cy-diGMP complex  
Authors : Barends, T.; Hartmann, E.; Griese, J.; Beitlich, T.; Kirienko, N.; Ryjenkov, D.; Reinstein, J.; Shoeman, R.; Gomelsky, M.; Schlichting, I.  
Deposited on : 2009-02-27  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	C2E	A	501	X	-	-	-
2	C2E	B	501	X	-	-	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6678 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Klebsiella pneumoniae BlrP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	393	3080	1973	533	556	18	0	0	0
1	B	394	3094	1983	534	559	18	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	ILE	-	expression tag	UNP A6T8V8
A	-6	SER	-	expression tag	UNP A6T8V8
A	-5	GLU	-	expression tag	UNP A6T8V8
A	-4	PHE	-	expression tag	UNP A6T8V8
A	-3	GLY	-	expression tag	UNP A6T8V8
A	-2	SER	-	expression tag	UNP A6T8V8
A	-1	SER	-	expression tag	UNP A6T8V8
A	0	ARG	-	expression tag	UNP A6T8V8
B	-7	ILE	-	expression tag	UNP A6T8V8
B	-6	SER	-	expression tag	UNP A6T8V8
B	-5	GLU	-	expression tag	UNP A6T8V8
B	-4	PHE	-	expression tag	UNP A6T8V8
B	-3	GLY	-	expression tag	UNP A6T8V8
B	-2	SER	-	expression tag	UNP A6T8V8
B	-1	SER	-	expression tag	UNP A6T8V8
B	0	ARG	-	expression tag	UNP A6T8V8

- Molecule 2 is 9,9'-[ $(2R,3R,3aS,5S,7aR,9R,10R,10aS,12S,14aR)-3,5,10,12\text{-tetrahydroxy-}5,12\text{-dioxidoctahydro-}2H,7H\text{-difuro}[3,2-d:3',2'-j][1,3,7,9,2,8]\text{tetraoxadiphosphacyclododecine-}2,9\text{-diyl}]$ bis(2-amino-1,9-dihydro-6H-purin-6-one) (three-letter code: C2E) (formula:  $C_{20}H_{24}N_{10}O_{14}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total C N O P					0	0
			46 20 10 14 2						
2	B	1	Total C N O P					0	0
			46 20 10 14 2						

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total Mn		0	0
			2 2			
3	B	2	Total Mn		0	0
			2 2			

- Molecule 4 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total C N O P					0	0
			31 17 4 9 1						
4	B	1	Total C N O P					0	0
			31 17 4 9 1						


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	208	Total O 208 208		0	0
5	B	138	Total O 138 138		0	0


### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Klebsiella pneumoniae BlrP1



- Molecule 1: Klebsiella pneumoniae BlrP1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.45 Å    97.09 Å    127.17 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	48.54 – 2.05 48.55 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.1 (48.54-2.05) 97.1 (48.55-2.05)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.77 (at 2.05 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
$R$ , $R_{free}$	0.229 , 0.272 0.232 , 0.275	Depositor DCC
$R_{free}$ test set	2618 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.4	Xtriage
Anisotropy	0.522	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 39.5	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.47$ , $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6678	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MN, FMN, C2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3148	0.65	1/4266 (0.0%)
1	B	0.39	0/3163	0.61	0/4287
All	All	0.40	0/6311	0.63	1/8553 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
All	All	0	5

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	316	ARG	CB-CA-C	-5.16	100.08	110.40

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	308	TYR	Peptide
1	A	309	SER	Peptide
1	A	316	ARG	Peptide
1	B	89	GLY	Peptide
1	B	90	THR	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3080	0	3100	13	0
1	B	3094	0	3110	20	0
2	A	46	0	17	1	0
2	B	46	0	18	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	31	0	19	0	0
4	B	31	0	19	0	0
5	A	208	0	0	0	0
5	B	138	0	0	0	0
All	All	6678	0	6283	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:THR:HG23	1:B:91:GLY:C	1.95	0.87
1:B:90:THR:HG23	1:B:91:GLY:O	1.83	0.77
1:A:310:GLY:O	1:A:313:LEU:N	2.27	0.64
1:B:314:LEU:O	1:B:318:GLN:NE2	2.36	0.59
1:B:89:GLY:O	1:B:90:THR:CB	2.54	0.56
1:A:311:LEU:HD23	1:B:341:ILE:HG23	1.87	0.56
1:B:149:ARG:O	1:B:400:ARG:N	2.39	0.55
1:A:24:VAL:HG21	1:A:94:ILE:HB	1.91	0.52
1:B:28:SER:HB3	1:B:91:GLY:HA2	1.92	0.52
1:B:41:LEU:HB3	1:B:97:LEU:HD21	1.92	0.51
1:A:317:PHE:N	1:A:317:PHE:CD1	2.80	0.50
1:B:89:GLY:O	1:B:90:THR:OG1	2.29	0.50
1:B:246:TYR:CD2	1:B:284:GLN:HB2	2.48	0.49
1:B:275:GLU:O	1:B:279:ILE:HG22	2.13	0.49
1:A:272:GLU:OE1	1:A:323:LYS:NZ	2.41	0.48
1:B:161:PRO:HG3	1:B:387:CYS:SG	2.54	0.47
1:A:39:ILE:HD11	1:A:129:LEU:HD21	1.97	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:SER:CB	1:B:91:GLY:HA2	2.47	0.44
1:A:304:PHE:CE1	1:A:310:GLY:HA3	2.52	0.44
1:A:174:GLN:OE1	2:A:501:C2E:H8	2.17	0.43
1:A:312:SER:HB3	1:B:307:GLY:O	2.18	0.43
1:B:246:TYR:CE2	1:B:284:GLN:HB2	2.54	0.42
1:A:308:TYR:HA	1:B:312:SER:OG	2.18	0.42
1:A:311:LEU:HD13	1:B:304:PHE:CE2	2.55	0.42
1:B:174:GLN:OE1	2:B:501:C2E:H8	2.19	0.42
1:A:275:GLU:CD	1:A:309:SER:HA	2.40	0.41
1:A:322:ILE:HD11	1:A:345:VAL:CG1	2.50	0.41
1:B:90:THR:CG2	1:B:91:GLY:C	2.78	0.40
1:B:90:THR:HA	1:B:91:GLY:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	387/413 (94%)	378 (98%)	7 (2%)	2 (0%)	29 18
1	B	390/413 (94%)	381 (98%)	6 (2%)	3 (1%)	19 10
All	All	777/826 (94%)	759 (98%)	13 (2%)	5 (1%)	25 15

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	90	THR
1	A	311	LEU
1	A	309	SER
1	B	102	THR
1	B	231	LEU

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	330/346 (95%)	325 (98%)	5 (2%)	65 62
1	B	332/346 (96%)	327 (98%)	5 (2%)	65 62
All	All	662/692 (96%)	652 (98%)	10 (2%)	65 62

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	HIS
1	A	80	ASP
1	A	173	LEU
1	A	312	SER
1	A	316	ARG
1	B	25	HIS
1	B	71	HIS
1	B	256	MET
1	B	270	LEU
1	B	298	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	66	GLN
1	B	268	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	C2E	B	501	3	44,52,52	3.20	11 (25%)	52,82,82	1.94	18 (34%)
4	FMN	B	504	-	33,33,33	1.03	2 (6%)	48,50,50	1.34	8 (16%)
4	FMN	A	504	-	33,33,33	1.04	2 (6%)	48,50,50	1.48	10 (20%)
2	C2E	A	501	3	44,52,52	3.27	12 (27%)	52,82,82	1.76	13 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C2E	B	501	3	1/1/10/10	6/22/62/62	0/6/7/7
4	FMN	B	504	-	-	2/18/18/18	0/3/3/3
4	FMN	A	504	-	-	3/18/18/18	0/3/3/3
2	C2E	A	501	3	1/1/10/10	5/22/62/62	0/6/7/7

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	C2E	C2'-C1'	-12.99	1.34	1.53
2	B	501	C2E	C2'-C1'	-12.97	1.34	1.53
2	A	501	C2E	C2A-C1A	-12.75	1.34	1.53
2	B	501	C2E	C2A-C1A	-12.22	1.35	1.53
2	A	501	C2E	O4'-C1'	5.04	1.48	1.41
2	B	501	C2E	O4'-C1'	4.64	1.47	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	C2E	C51-C61	-4.41	1.38	1.47
4	A	504	FMN	C4A-N5	4.13	1.38	1.30
2	A	501	C2E	C51-C61	-4.07	1.39	1.47
2	B	501	C2E	C5-C6	-3.99	1.39	1.47
2	A	501	C2E	O2'-C2'	-3.95	1.33	1.43
2	A	501	C2E	C5-C6	-3.85	1.39	1.47
2	B	501	C2E	O2'-C2'	-3.84	1.33	1.43
2	B	501	C2E	O2A-C2A	-3.78	1.34	1.43
4	B	504	FMN	C4A-N5	3.72	1.38	1.30
2	A	501	C2E	O2A-C2A	-3.61	1.34	1.43
2	B	501	C2E	C8-N7	-3.27	1.29	1.35
2	A	501	C2E	C8-N7	-2.77	1.30	1.35
2	B	501	C2E	O4A-C1A	2.64	1.44	1.41
4	B	504	FMN	C10-N1	2.56	1.38	1.33
2	A	501	C2E	C6-N1	-2.54	1.34	1.37
2	A	501	C2E	C2'-C3'	-2.53	1.47	1.52
2	A	501	C2E	P11-O11	2.39	1.59	1.50
2	A	501	C2E	O4A-C1A	2.34	1.44	1.41
2	B	501	C2E	P11-O11	2.33	1.59	1.50
2	B	501	C2E	C2'-C3'	-2.18	1.48	1.52
4	A	504	FMN	C10-N1	2.16	1.37	1.33

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	C2E	O4A-C1A-C2A	4.92	114.11	106.93
2	A	501	C2E	O4A-C1A-C2A	4.27	113.16	106.93
2	A	501	C2E	O2'-C2'-C1'	3.90	125.26	110.85
2	B	501	C2E	C8-N7-C5	3.88	110.38	102.99
2	A	501	C2E	C8-N7-C5	3.73	110.10	102.99
4	A	504	FMN	C4A-C10-N10	3.72	121.92	116.48
2	B	501	C2E	O2'-C2'-C1'	3.70	124.51	110.85
4	A	504	FMN	C4-N3-C2	-3.62	118.94	125.64
2	A	501	C2E	C3'-C2'-C1'	3.41	107.45	99.89
2	B	501	C2E	C3'-C2'-C1'	3.32	107.24	99.89
2	B	501	C2E	O5A-C5A-C4A	-3.20	97.99	108.99
2	A	501	C2E	C81-N71-C51	3.17	109.03	102.99
4	B	504	FMN	C4-N3-C2	-3.13	119.86	125.64
2	B	501	C2E	O61-C61-C51	-3.13	118.26	124.37
2	A	501	C2E	O21-P11-O5A	3.10	122.16	107.75
2	B	501	C2E	C51-C61-N11	3.09	119.41	113.95
4	A	504	FMN	C4A-C4-N3	2.93	120.64	113.19

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	504	FMN	C4A-C10-N10	2.93	120.76	116.48
2	A	501	C2E	C51-C61-N11	2.83	118.95	113.95
4	A	504	FMN	C10-C4A-N5	-2.77	118.98	124.86
2	B	501	C2E	C5'-C4'-C3'	-2.65	105.61	114.40
4	B	504	FMN	C10-C4A-N5	-2.61	119.31	124.86
2	A	501	C2E	C21-N11-C61	-2.61	120.29	125.10
4	B	504	FMN	C4A-C4-N3	2.59	119.75	113.19
2	A	501	C2E	C5'-C4'-C3'	-2.55	105.95	114.40
2	B	501	C2E	C5-C6-N1	2.53	118.42	113.95
2	B	501	C2E	N21-C21-N11	2.51	122.05	116.71
2	B	501	C2E	C21-N11-C61	-2.50	120.50	125.10
2	B	501	C2E	O21-P11-O5A	2.46	119.16	107.75
4	A	504	FMN	C4A-C10-N1	-2.41	119.13	124.73
2	A	501	C2E	O5'-P1-O1P	-2.37	99.80	109.07
2	B	501	C2E	O3A-P1-O1P	-2.37	100.59	109.47
2	A	501	C2E	C5-C6-N1	2.36	118.13	113.95
2	B	501	C2E	C81-N71-C51	2.33	107.44	102.99
4	A	504	FMN	C4-C4A-N5	2.31	121.53	118.23
2	A	501	C2E	O61-C61-C51	-2.28	119.91	124.37
2	A	501	C2E	O4'-C4'-C5'	2.26	116.82	109.37
2	B	501	C2E	O4'-C4'-C5'	2.26	116.81	109.37
4	B	504	FMN	O4-C4-C4A	-2.25	120.64	126.60
2	B	501	C2E	C2-N1-C6	-2.19	121.06	125.10
4	A	504	FMN	C10-N1-C2	2.19	121.28	116.90
4	B	504	FMN	C9A-C5A-N5	-2.18	120.06	122.43
4	A	504	FMN	O2-C2-N1	-2.17	118.24	121.83
2	B	501	C2E	O3A-C3A-C2A	2.13	119.42	111.68
4	B	504	FMN	C4A-C10-N1	-2.13	119.79	124.73
4	A	504	FMN	O4-C4-C4A	-2.09	121.05	126.60
4	A	504	FMN	C9A-C5A-N5	-2.08	120.17	122.43
2	B	501	C2E	O5A-P11-O11	-2.08	100.95	109.07
4	B	504	FMN	C4-C4A-N5	2.08	121.19	118.23

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	501	C2E	C1A
2	B	501	C2E	C1A

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	504	FMN	C5'-O5'-P-O2P
4	B	504	FMN	C5'-O5'-P-O3P
2	B	501	C2E	C2A-C3A-O3A-P1
2	B	501	C2E	C4A-C3A-O3A-P1
2	B	501	C2E	C4'-C5'-O5'-P1
4	A	504	FMN	C4'-C5'-O5'-P
2	A	501	C2E	C2A-C3A-O3A-P1
2	A	501	C2E	C3A-O3A-P1-O5'
2	A	501	C2E	C4'-C5'-O5'-P1
2	B	501	C2E	C3A-O3A-P1-O5'
2	A	501	C2E	C4A-C3A-O3A-P1
4	A	504	FMN	C2'-C3'-C4'-O4'
2	B	501	C2E	C3A-O3A-P1-O2P
2	A	501	C2E	O4A-C4A-C5A-O5A
2	B	501	C2E	C5'-O5'-P1-O1P
4	A	504	FMN	O3'-C3'-C4'-O4'


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	C2E	1	0
2	A	501	C2E	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	393/413 (95%)	1.12	52 (13%) <span style="border: 1px solid red; padding: 2px;">3</span> <span style="border: 1px solid red; padding: 2px;">2</span>	35, 41, 50, 55	0
1	B	394/413 (95%)	1.16	62 (15%) <span style="border: 1px solid red; padding: 2px;">2</span> <span style="border: 1px solid red; padding: 2px;">1</span>	34, 41, 49, 53	0
All	All	787/826 (95%)	1.14	114 (14%) <span style="border: 1px solid red; padding: 2px;">2</span> <span style="border: 1px solid red; padding: 2px;">2</span>	34, 41, 50, 55	0

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308	TYR	10.0
1	B	279	ILE	9.2
1	B	308	TYR	8.6
1	A	281	CYS	8.6
1	A	282	PHE	8.3
1	A	279	ILE	7.9
1	B	284	GLN	7.0
1	A	247	HIS	7.0
1	B	311	LEU	6.9
1	B	282	PHE	6.4
1	A	312	SER	6.1
1	B	102	THR	5.9
1	B	14	ASP	5.6
1	A	311	LEU	5.3
1	B	316	ARG	5.0
1	B	231	LEU	4.7
1	B	232	GLY	4.4
1	B	161	PRO	4.4
1	B	264	LEU	4.4
1	A	248	HIS	4.4
1	B	285	PHE	4.3
1	B	249	PRO	4.2
1	B	313	LEU	4.2
1	B	233	LYS	4.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	18	VAL	3.9
1	B	286	ARG	3.8
1	A	159	ALA	3.8
1	B	153	MET	3.7
1	A	280	THR	3.7
1	A	180	ALA	3.6
1	B	90	THR	3.6
1	B	229	LEU	3.6
1	A	278	VAL	3.6
1	A	26	ARG	3.5
1	B	314	LEU	3.5
1	A	233	LYS	3.5
1	A	1	MET	3.4
1	B	315	THR	3.4
1	B	230	PRO	3.2
1	A	114	THR	3.2
1	A	322	ILE	3.1
1	B	27	ALA	3.1
1	B	112	PHE	3.1
1	B	280	THR	3.1
1	B	312	SER	3.1
1	A	342	VAL	3.0
1	B	398	VAL	3.0
1	A	310	GLY	3.0
1	B	354	ILE	2.9
1	A	67	SER	2.9
1	B	261	ALA	2.9
1	A	341	ILE	2.9
1	A	14	ASP	2.8
1	A	317	PHE	2.8
1	B	319	PRO	2.8
1	B	256	MET	2.7
1	A	316	ARG	2.7
1	B	287	LYS	2.7
1	A	309	SER	2.7
1	A	285	PHE	2.7
1	B	163	HIS	2.7
1	B	262	ALA	2.7
1	B	306	ALA	2.7
1	A	393	GLU	2.7
1	A	346	VAL	2.6
1	B	309	SER	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	88	HIS	2.6
1	B	278	VAL	2.6
1	B	160	ALA	2.6
1	B	45	LEU	2.6
1	A	90	THR	2.5
1	A	343	ALA	2.5
1	A	163	HIS	2.5
1	B	352	LEU	2.5
1	A	332	ILE	2.5
1	B	289	LEU	2.4
1	A	249	PRO	2.4
1	A	77	LEU	2.4
1	B	393	GLU	2.4
1	A	336	GLY	2.4
1	B	343	ALA	2.3
1	A	337	THR	2.3
1	B	350	GLU	2.3
1	A	228	GLN	2.3
1	A	25	HIS	2.2
1	B	241	LEU	2.2
1	A	313	LEU	2.2
1	A	286	ARG	2.2
1	A	187	PHE	2.2
1	B	275	GLU	2.2
1	B	310	GLY	2.2
1	A	345	VAL	2.1
1	A	347	ARG	2.1
1	A	21	ASP	2.1
1	B	396	TRP	2.1
1	B	226	ALA	2.1
1	B	16	PRO	2.1
1	B	274	THR	2.1
1	B	391	ILE	2.1
1	A	155	ALA	2.1
1	A	287	LYS	2.1
1	A	260	LEU	2.1
1	B	50	VAL	2.1
1	B	288	VAL	2.1
1	B	331	ASP	2.1
1	A	304	PHE	2.1
1	B	317	PHE	2.1
1	A	354	ILE	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	270	LEU	2.0
1	B	346	VAL	2.0
1	A	47	PHE	2.0
1	B	151	MET	2.0
1	B	91	GLY	2.0
1	B	212	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	FMN	A	504	31/31	0.87	0.17	34,37,52,52	0
4	FMN	B	504	31/31	0.88	0.14	36,38,52,53	0
3	MN	A	503	1/1	0.92	0.07	49,49,49,49	0
2	C2E	A	501	46/46	0.92	0.11	36,39,44,47	0
2	C2E	B	501	46/46	0.92	0.11	36,40,43,46	0
3	MN	B	503	1/1	0.94	0.14	53,53,53,53	0
3	MN	A	502	1/1	0.96	0.05	43,43,43,43	0
3	MN	B	502	1/1	0.99	0.03	41,41,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.