



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2021 – 03:29 PM EDT

PDB ID : 3GQB  
Title : Crystal Structure of the A3B3 complex from V-ATPase  
Authors : Meher, M.; Akimoto, S.; Iwata, M.; Nagata, K.; Hori, Y.; Yoshida, M.;  
Yokoyama, S.; Iwata, S.; Yokoyama, K.  
Deposited on : 2009-03-24  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	578	 3% 52% 41% 7%
1	C	578	 3% 49% 44% 7%
2	B	464	 2% 56% 37% 6% .
2	D	464	 4% 59% 35% 5% .

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 16863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type ATP synthase alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	578	4477	2856	763	839	19	0	0	0
1	C	578	4477	2856	763	839	19	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	SER	CYS	engineered mutation	UNP Q56403
A	235	SER	THR	engineered mutation	UNP Q56403
A	255	SER	CYS	engineered mutation	UNP Q56403
A	507	SER	CYS	engineered mutation	UNP Q56403
C	28	SER	CYS	engineered mutation	UNP Q56403
C	235	SER	THR	engineered mutation	UNP Q56403
C	255	SER	CYS	engineered mutation	UNP Q56403
C	507	SER	CYS	engineered mutation	UNP Q56403

- Molecule 2 is a protein called V-type ATP synthase beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	460	3601	2280	623	689	9	0	0	0
2	D	460	3601	2280	623	689	9	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	264	SER	CYS	engineered mutation	UNP Q56404
D	264	SER	CYS	engineered mutation	UNP Q56404


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	199	Total 199	O 199	0	0
3	B	135	Total 135	O 135	0	0
3	C	239	Total 239	O 239	0	0
3	D	134	Total 134	O 134	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: V-type ATP synthase alpha chain




- Molecule 1: V-type ATP synthase alpha chain







• Molecule 2: V-type ATP synthase beta chain



• Molecule 2: V-type ATP synthase beta chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	199.37Å 199.37Å 179.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.95 – 2.80 19.94 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.6 (19.95-2.80) 97.7 (19.94-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 2.79Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.250 , 0.281 0.263 , 0.293	Depositor DCC
$R_{free}$ test set	4941 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.7	Xtriage
Anisotropy	0.005	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 8.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.077 for -h,-k,l	Xtriage
Reported twinning fraction	0.845 for 1.000H, 1.000K, L 0.155 for -1.000H, -1.000K, L	Depositor
Outliers	0 of 98400 reflections	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	16863	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.56	0/4573	0.71	1/6203 (0.0%)
1	C	0.58	0/4573	0.72	2/6203 (0.0%)
2	B	0.54	0/3668	0.67	0/4965
2	D	0.50	0/3668	0.65	0/4965
All	All	0.55	0/16482	0.69	3/22336 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	241	LEU	CA-CB-CG	5.47	127.89	115.30
1	A	241	LEU	CA-CB-CG	5.42	127.76	115.30
1	C	215	LEU	CA-CB-CG	-5.13	103.49	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4477	0	4494	367	0
1	C	4477	0	4494	374	1
2	B	3601	0	3627	193	1
2	D	3601	0	3627	197	0
3	A	199	0	0	120	1
3	B	135	0	0	60	1

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	239	0	0	135	0
3	D	134	0	0	70	0
All	All	16863	0	16242	1107	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 1107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:GLY:O	1:A:329:ARG:HD2	1.24	1.32
1:A:574:PHE:HB3	3:A:697:HOH:O	1.33	1.29
1:A:447:TYR:HA	3:A:651:HOH:O	1.17	1.27
1:C:124:LYS:HD3	3:C:714:HOH:O	1.25	1.26
1:C:256:GLY:O	1:C:329:ARG:HD2	1.22	1.25

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:410:ARG:NH2	1:C:478:ASP:OD2[6_555]	2.13	0.07
3:A:653:HOH:O	3:B:704:HOH:O[2_665]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	576/578 (100%)	512 (89%)	51 (9%)	13 (2%)	6	21
1	C	576/578 (100%)	504 (88%)	61 (11%)	11 (2%)	8	26
2	B	458/464 (99%)	407 (89%)	43 (9%)	8 (2%)	9	29

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	458/464 (99%)	408 (89%)	41 (9%)	9 (2%)	7	24
All	All	2068/2084 (99%)	1831 (88%)	196 (10%)	41 (2%)	7	24

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	GLY
1	A	235	SER
1	A	555	SER
2	B	281	ARG
2	B	346	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	468/468 (100%)	425 (91%)	43 (9%)	9	27
1	C	468/468 (100%)	419 (90%)	49 (10%)	7	20
2	B	386/390 (99%)	352 (91%)	34 (9%)	10	29
2	D	386/390 (99%)	354 (92%)	32 (8%)	11	32
All	All	1708/1716 (100%)	1550 (91%)	158 (9%)	9	26

5 of 158 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	432	THR
2	D	202	SER
1	C	461	GLU
2	D	78	ASP
2	D	336	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	496	GLN
2	D	142	ASN
2	D	129	GLN
2	D	198	GLN
2	B	198	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	578/578 (100%)	0.27	16 (2%) 53 43	9, 29, 57, 63	0
1	C	578/578 (100%)	0.31	15 (2%) 56 46	9, 29, 57, 63	0
2	B	460/464 (99%)	0.19	10 (2%) 62 52	5, 28, 54, 67	0
2	D	460/464 (99%)	0.26	20 (4%) 35 25	5, 28, 54, 67	0
All	All	2076/2084 (99%)	0.26	61 (2%) 51 41	5, 29, 56, 67	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	464	GLY	5.5
2	B	401	GLU	5.1
2	D	395	LEU	5.1
2	D	400	GLY	4.9
1	C	388	GLY	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.